CHAOS, PERIODICITY, AND SNAKELIKE CONTINUA

BY

MARCY BARGE AND JOE MARTIN¹

ABSTRACT. The results of this paper relate the dynamics of a continuous map f of the interval and the topology of the inverse limit space with bonding map f . These inverse limit spaces have been studied by many authors, and are examples of what Bing has called "snakelike continua". Roughly speaking, we show that when the dynamics of f are complicated, the inverse limit space contains indecomposable subcontinua. We also establish a partial converse.

Introduction. Let I be a closed interval, and let $f: I \rightarrow I$ be a continuous function. Associated with f is the inverse limit space $(I, f) = \{(x_0, x_1, \dots) | f(x_{n+1}) = x_n\}$. With a natural topology, (I, f) is a compact, connected, metric space, and is an example of what Bing [Bi] has called a snakelike continuum. In this paper we will investigate the relationship between behavior of the orbits $\{f^n(x) | n \geq 0\}$ of points of I under f , and the topological properties of the space (I, f) . These examples suggest some of the ideas which we will explore.

EXAMPLE 1. Let $I = [0, 1]$, and define $f: I \rightarrow I$ by

$$f(t) = \begin{cases} \frac{3}{2}t & \text{if } 0 \leq t \leq \frac{2}{3}, \\ \frac{5}{3} - t & \text{if } \frac{2}{3} \leq t \leq 1. \end{cases}$$

It can be verified that (i) if $x \in I$, then $\{f^n(x) | n \geq 0\}$ is not dense in I , and (ii) f has points of period 1 and 2, but no points of period $n \geq 3$. Now let X be the subspace of the plane defined by

$$X = \{(x, y) | 0 < x \leq 1 \text{ and } y = \sin(1/x)\} \cup \{(x, y) | x = 0 \text{ and } -1 \leq y \leq 1\}.$$

Received by the editors December 16, 1983 and, in revised form, July 18, 1984.

1980 *Mathematics Subject Classification.* Primary 58F13, 58F22, 55F20, 54H20.

¹Research supported in part by NSF grant 8120790.

©1985 American Mathematical Society
0002-9947/85 \$1.00 + \$.25 per page

It can be verified that X is homeomorphic with (I, f) and has the following property: if C is a nondegenerate subcontinuum of X , then C is the union of two of its proper subcontinua.

EXAMPLE 2. Let $I = [0, 1]$, and let $f: I \rightarrow I$ be defined by

$$f(t) = \begin{cases} 2t & \text{if } 0 \leq t \leq \frac{1}{2}, \\ 2 - 2t & \text{if } \frac{1}{2} \leq t \leq 1. \end{cases}$$

It can be verified that (i) there is a point $x \in I$ such that $\{f^n(x) | n \geq 0\}$ is dense in I , and (ii) for each positive integer k , f has points of period k . Now let X be the subspace of the plane described as follows: let C be the standard Cantor set on the x -axis. Let K_0 be the union of all semicircles in the upper half-plane with endpoints on C , which are symmetric with the line $x = \frac{1}{2}$. For each positive integer i , let K_i be the union of all semicircles in the lower half-plane with endpoints on C , which are symmetric about the line $x = 5/3^i \cdot 2$. Then X is $\bigcup_{i=0}^{\infty} K_i$.

It can be verified that X is homeomorphic with (I, f) and has the following property: X is not the union of two of its proper subcontinua.

Definitions and terminology. If a and b are distinct real numbers we will let $[a, b]$ denote the smallest closed interval containing both a and b , and let (a, b) denote the associated open interval. We will generically let I be a closed interval and will be considering continuous functions $f: I \rightarrow I$. All of the functions which we will consider are continuous.

If $f: I \rightarrow I$ and $x \in I$, then the *orbit* of x under f is $\{y | \text{for some integer } n, n \geq 0, y = f^n(x)\}$. We will be interested in functions $f: I \rightarrow I$ for which there is a point x whose orbit is dense in I . (In [AY] a function f is defined to be chaotic if there is a point whose orbit is dense and if every point is unstable. As a corollary to Lemma 2 we show that, for functions on the interval, the existence of a dense orbit implies that every point is unstable.)

If $f: I \rightarrow I$ and $x \in I$, the statement that x has *period* n , means that n is a positive integer, $f^n(x) = x$, and if k is an integer, $1 \leq k < n$, then $f^k(x) \neq x$.

Associated with $f: I \rightarrow I$ is the compact, connected metric space $(I, f) = \{(x_0, x_1, \dots) | f(x_i) = x_{i-1}\}$ with metric

$$d((x_0, x_1, \dots), (y_0, y_1, \dots)) = \sum_{i=0}^{\infty} \frac{|x_i - y_i|}{2^i}.$$

(I, f) is an example of what Bing [Bi] has called a *snakelike continuum*. The reason for this terminology is that for each $\varepsilon > 0$, there is a finite open covering $\{g_1, g_2, \dots, g_n\}$ of (I, f) such that (i) $\text{diam } g_i < \varepsilon$, and (ii) $g_i \cap g_j \neq \emptyset$ iff $j = i - 1$, i , or $i + 1$. We will denote elements of (I, f) by subbarred letters, as $\underline{x} = (x_0, x_1, \dots)$. The projection maps π_n of (I, f) onto I given by $\pi_n(\underline{x}) = x_n$ are continuous. If H is a subcontinuum (compact, connected subspace) of (I, f) we will let H_n denote $\pi_n(H)$. Note that H_n is a closed interval or point, and that $f(H_{n+1}) = H_n$.

If $f: I \rightarrow I$, then f induces a homeomorphism $\hat{f}: (I, f) \rightarrow (I, f)$ by $\hat{f}((x_0, x_1, \dots)) = (f(x_0), x_0, x_1, \dots)$. Notice that $f \circ \pi_n = \pi_n \circ \hat{f}$, $\pi_n = \pi_{n+1} \circ \hat{f}$, and $f \circ \pi_{n+1} = \pi_n$.

Here are important facts about snakelike continua which we will utilize. Suppose S is snakelike; then (i) the intersection of any collection of subcontinua of S is a subcontinuum of S , and (ii) if H is a subcontinuum of S , then $S - H$ has at most two components (see [Bi]).

If S is a continuum, the statement that S is *indecomposable* means that S is not the union of two of its proper subcontinua. Here are two conditions each of which is equivalent to the indecomposability of S (see [HY, pp. 139–141]).

(1) If H is a subcontinuum of S , then H contains no open set in S .

(2) S contains three distinct points x , y and z such that S is irreducible between each pair of these points. (*Irreducibility* between x and y means that no proper subcontinuum of S contains both x and y .)

We will utilize the following important construction due to Bing [Bi]. Suppose that (I, f) contains no indecomposable subcontinuum with interior. For each $\underline{x} \in (I, f)$ let $g_{\underline{x}}$ be the intersection of all subcontinua of (I, f) that contain interiorly a subcontinuum that contains $\underline{x}$ in its interior. Then $g_{\underline{x}}$ is a subcontinuum of (I, f) . Furthermore, the sets $g_{\underline{x}}$ partition (I, f) and if we let $G = \{g_{\underline{x}} | \underline{x} \in (I, f)\}$ with the quotient topology, then G is an arc (i.e. homeomorphic with I). Moreover, f induces a homeomorphism $\hat{f}$ of G onto G given by $\hat{f}(g_{\underline{x}}) = g_{\hat{f}(\underline{x})}$. Bing also shows that, for each $\underline{x} \in (I, f)$, $g_{\underline{x}}$ does not have interior.

If A is a set, we will let both $\bar{A}$ and $\text{cl}(A)$ denote the closure of A and let $\text{int } A$ denote the interior of A .

THEOREM 1. *Suppose that k and n are integers, $k \geq 0$, $n \geq 1$, and that f has a point of period $2^k(2n + 1)$; i.e., of period not a power of 2. Then (I, f) has an indecomposable subcontinuum that is invariant under $\hat{f}^{2^{k+1}}$.*

PROOF. It follows from Sarkovskii's Theorem [S, N] and the hypothesis that f has a point of period $(2^{k+1})(3)$. Then $f^{2^{k+1}}$ has a point of period 3. Now, for each positive integer j , the spaces (I, f) and (I, f^j) are homeomorphic. One such homeomorphism is $(x_0, x_1, \dots, x_j, \dots) \rightarrow (x_0, x_j, x_{2j}, \dots)$. Using this fact, we need only show that if $g: I \rightarrow I$ is continuous and g has a point of period 3, then (I, g) has an indecomposable subcontinuum that is invariant under $\hat{g}$. Let g be such a function and let x be a point of period 3. Then we have the point $\underline{x} = (x, g^2(x), g(x), x, \dots)$ of (I, g) with $\hat{g}^3(\underline{x}) = \underline{x}$ and $\hat{g}(\underline{x}) \neq \underline{x}$.

Now let S be the intersection of all subcontinua of (I, g) which contain $\{\underline{x}, \hat{g}(\underline{x}), \hat{g}^2(\underline{x})\}$. Since $\{\underline{x}, \hat{g}(\underline{x}), \hat{g}^2(\underline{x})\}$ is contained in both $\hat{g}(S)$ and $\hat{g}^{-1}(S)$, it follows that $S = g(S)$. So S is a subcontinuum of (I, g) which is invariant under $\hat{g}$.

We will next show that S is irreducible between each pair of $\underline{x}$, $\hat{g}(\underline{x})$ and $\hat{g}^2(\underline{x})$. It will follow that S is indecomposable. Suppose, for example, that S is reducible between $\hat{g}(\underline{x})$ and $\hat{g}^2(\underline{x})$. Then there is a proper subcontinuum H of S which contains $\hat{g}(\underline{x})$ and $\hat{g}^2(\underline{x})$. Now, because H is proper, $\underline{x} \notin H$, and it follows that there is an integer N such that if $n > N$, then $\pi_n(\underline{x}) \notin H_n$. However, for every third n , $\pi_n(\underline{x})$ is between $\pi_n(\hat{g}(\underline{x}))$ and $\pi_n(\hat{g}^2(\underline{x}))$, which are in the closed interval H_n . This contradiction establishes Theorem 1.

In what follows, if $x \in I$, and s and k are integers, $s \geq 1$, $k \geq 0$, we let $A_{s,k}(x) = A_{s,k}$ be $\{f^{sn+k}(x) | n \geq 0\}$.

LEMMA 2. *Suppose that f has a dense orbit, and that x is a point whose orbit is dense, i.e. $A_{1,0}$ is dense in I . Then one of the following occurs.*

- (i) $A_{2,0}$ is dense in I , in which case $A_{s,k}$ is dense in I for each $s \geq 1$, $k \geq 0$, or
- (ii) $A_{2,0}$ is not dense in I , in which case $I = \overline{A_{2,0}} \cup \overline{A_{2,1}}$, $\overline{A_{2,0}}$ and $\overline{A_{2,1}}$ are closed intervals which intersect in a point, and $f(\overline{A_{2,0}}) = \overline{A_{2,1}}$, $f(\overline{A_{2,1}}) = \overline{A_{2,0}}$. Moreover, for each $k \geq 1$, $A_{2k,0}$ is dense in $\overline{A_{2,0}}$ and $A_{2k,1}$ is dense in $\overline{A_{2,1}}$.

PROOF. Let s be an integer, $s \geq 1$, and for each integer r , $0 \leq r \leq s-1$, let $B_r = \overline{A_{s,r}}$. Then since $\bigcup_{r=0}^{s-1} A_{s,r} = A_{1,0}$, it follows that $\bigcup_{r=0}^{s-1} B_r = I$. From this we see that there is an r , $0 \leq r \leq s-1$, so that B_r has nonempty interior.

Next, notice that if J is a closed subinterval of I , then $f(J)$ is a closed interval, because if $f(J)$ is a point, then for some integer n , $f^n(x)$ is periodic. From this remark, and the fact that $f(B_r) \subset B_{r+1} \pmod{s}$ it follows that, for each integer i , $0 \leq i \leq r-1$, B_i has nonempty interior.

We next show that if the interiors of B_i and B_j intersect, then these interiors are identical. For, if $\text{int } B_i \cap \text{int } B_j \neq \emptyset$, then there is a positive integer n so that $f^{2n+i}(x) \in \text{int } B_i \cap \text{int } B_j$, and there is a sequence $n_1, n_2, n_3, \dots$ of positive integers such that $f^{n_k s + j}(x) \rightarrow f^{s n + i}(x)$. Then for every integer $l > 0$ we have $f^{s(n_k + l) + j}(x) \rightarrow f^{s(n + l) + i}(x)$. From this it follows that $\text{cl}\{f^{s n + i}(x), f^{s(n+1) + i}(x), \dots\} \subset B_j$ and hence that

$$B_i \subset B_j \cup \{f^i(x), f^{s+i}(x), \dots, f^{(n-1)s+i}(x)\}.$$

From this we see that $\text{int } B_i \subset \text{int } B_j$. A similar argument shows that $\text{int } B_j \subset \text{int } B_i$, and hence $\text{int } B_i = \text{int } B_j$.

Now let $G = \{g | \text{for some } r, 0 \leq r \leq s-1, g \text{ is a component of } \text{int } B_r\}$. Notice that G is a collection of disjoint open intervals whose union is dense in I . Since G is countable we list G as $\{g_1, g_2, \dots\}$. Now for each $g_i \in G$ let $C_i = \overline{g_i}$. Then C_i is a closed interval, $f(C_i)$ is a closed interval by an earlier remark, and there is an r , $0 \leq r \leq s-1$, such that $f(C_i) \subset B_r$. Then $\text{int } f(C_i) \subset \text{int } B_r$ and there is an integer k so that $f(C_i) \subset C_k$. Because x has a dense orbit, we see that if i and k are integers which are subscripts of elements of G , then there is a positive integer j so that $f^j(C_i) \subset C_k$. Since we have this transitivity, it follows that G is finite. Thus we may

list G as $\{g_1, g_2, \dots, g_n\}$ and their closures as $C_1, C_2, \dots, C_n$. Notice that because of the above transitivity condition, the set $\{C_1, C_2, \dots, C_n\}$ is permuted by f .

We next show that $n \leq 2$. Let y be a fixed point of f . Now, if $y \in \text{int } C_i$, then $f(C_i) = C_i$, which is impossible unless $n = 1$. Similarly, if y is an endpoint of I , then $n = 1$. If y is a common endpoint of C_i and C_j , then $f(C_i) = C_j$ and $f(C_j) = C_i$ which is impossible unless $n = 2$. Notice that the integer n depends on s . In what follows we will refer to n as $n(s)$.

We now verify the conclusion. First, assume that $A_{2,0}$ is dense in I . Let s be an integer, $s \geq 1$, and suppose that $n(s) = 2$. Then, there are closed intervals C_1 and C_2 with $C_1 \cup C_2 = I$, $C_1 \cap C_2 = \{\text{pt}\}$, $f(C_1) = C_2$ and $f(C_2) = C_1$. Assuming that $x \in C_2$ we see that, for each j , $f^{2j}(x) \in C_2$ and hence $A_{2,0} \cap \text{int } C_1 \neq \emptyset$. This contradicts the fact that $A_{2,0}$ is dense in I , and hence $n(s) = 1$. Then, for each r , $0 \leq r \leq s - 1$, $B_r = I$. Then $\bar{A}_{s,r} = I$, and so $A_{s,r}$ is dense in I . From this, we see that for any integer $k \geq 0$, $A_{s,k}$ is dense in I .

Next, assume that $A_{2,0}$ is not dense in I . Let $s = 2$. Since $A_{2,0}$ is not dense, $B_0 \neq I$ and so $n(2) = 2$. Now let j be an integer, $j \geq 1$. Then, for each integer l , $A_{2j,l} \subset A_{2,l}$ and since $\bar{A}_{2,0} \neq I$, we have $n(2j) = 2$. Now, notice that the intervals C_1 and C_2 which we construct for $s = 2j$ are independent of j . This is because their common endpoint is the only fixed point for the function f . Then, assuming $x \in C_2$, we have $C_2 = \bar{A}_{2,0}$, $C_1 = \bar{A}_{2,1}$, and, for each integer $k \geq 1$, $\bar{A}_{2k,0} = C_2$ and $\bar{A}_{2k,1} = C_1$. This establishes Lemma 2.

The following result is known [N], but we include it for completeness.

COROLLARY. *Suppose that f has a dense orbit. Then the set of periodic points of f is dense in I .*

PROOF. Let V be an open interval in I . Let x be a point of V whose orbit is dense in I . If $\{f^{2n}(x) | n \geq 0\}$ is not dense in I , we may assume from Lemma 1 that $V \subset \text{cl}\{f^{2n}(x) | n \geq 0\}$. Let j be an integer such that $f^j(x) \in V$. We may assume that $x < f^j(x)$. Let $g: I \rightarrow I$ be the function $g = f^j$. From Lemma 2, it follows that $\{g^k(x) | k \geq 0\}$ is dense in V . Now let l be the smallest positive integer such that $g^l(g(x)) < g(x)$. Then $g^l(x) = g^{l-1}(g(x)) \geq g(x) > x$ and $g^l(g(x)) < g(x)$. So $g^l(x) > x$ and $g^l(g(x)) < g(x)$. Consequently g^l has a fixed point y , $x < y < g(x)$. Since $g^l(y) = y$, $f^{kl}(y) = y$ and, since $y \in V$, V contains a periodic point of f .

DEFINITION If $y \in I$, the statement that y is *topologically stable* means that if $\varepsilon > 0$, then there is a $\delta > 0$ such that if $z \in I$ and $|y - z| < \delta$ then for each positive integer n , $|f^n(y) - f^n(z)| < \varepsilon$. If y is not topologically stable, then y is called *topologically unstable*.

COROLLARY. *Suppose that f has a dense orbit. Then every point of I is topologically unstable.*

PROOF. Suppose that $y \in I$ and that y is topologically stable. Let x be a point of I whose orbit under f is dense. We first show that the orbit of y is dense. Suppose that U is an open interval in I and, for each n , $f^n(y) \notin U$. Let V be the open interval which is the open middle third of U . Let $\varepsilon = \frac{1}{3} \text{diam } U$. Then, since y is topologically

stable, there is a $\delta > 0$ such that if $|z - y| < \delta$ then, for each n , $|f^n(y) - f^n(z)| < \varepsilon$. In particular, if $|z - y| < \delta$ then, for each n , $f^n(z) \notin V$. Now since x has a dense orbit, there is a j such that $|f^j(x) - y| < \delta$. Then there is an integer $k > j$ such that $f^k(x) \in V$. But then $f^{k-j}(f^j(x)) \in V$ and this is a contradiction. Therefore, the orbit of y is dense.

Now it follows from Lemma 2 that there is a positive number ε and a subinterval C of I such that $\text{diam } C > 3\varepsilon$, and, for each positive integer n , $\{f^{kn}(y) | k \geq 0\}$ is dense in C . Now choose δ such that if $|z - y| < \delta$, then for each j , $|f^j(z) - f^j(y)| < \varepsilon$. Using the previous corollary, let t be a periodic point such that $|t - y| < \delta$. Let n be the period of t . Then, for each k , $|f^{kn}(t) - f^{kn}(y)| < \varepsilon$, so $|t - f^{kn}(y)| < \varepsilon$. But then $\{f^{kn}(y) | k \geq 0\}$ is not dense in C . This establishes the Corollary.

EXAMPLE 3. Let $I = [0, 1]$, and let $f: I \rightarrow I$ be defined by

$$f(x) = \begin{cases} 2x + \frac{1}{2}, & 0 \leq x \leq \frac{1}{4}, \\ -2x + \frac{3}{2}, & \frac{1}{4} \leq x \leq \frac{3}{4}, \\ 2x - \frac{3}{2}, & \frac{3}{4} \leq x \leq 1. \end{cases}$$

Then f has the following properties. (i) There is a number x such that $\{f^n(x) | n \geq 0\}$ is dense in I , and (ii) if $y \in I$, then $\{f^{2n}(y) | n \geq 0\}$ is not dense in I .

It can be shown that (I, f) is homeomorphic with Example 2 together with its reflection through the origin.

THEOREM 3. Suppose that $x \in I$, and that x has a dense orbit under f . Then one of the following occurs.

- (a) $\{f^{2n}(x) | n \geq 0\}$ is dense in I , in which case (I, f) is indecomposable, or
- (b) $\{f^{2n}(x) | n \geq 0\}$ is not dense in I , in which case there are proper subcontinua H and K of (I, f) such that (i) H and K are indecomposable, (ii) $H \cup K = (I, f)$, (iii) $H \cap K$ is a point, (iv) $\hat{f}(H) = K$, and (v) $\hat{f}(K) = H$.

PROOF. First, assume that $\{f^{2n}(x) | n \geq 0\}$ is dense in I . Assume further that (I, f) has no indecomposable subcontinuum with interior. Then Bing's construction [Bi] applies to yield a homeomorphism $\hat{f}: G \rightarrow G$ of the arc G onto itself. Since x has a dense orbit, the function f is onto, and by choosing inverse images we may construct the point $\underline{x} = (x, f^{-1}(x), \dots)$ of (I, f) . It is clear that $\{\hat{f}^n(x) | n \geq 0\}$ is dense in (I, f) . From this it follows that $\{\hat{f}^n(g_{\underline{x}}) | n \geq 0\}$ is dense in G . This is impossible since $\hat{f}$ is a homeomorphism and G is an arc. Therefore there is a subcontinuum S of (I, f) such that S is indecomposable and has interior.

We will next show that $S = (I, f)$. First, suppose that for each positive integer k , $\text{int}(\hat{f}^k(S)) \cap \text{int } S = \emptyset$. As above, let $\underline{x} = (x, f^{-1}(x), f^{-2}(x), \dots)$. The previous assumption makes it impossible for $\{\hat{f}^n(\underline{x}) | n \geq 0\}$ to be dense in (I, f) . It follows that there is a positive integer k such that $\text{int}(\hat{f}^k(S)) \cap \text{int } S \neq \emptyset$. Then $\hat{f}^k(S) \cap S$ is a subcontinuum of both $\hat{f}^k(S)$ and S which has interior, and, since S is indecomposable, $\hat{f}^k(S) = S$. Then, for each positive integer j , $\hat{f}^{jk}(S) = S$. Now let l be a positive integer such that $\hat{f}^l(\underline{x}) \in S$. From Lemma 2 we have $\{f^{j(kl)}(x) | j \geq 1\}$ is dense in I and hence $\{\hat{f}^{j(kl)}(\underline{x}) | j \geq 1\}$ is dense in (I, f) . Since $\hat{f}^{j(kl)}(\underline{x}) \in S$ we see that $S = (I, f)$.

Next, we consider the case where $\{f^{2n}(x) | n \geq 0\}$ is not dense in I . By Lemma 2, there are closed subintervals C_1 and C_2 such that $I = C_1 \cup C_2$, $C_1 \cap C_2 = \{p\}$, $f(C_1) = C_2$ and $f(C_2) = C_1$. Now let

$$H = \{\underline{y} | \underline{y} \in (I, f), y_{2n} \in C_1 \text{ and } y_{2n+1} \in C_2 \text{ if } n \geq 0\}$$

and

$$K = \{\underline{y} | \underline{y} \in (I, f), y_{2n+1} \in C_1 \text{ and } y_{2n} \in C_2 \text{ if } n \geq 0\}.$$

Then H and K are subcontinua of (I, f) , $H \cup K = (I, f)$, $H \cap K = (p, p, p, \dots)$, $\hat{f}(H) = K$ and $\hat{f}(K) = H$. In order to see that K is indecomposable, consider the function $h = f^2: C_2 \rightarrow C_2$. Then, assuming that $x \in C_2$, it follows from Lemma 2 that both $\{h^n(x) | n \geq 0\}$ and $\{h^{2n}(x) | n \geq 0\}$ are dense in C_2 . By the first part of this theorem, (C_2, h) is indecomposable. The correspondence

$$(y, h^{-1}(y), h^{-2}(y), \dots) \leftrightarrow (y, f(h^{-1}(y)), h^{-1}(y), f(h^{-2}(y)), h^{-2}(y), \dots)$$

is a homeomorphism between (C_2, h) and K . Therefore K is indecomposable and, as $H = \hat{f}(K)$, H is indecomposable. This establishes Theorem 3.

DEFINITION. Suppose that y is a fixed point of f . This statement that x is *homoclinic to the fixed point* y means that $x \neq y$ and there is a choice of inverse images $f^{-1}(x), f^{-2}(x), \dots$ such that both $f^n(x) \rightarrow y$ and $f^{-n}(x) \rightarrow y$. If y is a periodic point of f with period s , then the statement that x is *homoclinic to* y means that x is homoclinic to the fixed point y under f^s .

The next result can be obtained from Theorem 1 and [BI]. We include a direct proof.

THEOREM 4. *If f has a point homoclinic to a periodic point, then (I, f) contains an indecomposable subcontinuum.*

PROOF. Since, for each positive integer s , (I, f) is homeomorphic with (I, f^s) , we will assume that f has a point homoclinic to a fixed point. Let y be a fixed point and let $x \neq y$, together with a choice of inverse images, be such that $f^n(x) \rightarrow y$ and $f^{-n}(x) \rightarrow y$. In (I, f) let $\underline{y} = (y, y, y, \dots)$ and $\underline{x} = (x, f^{-1}(x), f^{-2}(x), \dots)$. Then $\hat{f}(\underline{y}) = \underline{y}$, $\hat{f}^n(\underline{x}) \rightarrow \underline{y}$ and $\hat{f}^{-n}(\underline{x}) \rightarrow \underline{y}$.

Now let S be the intersection of all subcontinua of (I, f) which contain $\{\underline{y}\} \cup \{\hat{f}^n(\underline{x}) | -\infty < n < \infty\}$. Since both $\hat{f}(S)$ and $\hat{f}^{-1}(S)$ contain $\{\underline{y}\} \cup \{\hat{f}^n(\underline{x}) | -\infty < n < \infty\}$, we see that $\hat{f}(S) = S$.

Now, for each $n \geq 0$, $\pi_n \circ \hat{f} = f \circ \pi_n$, and so $f(\pi_n(S)) = \pi_n(\hat{f}(S)) = \pi_n(S)$. Thus $\pi_n(S)$ is invariant under f . Now let $J = \pi_0(S)$. Then $f(J) = J$ and $S = (J, f)$.

Suppose now that $S = (J, f)$ contains no indecomposable subcontinuum with interior. Then Bing's construction [Bi] yields an arc G , and $\hat{f}: G \rightarrow G$ is a homeomorphism. Now $\hat{f}^n(g_x) \rightarrow g_y$ and $\hat{f}^{-n}(g_x) \rightarrow g_y$. Since G is an arc and $\hat{f}$ is a homeomorphism, this is impossible unless, for each j , $\hat{f}^j(g_x) = g_y$. This implies that, for each j , $g_{\hat{f}^j(x)} = g_y$. But then g_y contains $\{y\} \cup \{\hat{f}^j(x) | -\infty < j < \infty\}$ and hence $g_y = S$. But then G is degenerate, and this is impossible.

Therefore S contains an indecomposable subcontinuum with interior, and (I, f) contains an indecomposable subcontinuum.

DEFINITION. If $f: I \rightarrow I$ is continuous, then the statement that f is *organic* means that if $\underline{x} \in (I, f)$, $\underline{y} \in (I, f)$ and (I, f) is irreducible from $\underline{x}$ to $\underline{y}$, then there is a positive integer n such that $f^n([\pi_n(\underline{x}), \pi_n(\underline{y})]) = I$. The statement that f is *inorganic* means that f is not organic.

EXAMPLE 4. The accompanying figure is a sketch of a function which is inorganic.

In [H], Henderson shows that (I, f) is a pseudo-arc, a particular snakelike continuum which is hereditarily indecomposable. Notice that f has no points of period greater than one.

LEMMA 5. Let $I = [a, b]$ and suppose that $f: I \rightarrow I$ is continuous and onto. Further, suppose that (I, f) is irreducible between $\underline{x} = (x_0, x_1, \dots)$ and $\underline{y} = (y_0, y_1, \dots)$. Then if c and d are numbers, $a < c < d < b$, then there is an integer N such that if $n > N$, then $[c, d] \subset f^n([x_n, y_n])$.

PROOF. Recall that $[x_n, y_n]$ is the smallest closed interval containing x_n and y_n .

First, notice that if $n_2 > n_1$, then $f^{n_1}([x_{n_1}, y_{n_1}]) \subset f^{n_2}([x_{n_2}, y_{n_2}])$. This is because $f^{n_2-n_1}(x_{n_2}) = x_{n_1}$ and $f^{n_2-n_1}(y_{n_2}) = y_{n_1}$. Then we have $[x_0, y_0] \subset f([x_1, y_1]) \subset f^2([x_2, y_2]) \subset \dots$. Now if k is an integer, $k \geq 0$, let J_k be $\text{cl}(\bigcup_{n \geq k} f^{n-k}([x_n, y_n]))$. Then for each k , J_k is a closed subinterval of I , and $f(J_{k+1}) = J_k$.

Now let J be the subcontinuum of (I, f) defined by $J = \{(z_0, z_1, z_2, \dots) | z_k \in J_k \text{ and } f(z_{k+1}) = z_k\}$. Since $\underline{x}$ and $\underline{y}$ belong to J , and (I, f) is irreducible from $\underline{x}$ to $\underline{y}$, it follows that $J = (I, f)$. Now, since f is onto, $J_0 = I$. Then $I = \text{cl}(\bigcup_{n=0}^{\infty} f^n([x_n, y_n]))$, and since $f^n([x_n, y_n]) \subset f^{n+1}([x_{n+1}, y_{n+1}])$, the conclusion follows.

LEMMA 6. Suppose that $I = [a, b]$ and that $f: I \rightarrow I$ is continuous. If there are numbers p and q , $a < p < b$, $a < q < b$, and integers r and s such that $f^r(p) = a$, $f^s(q) = b$, then f is organic.

PROOF. Suppose that (I, f) is irreducible between $\underline{x} = (x_0, x_1, \dots)$ and $\underline{y} = (y_0, y_1, \dots)$. It follows from the argument given in Lemma 5 that there is an integer

N_r such that if $n > N_r$, then $p \in f^{n-r}([x_n, y_n])$, and an integer N_s such that if $n > N_s$, then $q \in f^{n-s}([x_n, y_n])$. Then, if $n > N_r$, $a \in f^n([x_n, y_n])$, and if $n > N_s$, then $b \in f^n([x_n, y_n])$.

Now if $n > N_r + N_s$, then $I = f^n([x_n, y_n])$, and so f is organic.

THEOREM 7. *If $f: I \rightarrow I$ is organic, and (I, f) is indecomposable, then f has a periodic point whose period is not a power of 2.*

PROOF. Since (I, f) is indecomposable, there are three points $\underline{x} = (x_0, x_1, \dots)$, $\underline{y} = (y_0, y_1, \dots)$ and $\underline{z} = (z_0, z_1, \dots)$ in (I, f) , such that (I, f) is irreducible between any two of them. Because f is organic, there is a positive integer n such that $f^n([x_n, y_n]) = f^n([x_n, z_n]) = f^n([y_n, z_n]) = I$. We will assume that the notation is chosen so that $x_n < y_n < z_n$. Now since $[x_n, y_n] \subset f^n([y_n, z_n])$, there is a closed subinterval J_1 of $[y_n, z_n]$ such that $f^n(J_1) = [x_n, y_n]$. Now there is a closed subinterval J_2 of $[y_n, z_n]$ such that $f^n(J_2) = J_1$. Notice that $y_n \notin J_1 \cap J_2$. Finally, let J_3 be a closed subinterval of $[x_n, y_n]$ such that $f^n(J_3) = J_2$.

Now $J_3 \subset f^{3n}(J_3)$, and so there is a point p of J_3 such that $f^{3n}(p) = p$.

Now suppose $f^n(p) = p$. Then $p = y_n$ and $f^n(y_n) = f^{2n}(y_n) = f^{3n}(y_n) = y_n$. But then $y_n \in J_1 \cap J_2$, which is a contradiction. Thus the points p , $f^n(p)$ and $f^{2n}(p)$ are distinct.

Let s be the period of p . Then $f^{3n}(p) = p$ and it follows that 3 divides s . Therefore, s is not a power of 2. This establishes Theorem 7.

LEMMA 8. *Suppose $f: I \rightarrow I$ is continuous and onto. Suppose that J is a proper closed subinterval of I and, for each $n \geq 1$, $f^{-n}(J)$ is an interval. Then (I, f) is decomposable.*

PROOF. Let $H = \{\underline{x} | \underline{x} \in (I, f) \text{ and } \pi_0(\underline{x}) \in J\}$. Since J is proper and f is onto, H is a proper subset of (I, f) . Since, for each n , $f^{-n}(J)$ is an interval, $H_n = \pi_n(H) = f^{-n}(J)$, and so H is a subcontinuum of (I, f) . Now let $U = \pi_0^{-1}(\text{int } J)$. Then U is open in (I, f) and $U \subset H$. Therefore H is a proper subcontinuum of (I, f) with interior, and it follows that (I, f) is decomposable.

The following lemma is well known.

LEMMA 9. *Let $f: I \rightarrow I$ be continuous, $h: I \rightarrow I$ be a homeomorphism, and $f_1 = h \circ f \circ h^{-1}$. Then (I, f) and (I, f_1) are homeomorphic.*

PROOF. Define $H: (I, f) \rightarrow (I, f_1)$ by $H((x_0, x_1, \dots)) = (h(x_0), h(x_1), \dots)$. It is clear that H is a homeomorphism.

DEFINITION. If $f: I \rightarrow I$ is continuous, then the statement that f has *finitely many turning points* means that there is a finite set $\{a_0, a_1, \dots, a_l\}$, $a = a_0 < a_1 < \dots < a_l = b$ in $I = [a, b]$ such that f is monotone on $[a_{i-1}, a_i]$ for $i = 1, 2, \dots, l$.

THEOREM 10. *Suppose that $f: I \rightarrow I$ is continuous, onto, and has finitely many turning points. Then if (I, f) is indecomposable, f has a periodic point whose period is not a power of 2.*

PROOF. If f is organic, then the conclusion follows from Theorem 7. We will show that if f is inorganic, then f has a point of period 3. Suppose that f is inorganic. Then it follows from Lemma 6 that either

- (i) $f^{-1}(\{a\}) = \{a\}$,
- (ii) $f^{-1}(\{b\}) = \{b\}$, or
- (iii) $f(a) = b, f(b) = a$ and $f^{-1}(\{a, b\}) = \{a, b\}$.

Now if (ii) holds, define $h: I \rightarrow I$ by $h(x) = (a + b) - x$, and let $f_1 = h \circ f \circ h^{-1}$. Then $f_1^{-1}(\{a\}) = \{a\}$. It follows from Lemma 8 that $(I, f) \sim (I, f_1)$, and so the hypotheses of the theorem hold for f_1 .

If (iii) holds let $f_1 = f^2$. Then $f_1^{-1}(\{a\}) = \{a\}$ and, since $(I, f) \sim (I, f^2)$, the hypotheses of the theorem hold for f_1 .

It follows from the preceding discussion that we may assume that $f^{-1}(\{a\}) = \{a\}$. Let the turning points of f be $a = a_0 < a_1 < \dots < a_l = b$. We will consider two cases.

Case 1. $f^{-1}(\{b\}) = \{b\}$. Now suppose that there is a number c in (a, b) such that if $x \in [a, c]$, then $f(x) > x$. Since $f^{-1}(\{a\}) = \{a\}$, we may assume that c is chosen so that $f([c, b]) \cap [a, c] = \emptyset$. Then, for each $n \geq 0$, $f^{-n}([a, c])$ is an interval, and it follows from Lemma 8 that (I, f) is decomposable. Therefore for each $c \in (a, b)$ there is an $x \in (a, c)$ such that $f(x) \leq x$. Similarly, for each $c \in (a, b)$ there is an $x \in (c, b)$ such that $f(x) \geq x$. It follows that there are fixed points for f in (a, b) .

Now let $\mathcal{C} = \{c | a < c < b \text{ and } f([a, c]) = [a, c]\}$. We will show that $\mathcal{C}$ is nonempty. Let q be a point of (a, b) such that $f(q) = q$. Then either $f([a, q]) = [a, q]$ or there is a turning point a_i , $a < a_i < q$, with $f(a_i) > a_i$. But in this case there is a point q_1 , $a < q_1 < a_1$, with $f(q_1) = q_1$. Since there are only finitely many turning points there is a number c_0 , $a < c_0 < b$, such that $f(c_0) = c_0$, and $f([a, c_0]) = [a, c_0]$.

Now since $f([a, c_0]) = [a, c_0]$ and (I, f) is indecomposable, it follows from Lemma 8 that $f([c_0, b]) \cap [a, c_0] \neq \emptyset$. Let a_{n_0} be the smallest turning point such that $f(a_{n_0}) < c_0 < a_{n_0}$. Notice that $a_{n_0} \neq b$. Now, there is a q , $a_{n_0} < q < b$, such that $f(q) = q$. Let c_1 be the smallest such q . Now either $f([a, c_1]) = [a, c_1]$, or there is a number x , $c_0 < x < c_1$, with $f(x) > c_1$. If $x > a_{n_0}$, there is a q , $a_{n_0} < q < x$, such that $f(q) = q$, which contradicts the choice of c_1 . Therefore $c_0 < x < a_{n_0}$. Again, there is a q , $x < q < a_{n_0}$, such that $f(q) = q$. Now, we have $[c_0, c_1] \subset f([c_0, q])$ and $[c_0, c_1] \subset f([q, c_1])$, and using the same argument as in the proof of Theorem 7, we have a point of period 3. Thus we have the conclusion of the theorem, or $c_1 \in \mathcal{C}$. If $c_1 \in \mathcal{C}$ we repeat the argument, replacing c_0 with c_1 . Continuing this way we get $a_{n_1}, c_2, a_{n_3}, c_3, \dots$ with $a_{n_0} < a_{n_1} < a_{n_2} < \dots$. Since there are only finitely many turning points, the process will end in a finite number of steps with a point of period 3.

Case 2. There is a point $p \in (a, b)$ with $f(p) = b$. We proceed as in Case 1. We have $\mathcal{C} \cap (a, p) \neq \emptyset$. Choose c_0 such that $f(c_0) = c_0$ and $f([a, c_0]) = [a, c_0]$. Let a_{n_0} be as in Case 1, except now it might be that $a_{n_0} = b$. If $a_{n_0} = b$, there is a q , $p < q < b$, such that $f(q) = q$. Then $[c_0, a_{n_0}] \subset f([c_0, q])$ and $[c_0, q] \subset f([q, a_{n_0}])$. As before, it follows that f has a point of period 3. In fact, if $a_{n_0} > p$, the same result holds.

Thus we may assume that $a_{n_0} < p$. We may then find c_1 , $a_{n_0} < c_1 < p$, and proceed as in Case 1. Thus f has a point of period 3.

EXAMPLE 5. Let f be as sketched in the accompanying figure. Then it can be shown that (I, f) is indecomposable and that f is inorganic. It follows from the argument in Theorem 10 that f has a point of period 3.

COROLLARY 11. Suppose that $f: I \rightarrow I$ is continuous and has finitely many turning points. Then, there is an integer $l \geq 0$ and an indecomposable subcontinuum of (I, f) which is invariant under $\hat{f}^{2^l}$ if and only if f has a periodic point whose period is not a power of 2.

PROOF. Theorem 1 shows that if f has the required periodic point, then (I, f) has the required indecomposable subcontinuum.

Suppose then that S is an indecomposable subcontinuum of (I, f) which is invariant under $\hat{f}^{2^l}$. Let $g = f^{2^l}$. Then (I, f) is homeomorphic with (I, g) . Let S_1 be the image of S under the natural homeomorphism. Then S_1 is invariant under $\hat{g}$. Let $J = \pi_0(S_1)$. Then J is invariant under g , and g has finitely many turning points in J . We now apply Theorem 10 to $g: J \rightarrow J$ and find that g has a periodic point whose period is not a power of 2, and it follows that f has a periodic point whose period is not a power of 2.

REFERENCES

- [AY] J. Auslander and J. Yorke, *Interval maps, factors of maps, and chaos*, Tôhoku Math. J. (2) **32** (1980), 177–188.
- [Bi] R. H. Bing, *Snake-like continua*, Duke Math. J. **18** (1951), 653–663.
- [Bl] L. Block, *Homoclinic points of mappings of the interval*, Proc. Amer. Math. Soc. **72** (1978), 576–580.
- [H] G. W. Henderson, *The pseudo-arc as an inverse limit with one binding map*, Duke Math. J. **31** (1964), 421–425.
- [HY] J. Hocking and G. Young, *Topology*, Addison-Wesley, Reading, Mass., 1961.
- [N] Z. Nitecki, *Topological dynamics on the interval*, Ergodic Theory and Dynamical Systems. II, Proceedings, Special Year, Maryland 1979–80 (A. Katok, ed.), Birkhäuser, Basel, 1982, pp. 1–73.
- [S] A. Sarkovskii, *Coexistence of cycles of a continuous map of the line into itself*, Ukrain. Mat. Zh. **16** (1964), 61–71.

MATHEMATICAL SCIENCES RESEARCH INSTITUTE, 2223 FULTON STREET, BERKELEY, CALIFORNIA 94720

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WYOMING, LARAMIE, WYOMING 82071 (Current address of both authors)