LINEARIZING CERTAIN REDUCTIVE GROUP ACTIONS

BY

H. BASS AND W. HABOUSH

ABSTRACT. Is every algebraic action of a reductive algebraic group G on affine space $\mathbf{C}^n$ equivalent to a linear action? The “normal linearization theorem” proved below implies that, if each closed orbit of G is a fixed point, then $\mathbf{C}^n$ is G -equivariantly isomorphic to $(\mathbf{C}^n)^G \times \mathbf{C}^m$ for some linear action of G on $\mathbf{C}^m$.

Introduction. Let G be a reductive algebraic group acting (algebraically) on affine space $\mathbf{C}^n$. Is the action equivalent to a linear action, i.e. is the action linear after some global polynomial change of coordinates on $\mathbf{C}^n$? This question, posed in print by Kambayashi [K, Conjecture (3.1)], motivates the present paper. The first step in the problem is to find a fixed point (to play the role of the origin for the linear action).

Fixed points are known to exist in the following cases.

1. G is a finite p -group. (Classical; see [B-B1] for a simple algebraic proof due to Shafarevič.)

2. G is a torus. (Bialynicki-Birula [B-B1] deduces this easily from case 1, using Zariski density of the p -torsion subgroup of G .)

3. $G = \mathrm{SL}_2$ acting on $\mathbf{C}^n$ without three-dimensional orbits (Panyushev [P]).

Linearizability of the action has been proved in the following cases.

1. G is a torus acting effectively on $\mathbf{C}^n$ and either

(a) $\dim G = n$ or $n - 1$ [B-B1 and B-B2], or

(b) The action is “unmixed,” in the sense that no nontrivial character, together with its inverse, occur as weights of semi-invariant polynomials, and $\dim(\mathbf{C}^n/G) \leq 2$ [K-R].

2. G is connected semisimple and $n \leq 4$. The case $n = 1$ is trivial. The case $n = 2$ is an easy consequence of Jung’s Theorem [J] presenting the group of polynomial automorphisms of $\mathbf{C}^2$ as an amalgamated free product (cf. [Sh]). The case $n = 3$ is due to Kraft and Popov [KP], and the case $n = 4$ to D. I. Panyushev [P]. The latter results are recent, and make heavy use of Luna’s Étale Slice Theory [L]. They are the most penetrating results to date.

The linearization problem subsumes the “cancellation problem” for cartesian factors of $\mathbf{C}^n$: If $Y \times \mathbf{C}^m \cong \mathbf{C}^n$, must the variety Y be isomorphic to $\mathbf{C}^{n-m}$? This is known to be so only for $n - m \leq 2$ (cf. [Ru]). Suppose however that G is a reductive group with a linear representation on $\mathbf{C}^m$ fixing only the origin. Then G acts on

Received by the editors July 9, 1984.

1980 *Mathematics Subject Classification*. Primary 13B10, 14L17; Secondary 13F20, 20G05, 13C10.

©1985 American Mathematical Society
0002-9947/85 \$1.00 + \$.25 per page

$Y \times \mathbf{C}^m$ by $g(y, z) = (y, gz)$ ($g \in G$, $y \in Y$, $z \in \mathbf{C}^m$) and we have $(Y \times \mathbf{C}^m)^G = Y \times \{0\} \cong Y$. If, viewed as an action on $\mathbf{C}^n$ ($\cong Y \times \mathbf{C}^m$), this action can be linearized, then Y must be isomorphic to the linear space $(\mathbf{C}^n)^G$, whence $Y \cong \mathbf{C}^{n-m}$. Conversely if $Y \cong \mathbf{C}^{n-m}$, then the action can clearly be linearized.

Our main result, the “Normal Linearization Theorem” (Theorem (9.1)), is a generalization to arbitrary reductive groups of the Kambayashi-Russell result on “unmixed” torus actions, cited above. The appropriate interpretation of unmixedness is what we call a “fix pointed action” of a reductive group G on an affine scheme X , namely an action for which $\pi: X \rightarrow X/G$ induces an isomorphism $X^G \rightarrow X/G$. Assuming this plus a smoothness condition on $X^G \subset X$ we conclude that $\pi: X \rightarrow X/G$ admits the structure of a G -vector bundle; in particular the fibers are linear with linear G -action.

Supposing that X is an affine space, or, more generally (by the Quillen-Suslin proof of the Serre Conjecture) that vector bundles on X are trivial, we conclude further that X is G -isomorphic to $(X/G) \times W$, where W is some rational G -module with $W^G = 0$. The action is then linearizable iff the variety X/G is an affine space, as in the cancellation problem discussed above.

The Normal Linearization Theorem is actually formulated for situations more general than fix pointed actions. Moreover we make some observations en route on the structure of G -vector bundles on affine G -schemes X . Some of these are related to results of Richardson [R1 and R2], though our proofs are different and, we feel, somewhat easier.

A final section explains how the present results might be used to approach the Jacobian Conjecture [BCW], stating that polynomial endomorphisms of $\mathbf{C}^n$ with Jacobian determinant 1 are invertible.

1. The affine quotient $\pi: X \rightarrow X/G$. We fix an algebraically closed field k of characteristic zero. By a *scheme* X we shall mean a scheme of finite type over k . If X is affine its affine algebra will be denoted A_X , so that $X = \text{Spec}(A_X)$.

Let G be an algebraic group (over k); all such groups will be understood to be affine. We shall often confuse G with its group $G(k)$ of rational points over k . A scheme X together with an action of G , $G \times X \rightarrow X$, will be called a G -scheme.

Let X be an affine G -scheme. Suppose that the ring A_X^G of G -invariants is finitely generated over k . Classical invariant theory (cf. [Sp]) guarantees this whenever the group G is reductive. Then we put $X/G = \text{Spec}(A_X^G)$ and call $\pi = \pi_{X,G}: X \rightarrow X/G$ (induced by $A_X^G \subset A_X$) the *affine quotient* of the G -scheme X .

An orbit Gx is open in its closure $\overline{Gx}$, which consists of $\overline{Gx}$ plus lower dimensional orbits. In particular an orbit of minimal dimension in $\overline{Gx}$ is closed, so nonempty closed G -stable subsets of X contain closed orbits. Equivalently: *An open G -stable subset of X containing all closed orbits is all of X .* Part of the philosophy of this paper is to develop the latter as the geometric form of an “Equivariant Nakayama Lemma” (see §§5 and 6).

Suppose that G is reductive. If C_0, C_1 are disjoint closed G -stable subschemes of X , then there is an $f \in A_X^G$ such that $f|_{C_i} = i$ ($i = 0, 1$). Consequently, each fiber

$\pi^{-1}(\xi)$ ($\xi \in X/G$) contains a unique closed orbit, denoted $C(\xi)$, and

$$\pi^{-1}(\xi) = \{x \in X \mid C(\xi) \subset \overline{Gx}\}.$$

2. A - G -modules. Let G be an (affine) algebraic group (over k). A finite dimensional k -module V on which $G(k)$ acts k -linearly is called *rational* if, relative to a basis of V , the matrix entries of the representation belong to A_G . In general we understand a *rational G -module* to be a union of finite dimensional ones. We shall call these simply *k - G -modules*.

Let $X = \text{Spec}(A)$ be an affine scheme. An action of G on X corresponds to a (translation) action of G on $A = A_X$ as k -algebra automorphisms, making A a k - G -module as above. Conversely any such action of G on A defines an action on X .

By an *A - G -module* then we mean an A -module M which is simultaneously a k - G -module so that

$$s(am) = s(a) \cdot s(m)$$

for all $s \in G$, $a \in A$, $m \in M$. These form an abelian category. If M and N are A - G -modules so also is $M \otimes_A N$, with G -action $s(m \otimes n) = sm \otimes sn$. Further the symmetric algebra $S_A(M)$ and exterior algebra $\Lambda_A(M)$ similarly become A - G -modules. The G -action on $\text{Hom}_A(M, N)$, $(sf)(m) = s(f(s^{-1}m))$, is rational, and so makes $\text{Hom}_A(M, N)$ an A - G -module, provided that M is finitely A -generated. Further we have

$$\text{Hom}_A(M, N)^G = \text{Hom}_{A-G}(M, N).$$

3. Reductive groups. Let G be a reductive group. Then the k - G -modules are semisimple (cf. [N, Chapter VI]). Let Λ denote the set of classes of simple k - G -modules, and, for $\lambda \in \Lambda$, let V_λ be a simple k - G -module of type λ .

Let M be a k - G -module. Then

$$M = \bigoplus_{\lambda \in \Lambda} M_\lambda,$$

where M_λ is the isotypic component of type λ . Put $H_\lambda(M) = \text{Hom}_k(V_\lambda, M)$, and $H_\lambda^G(M) = H_\lambda(M)^G = \text{Hom}_{k-G}(V_\lambda, M)$. The canonical (evaluation) homomorphism

$$\varepsilon_{\lambda, M}: H_\lambda^G(M) \otimes_k V_\lambda \rightarrow M_\lambda$$

$$\varepsilon_{\lambda, M}(h \otimes v) = h(v)$$

is an isomorphism of k - G -modules (with trivial G -action on $H_\lambda^G(M)$). This reduces, by additivity, to the case $M = V_\lambda$, where it follows from Schur's Lemma (since k is algebraically closed).

Now suppose that G acts on $X = \text{Spec}(A)$ and that M is an A - G -module. Then $H_\lambda(M) = \text{Hom}_k(V_\lambda, M)$ is an A - G -module, finitely A -generated if M is so, $H_\lambda^G(M)$ is an A^G -module (with trivial G -action), and $\varepsilon_{\lambda, M}: H_\lambda^G(M) \otimes_k V_\lambda \rightarrow M_\lambda$ is an A^G - G -isomorphism.

(3.1) THEOREM (FINITE MULTIPLICITY). *Let G be a reductive group acting on the affine scheme $X = \text{Spec}(A)$. Let M be an A - G -module which is finitely A -generated. For each $\lambda \in \Lambda$, $H_\lambda^G(M) = \text{Hom}_{k-G}(V_\lambda, M)$ and $M_\lambda \cong H_\lambda^G(M) \otimes_k V_\lambda$ are finitely generated A^G -modules.*

PROOF. Form the symmetric algebra $S = S_A(M) = S_0 \oplus S_1 \oplus S_2 \oplus \cdots$ ($S_0 = A$, $S_1 = M$). It is a finitely generated k -algebra on which G acts rationally as graded algebra automorphisms. By classical invariant theory (cf. [Sp, Theorem 2.4.9]) S^G is a finitely generated k -algebra. Since $S^G = A^G \oplus M^G \oplus S_2^G \oplus \cdots$ it follows easily that M^G is a finitely generated A^G -module. Applying this to $H_\lambda(M) = \text{Hom}_k(V_\lambda, M)$, which is a finitely A -generated A - G -module, in place of M , we conclude that $H_\lambda^G(M)$ is finitely A^G -generated, whence likewise for $M_\lambda \cong H_\lambda^G(M) \otimes_k V_\lambda$.

(3.2) COROLLARY. *If the reductive group G acts on the affine scheme $X = \text{Spec}(A)$, then for each $\lambda \in \Lambda$, A_λ is a finitely generated A^G -module.*

Apply the theorem to $M = A$.

(3.3) REMARKS. Let $\xi \in X/G$ correspond to the maximal ideal $m \subset A^G$. Then it is clear that $H_\lambda(A/mA) = H_\lambda(A)/mH_\lambda(A)$, and, by semisimplicity of k - G -modules, $H_\lambda^G(A/mA) = H_\lambda^G(A)/mH_\lambda^G(A)$. Thus the (finite) multiplicity of V_λ in $A_{\pi^{-1}(\xi)} = A/mA$ is

$$m_\lambda(\xi) = \dim_k(H_\lambda^G(A)/mH_\lambda^G(A)).$$

In case $H_\lambda^G(A)$ is a flat, hence (by (3.2)) projective, A^G -module, then $m_\lambda(\xi)$ is a locally constant function of ξ , hence constant if X/G is connected. In this case $m_\lambda = m_\lambda(\xi)$ can be computed on any convenient fiber $\pi^{-1}(\xi)$.

Suppose that a fiber $\pi^{-1}(\xi)$ is a (reduced) closed orbit $Gx \cong G/H$, where $H = G_x$. Then A/mA is G -isomorphic to $A_{G/H} = (A_G)^H$, where H acts on A_G via right translation on G . Now as a $G \times G$ -module, via left and right translations, $A_G \cong \bigoplus_{\lambda \in \Lambda} (V_\lambda \otimes_k V_\lambda^*)$, where $(s, t) \in G \times G$ acts on $V_\lambda \otimes_k V_\lambda^*$ as $\rho_\lambda(s) \otimes \rho_\lambda^*(t)$. Moreover, $V_\lambda \otimes_k V_\lambda^*$ is, for left translation, the λ -isotypic component $(A_G)_\lambda$ of A_G . Thus we have $(A_{G/H})_\lambda \cong (V_\lambda \otimes_k V_\lambda^*)^H = V_\lambda \otimes_k (V_\lambda^*)^H$, so the multiplicity of V_λ in $A_{G/H}$ is

$$m_\lambda(G/H) = \dim_k(V_\lambda^*)^H,$$

where $(V_\lambda^*)^H = \text{Hom}_{k-H}(V_\lambda, k)$.

4. Projective A - G -modules. Consider a reductive group G acting on an affine scheme $X = \text{Spec}(A)$.

(4.1) PROPOSITION ("SPLITTING LEMMA"). *Let*

$$(*) \quad 0 \rightarrow M' \rightarrow M \xrightarrow{p} M'' \rightarrow 0$$

be an exact sequence of A - G -modules with M'' finitely A -generated. If $()$ splits A -linearly, then it splits A - G -linearly.*

PROOF. Since $(*)$ splits A -linearly,

$$\text{Hom}_A(M'', M) \xrightarrow{p \circ -} \text{Hom}_A(M'', M'') \rightarrow 0$$

is exact. Since M'' is finitely A -generated the latter are A - G -modules, hence semisimple G -modules since G is reductive. Thus we obtain a still exact sequence of G -invariants.

$$\mathrm{Hom}_{A-G}(M'', M) \xrightarrow{p \circ -} \mathrm{Hom}_{A-G}(M'', M'') \rightarrow 0.$$

Lifting $1_{M''}$ to $\mathrm{Hom}_{A-G}(M'', M)$ gives the desired splitting of $(*)$.

(4.2) COROLLARY. *Let P be a finitely A -generated A - G -module. The following conditions are equivalent.*

(a) P is A -projective.

(b) P is A - G -projective.

(c) P is a direct summand of $V_A \stackrel{\mathrm{def}}{=} A \otimes_k V$ for some finite dimensional k - G -module V .

PROOF. Let V be a k - G -module and M an A - G -module. The natural isomorphism $\mathrm{Hom}_k(V, M) \cong \mathrm{Hom}_A(V_A, M)$ is an A - G -isomorphism, whence $\mathrm{Hom}_{A-G}(V_A, M) \cong \mathrm{Hom}_{k-G}(V, M)$. Since k - G -modules are semisimple, $\mathrm{Hom}_{k-G}(V, -)$ is exact, hence $\mathrm{Hom}_{A-G}(V_A, -)$ is exact, so V_A is A - G -projective, whence (c) $\Rightarrow$ (b). Since V_A is A -free, (c) $\Rightarrow$ (a).

Choose in P a finite dimensional k - G -module V that generates P as an A -module. The inclusion $V \subset P$ defines an A - G -homomorphism $V_A \rightarrow P$ that is surjective. Assuming (b), it splits, whence (b) $\Rightarrow$ (c). Assuming (a) also, it splits, by the splitting lemma, whence (a) $\Rightarrow$ (c).

(4.3) THEOREM. *Let M be a finitely A -generated A - G -module. The following conditions are equivalent.*

(a) M is A^G -flat.

(b) M is A^G -projective.

(c) M is A^G - G -projective.

(d) $H_\lambda^G(M)$ is A^G -projective for each $\lambda \in \Lambda$.

PROOF. An A^G - G -module M decomposes as

$$M = \bigoplus_{\lambda} M_{\lambda} = \bigoplus_{\lambda} H_{\lambda}^G(M) \otimes_k V_{\lambda}$$

with each M_{λ} and $H_{\lambda}^G(M)$ finitely A^G -generated. Whence

$$\begin{aligned} M \text{ is } A^G\text{-flat} &\Leftrightarrow M_{\lambda} \text{ is } A^G\text{-flat} \quad \forall \lambda \\ &\Leftrightarrow H_{\lambda}^G(M) \text{ is } A^G\text{-flat} \quad \forall \lambda \\ &\Leftrightarrow H_{\lambda}^G(M) \text{ is } A^G\text{-projective} \quad \forall \lambda \\ &\Leftrightarrow M_{\lambda} \text{ is } A^G\text{-projective} \quad \forall \lambda \\ &\Leftrightarrow M \text{ is } A^G\text{-projective.} \end{aligned}$$

By Corollary (4.2), M_{λ} is A^G -projective $\forall \lambda \Leftrightarrow M_{\lambda}$ is A^G - G -projective $\forall \lambda \Leftrightarrow M$ is A^G - G -projective. Whence the theorem.

(4.4) COROLLARY. Suppose that projective A^G -modules are free. If M is a finitely A -generated A - G -module that is A^G -flat, then there is a k - G -module W with finite multiplicities (i.e. $\dim_k W_\lambda < \infty \forall \lambda$) such that M is isomorphic as an A^G - G -module to $A^G \otimes_k W$.

PROOF. The hypotheses and the theorem imply that, for each λ , $H_\lambda^G(M) \cong (A^G)^{n_\lambda}$ for some $n_\lambda \geq 0$. Thus M , as an A^G - G -module, is isomorphic to

$$\begin{aligned} \bigoplus_\lambda (H_\lambda^G(M) \otimes_k V_\lambda) &\cong \bigoplus_\lambda ((A^G)^{n_\lambda} \otimes_k V_\lambda) \\ &\cong \bigoplus_\lambda (A^G \otimes_k V_\lambda^{n_\lambda}) \cong A^G \otimes_k W, \end{aligned}$$

where $W = \bigoplus_\lambda V_\lambda^{n_\lambda}$.

(4.5) REMARK. In case projective A^G -modules are free and A is A^G -flat, then, by (4.4), $A \cong A^G \otimes_k W$ for some k - G -module $W \subset A$ with finite multiplicities. Cases where this situation prevails are treated by Richardson (see [R1, and R2, Theorem B]).

5. The G -radical. Let G be a reductive group. For any k - G -module V we have

$$V = V^G \oplus \Delta_G(V),$$

where $\Delta_G(V)$ is the sum of the nontrivial isotypic components of V , or, alternatively, it is the sub- k -module generated by all $sv - v$ ($s \in G, v \in V$).

Suppose that G acts on the affine scheme $X = \text{Spec}(A)$. Then $A = A^G \oplus \Delta_G(A)$ is a decomposition as an A^G -module. We put

$$\begin{aligned} \Delta_G^+(A) &= \text{the smallest ideal of } A \text{ containing } \Delta_G(A) \\ &= \text{the ideal of } A \text{ generated by } \Delta_G(A), \end{aligned}$$

and

$$\begin{aligned} \Delta_G^-(A) &= \text{the largest ideal of } A \text{ contained in } \Delta_G(A) \\ &= \text{the sum of all } A\text{-ideals contained in } \Delta_G(A). \end{aligned}$$

These are G -stable ideals of A , and

$$\Delta_G^-(A) \subset \Delta_G(A) \subset \Delta_G^+(A).$$

Clearly $\Delta_G^+(A)$ is the smallest ideal of A modulo which the action of G is trivial. Hence it defines the scheme of fixed points,

$$X^G = \text{Spec}(A/\Delta_G^+(A)) \subset \text{Spec}(A) = X.$$

We have $\Delta_G^+(A) = \Delta_G^+(A)^G \oplus \Delta_G(A)$, so that $A/\Delta_G^+(A) = A^G/\Delta_G^+(A)^G$, and the diagram

$$\begin{array}{ccc} X^G & \subset & X \\ & \searrow & \downarrow \pi \\ & & X/G \end{array}$$

corresponding to

$$\begin{array}{ccc} A/\Delta_G^+(A) & \leftarrow & A \\ & \nwarrow & \cup \\ & & A^G \end{array}$$

identifies X^G with a closed subscheme of X/G .

We now interpret $\Delta_G^-(A)$ geometrically.

If $x \in X(k)$ let $m_x \subset A$ denote its maximal ideal. If $s \in G$, then $s(m_x) = m_{sx}$.

Put

$$P_x = \bigcap_{s \in G} s(m_x) = \bigcap_{s \in G} m_{sx},$$

the largest G -stable ideal contained in m_x . Then every G -stable proper ideal is contained in some P_x . Further $\text{Spec}(A/P_x) \subset X$ is just the (reduced) orbit closure $\overline{Gx}$. Hence if G is connected, then P_x is prime. Further we have

$$P_x \subset P_y \Leftrightarrow \overline{Gy} \subset \overline{Gx}.$$

Hence

(5.1) PROPOSITION. P is a maximal G -stable proper ideal of A iff $P = P_x$ for some $x \in X$ such that Gx is closed, and then $Gx = \text{Spec}(A/P_x) \subset X$.

The G -radical of A is, by definition,

$$G\text{-rad}(A) = \bigcap_{Gx \text{ closed}} P_x.$$

Thus $\text{Spec}(A/G\text{-rad}(A))$ is the reduced subscheme corresponding to the closure of the union of the closed orbits; we shall call it briefly the “closed orbit closure”. It is the largest reduced closed subscheme of X on which the G -action is “stable” in the sense of Popov [Po]. Since $G\text{-rad}(A)$ is G -stable we have

$$G\text{-rad}(A) = G\text{-rad}(A)^G \oplus \Delta_G(G\text{-rad}(A)).$$

(5.2) PROPOSITION. (a) $G\text{-rad}(A)^G = \text{nil}(A^G)$, the nil radical of A^G .

(b) $G\text{-rad}(A) = \sqrt{\Delta_G^-(A)}$, the radical of $\Delta_G^-(A)$.

(c) If A^G is reduced, e.g. if A is reduced, then $G\text{-rad}(A) = \Delta_G^-(A)$.

(d) If A^G is an integral domain and G is connected, then $G\text{-rad}(A) (= \Delta_G^-(A))$ is a prime ideal.

PROOF. Let $\xi \in X/G$ with maximal ideal $m_\xi \subset A^G$. Then $\pi^{-1}(\xi) = \text{Spec}(A/m_\xi A)$. Choose $x \in C(\xi)$, the closed orbit in $\pi^{-1}(\xi)$, and write P_ξ for P_x ; P_ξ is the unique maximal G -stable ideal of A containing $m_\xi A = m_\xi \oplus m_\xi \Delta_G(A)$. We have

$$\begin{aligned} G\text{-rad}(A) &= \bigcap_{\xi \in X/G} P_\xi = \bigcap_{\xi \in X/G} (m_\xi \oplus m_\xi \Delta_G(A)) \\ &= \left(\bigcap_{\xi \in X/G} m_\xi \right) \oplus \left(\bigcap_{\xi \in X/G} m_\xi \Delta_G(A) \right) \\ &= \text{nil}(A^G) \oplus \Delta_G(G\text{-rad}(A)), \end{aligned}$$

by the Nullstellensatz (applied to A^G); whence (a).

If A^G is reduced, then $G\text{-rad}(A) = \Delta_G(G\text{-rad}(A))$ is an ideal contained in $\Delta_G(A)$, hence contained in $\Delta_G^-(A)$, by definition. Now (c) follows by showing that $\Delta_G^-(A) \subset G\text{-rad}(A)$. It suffices to show that each P_ξ ($\xi \in X/G$) contains $\Delta_G^-(A)$. If not, then by maximality of P_ξ we have $A = P_\xi + \Delta_G^-(A) = m_\xi \oplus (\Delta_G(P_\xi) + \Delta_G^-(A)) \subset m_\xi \oplus \Delta_G(A)$, which is absurd; this proves (b).

To show that $G\text{-rad}(A) \subset \sqrt{\Delta_G^-(A)}$ note first that both sides contain $\text{nil}(A)$, which is a G -stable ideal. Moreover $G\text{-rad}(A/\text{nil}(A)) = G\text{-rad}(A)/\text{nil}(A)$ and $\Delta_G^-(A/\text{nil}(A)) = \Delta_G^-(A)/(\Delta_G^-(A) \cap \text{nil}(A))$. Thus we may pass to $A/\text{nil}(A)$, and reduce to the case when A is reduced. Then, by (a), $G\text{-rad}(A) \subset \Delta_G(A)$ so, by definition of $\Delta_G^-(A)$, $G\text{-rad}(A) \subset \Delta_G^-(A)$. This concludes the proof of (c).

Suppose finally that A^G is an integral domain. Let $\Delta_G^-(A) = Q_1 \cap \cdots \cap Q_n$ be the primary decomposition of $\Delta_G^-(A)$ ($= G\text{-rad}(A)$, by (c)). Since $G\text{-rad}(A)$ is a radical ideal each Q_i is prime. Since $G\text{-rad}(A)$ is G -stable, G permutes the Q_i 's. Suppose that G is connected. Then G stabilizes each Q_i . Hence $Q_i = Q_i^G \oplus \Delta_G(Q_i)$. Then $Q_1^G \cap \cdots \cap Q_n^G = G\text{-rad}(A)^G = 0$. Since A^G is an integral domain there is an i such that $Q_i^G = 0$. Then $Q_i = \Delta_G(Q_i) \subset \Delta_G(A)$ so, by definition of $\Delta_G^-(A)$, $Q_i \subset \Delta_G^-(A)$. But then $Q_i = \Delta_G^-(A)$, so $\Delta_G^-(A)$ is prime.

(5.3) PROPOSITION. *Let J be a G -stable ideal of A and put $X_0 = \text{Spec}(A/J) \subset X = \text{Spec}(A)$. The following conditions are equivalent.*

- (i) X_0 contains all closed orbits in X .
- (ii) $J \subset G\text{-rad}(A)$.
- (iii) J^G is a nilpotent ideal in A^G .

The remarks preceding (5.2) show that (i) $\Leftrightarrow$ (ii), and (ii) $\Rightarrow$ (iii) by (5.2)(a). To show that (iii) $\Rightarrow$ (ii), assume that J^G is nilpotent. Put $A_{\text{red}} = A/\text{nil}(A)$ and $J_{\text{red}} =$ the image of J in A_{red} . Then $J_{\text{red}}^G = 0$ so $J_{\text{red}} \subset \Delta_G^-(A_{\text{red}}) \subset G\text{-rad}(A_{\text{red}})$ (by (5.2)(c)). Therefore J is contained in the inverse image in A of $G\text{-rad}(A_{\text{red}})$, which is $G\text{-rad}(A)$.

6. Equivariant Nakayama Lemma. Let G be a reductive group acting on $X = \text{Spec}(A)$. We fix a G -stable ideal, $J \subset G\text{-rad}(A)$.

(6.1) PROPOSITION ("EQUIVARIANT NAKAYAMA LEMMA"). *Let M be a finitely A -generated A - G -module.*

- (a) *If $JM = M$, then $M = 0$.*
- (b) *If N is an A - G -submodule of M and if $M = N + JM$, then $M = N$.*

PROOF. Applying (a) to M/N one obtains (b). To prove (a) recall that $JM = M$ iff $J + \text{ann}_A(M) = A$, where $\text{ann}_A(M) = \{a \in A \mid aM = 0\}$, a G -stable ideal. (See [NB1, II, §4, no. 4, Corollaire de la Proposition 18].) Since J is contained in every maximal G -stable ideal, $\text{ann}_A(M)$ cannot be contained in a maximal G -stable ideal, so $\text{ann}_A(M) = A$, i.e. $M = 0$.

(6.2) PROPOSITION. *Let P, Q be finitely A -generated A - G -modules. Assume that P is A -projective.*

(a) *Any A - G -homomorphism $u_0: P/JP \rightarrow Q/JQ$ lifts to an A - G -homomorphism $u: P \rightarrow Q$.*

(b) If u_0 is surjective, then u is surjective.

(c) If u_0 is an isomorphism and if Q also is A -projective, then u is an isomorphism.

PROOF. (a) In the commutative diagram of A - G -homomorphisms

$$\begin{array}{ccccc} & & & & Q \\ & & & \nearrow u & \downarrow \\ P & \longrightarrow & P/JP & \longrightarrow & Q/JQ \\ & & & \searrow u_0 & \\ & & & & \end{array}$$

u exists because P is A - G -projective (Corollary (4.2)).

(b) If u_0 is surjective and $C = \text{Coker}(u)$, then $C/JC = \text{Coker}(u_0) = 0$, so $C = 0$ by the Equivariant Nakayama Lemma.

(c) Suppose that u_0 is an isomorphism and that Q is A -projective. Then u is surjective, by (b), and so splits A -linearly (in fact A - G -linearly, by Corollary (4.2)). It follows that $K = \text{Ker}(u)$ is a direct summand of P , so that $K/JK = \text{Ker}(u_0) = 0$, whence $K = 0$ by Nakayama again.

(6.3) COROLLARY. Let P be an A - G -module, finitely generated and projective as an A -module. Let V be a finite dimensional k - G -module. Suppose that P/JP and $V_{A/J} = (A/J) \otimes_k V$ are isomorphic as (A/J) - G -modules. Then P and $V_A = A \otimes_k V$ are isomorphic as A - G -modules.

This follows from the proposition, part (c), with V_A in the role of Q .

(6.4) COROLLARY. Suppose that $A = A_0 \oplus J$ for some G -stable subalgebra A_0 . Let P be an A - G -module finitely generated and projective as an A -module. Then P is A - G -isomorphic to $A \otimes_{A_0} P_0$, where P_0 is the A_0 - G -module P/JP (and we identify A/J with A_0).

We apply the proposition, part (c), with $Q = Q \otimes_{A_0} P_0$, noting that $Q/JQ \cong P_0$.

(6.5) Geometric interpretation. Let $X_0 = \text{Spec}(A/J) \subset X = \text{Spec}(A)$. The condition that J is G -stable and contained in $G\text{-rad}(A)$ signifies that X_0 is G -stable and contains all closed orbits. Then A - G -modules P, Q define coherent G -sheaves $\tilde{P}, \tilde{Q}$ on X , with $\tilde{P}$ locally free, i.e. a “ G -vector bundle”. Their restrictions to X_0 are $\tilde{P}_0, \tilde{Q}_0$, where $P_0 = P/JP, Q_0 = Q/JQ$.

(a) Any G -morphism $u_0: \tilde{P}_0 \rightarrow \tilde{Q}_0$ extends to a G -morphism $u: \tilde{P} \rightarrow \tilde{Q}$.

(b) If u_0 is surjective, then u is surjective.

(c) If u_0 is an isomorphism and $\tilde{Q}$ also is locally free, then u is an isomorphism.

7. Luna’s Equivariant Zariski Main Theorem. We shall make use of the following results, quoted for reference.

(7.1) THEOREM (LUNA [L, I, §4]). Let G be a reductive group. Let X, Y be affine G -schemes. Let $\varphi: X \rightarrow Y$ be a G -morphism with finite fibers. Then there is a factorization

$$\begin{array}{ccccc} & & Z & & \\ & \nearrow i & & \searrow \psi & \\ X & & \rightarrow & & Y \\ & & \varphi & & \end{array}$$

where Z is an affine G -scheme, i and ψ are G -morphisms, i is an open immersion with $i^0: A_Z \rightarrow A_X$ injective, and ψ is a finite morphism.

(7.2) COROLLARY (LUNA [L, I, §4]). Assume above that:

(a) φ preserves closed orbits (i.e. Gx closed in $X \Rightarrow G\varphi(x)$ is closed in Y).

(b) φ is quasi-finite (i.e. has finite fibers).

(c) $\varphi/G: X/G \rightarrow Y/G$ is finite.

Then φ is finite.

8. Local complete intersections (LCI) and retracts. We fix an affine scheme $X = \text{Spec}(A)$, an ideal $J \subset A$, $A_0 = A/J$, and the closed subscheme $X_0 = \text{Spec}(A_0)$ of X . Consider the A_0 -module $N = J/J^2$. The graded A_0 -algebra

$$\text{gr}_J(A) = \bigoplus_{n \geq 0} J^n/J^{n+1}$$

is generated by N in degree 1, so there is a canonical surjection of graded A_0 -algebras

$$\gamma: S = S_{A_0}(N) \rightarrow \text{gr}_J(A).$$

One says that X_0 is a *local complete intersection* in X , or that J is a *local complete intersection ideal* in A , if the following conditions hold:

(i) The A_0 -module $N = J/J^2$ is projective.

(ii) $\gamma: S = S_{A_0}(N) \rightarrow \text{gr}_J(A)$ is an isomorphism.

(8.1) EXAMPLE. A (closed) point $x \in X$ is a local complete intersection in X iff x is a *regular* (or *smooth*) point of X , i.e. its local ring A_{m_x} is regular. (See [NB2, VIII, §5, Theorem 1].) We shall write X_{reg} for the set of regular points of X , and call X (and A) *regular* if $X_{\text{reg}} = X$. In general X_{reg} is open in X (cf. [EGA, IV, (6.12.5)]).

(8.2) REMARK. Suppose that a reductive group G acts on X . Then X_{reg} is G -stable. If X_{reg} contains all closed orbits, then, since the G -stable closed set $X - X_{\text{reg}}$ contains no closed orbits, it must be empty, i.e. X is regular.

(8.3) LEMMA. If X_0 is regular and X_0 is a local complete intersection in X , then $X_0 \subset X_{\text{reg}}$.

(A background reference for this proof is [NB2, VIII, §5].)

Let m be a maximal ideal of A containing J . We must show that A_m is regular. By assumption, $(A_0)_{m_0} = A_m/J_m$ is regular, where $m_0 = m/J$. Further it is clear that J_m is a local complete intersection ideal in A_m , by exactness of localization. It follows that J_m is generated by a regular sequence $a_1, \dots, a_r$ in A_m ; in particular $\dim(A_0)_{m_0} = \dim A_m - r$. Since $m_0(A_0)_{m_0} = mA_m/J_m$ is generated by $\dim(A_0)_{m_0}$ elements (by regularity) it follows that mA_m is generated by $r + \dim(A_0)_{m_0} = \dim A_m$ elements, so A_m is regular.

(8.4) PROPOSITION. Suppose that $A = A_0 \oplus J$ for some subalgebra A_0 of A (i.e. X_0 is a retract of X). Then the following conditions are equivalent.

(a) $X_0 \subset X_{\text{reg}}$.

(b) (0) X_0 is regular,

(1) X_0 is a local complete intersection in X .

The lemma gives (b) $\Rightarrow$ (a). Assume (a), let m_0 be a maximal ideal of A_0 , and put $m = m_0 \oplus J$. We are given that A_m is regular and must show that $(A_0)_{m_0}$ is regular and that J_m is a local complete intersection ideal in A_m . We have

$$m/m^2 = (m_0/m_0^2) \oplus (J/mJ)$$

and $J/mJ = J/(m_0J + J^2) = N/m_0N$. Choose $a_1, \dots, a_{d_0} \in m_0$ to lift a k -basis of m_0/m_0^2 and $a_{d_0+1}, \dots, a_d \in J$ to lift a k -basis of J/mJ . Then $a_1, \dots, a_d$ is a regular system of parameters in A_m , and, by Nakayama's Lemma, J_m is generated by the regular sequence $a_{d_0+1}, \dots, a_d$. Both conclusions now follow easily.

9. The Normal Linearization Theorem.

(9.1) THEOREM. Let G be a reductive group acting on the affine scheme $X = \text{Spec}(A)$. Let $X_0 = \text{Spec}(A/J)$ be a closed subscheme of X . Assume:

- (a) X_0 is G -stable and contains all closed orbits.
- (b) There is a G -equivariant retraction $\pi: X \rightarrow X_0$.
- (c) X_0 is a local complete intersection in X .

Then $\pi: X \rightarrow X_0$ admits the structure of a G -vector bundle over X_0 .

(9.2) REMARKS. 1. Algebraically the conditions of the theorem translate as follows.

- (a) J is G -stable and $J \subset G\text{-rad}(A)$.
- (b) $A = A_0 \oplus J$ for some G -stable subalgebra A_0 .
- (c) J is a local complete intersection ideal in A .

The conclusion then is that A , as an A_0 -algebra with G -action, is isomorphic to a symmetric algebra $S = S_{A_0}(N)$ where N is an A_0 - G -module finitely generated and projective as an A_0 -module.

2. By Corollary (6.4), conditions (a) and (b) imply that every G -vector bundle on X is the pullback of one on X_0 .

3. Suppose that $X_0 \subset X_{\text{reg}}$, and assume (b). Then Proposition (8.4) implies condition (c), that X_0 is a local complete intersection in X , as well as the regularity of X_0 . The conclusion of the theorem then further implies that X itself is regular.

4. Put $N = J/J^2$. Condition (c) means:

- (c)(i) N is a projective A_0 -module,
- (c)(ii) $\gamma: S = S_{A_0}(N) \rightarrow \text{gr}_J(A)$ is an isomorphism.

Consider the exact sequence of A_0 - G -modules,

$$0 \rightarrow J^2 \rightarrow J \rightarrow N \rightarrow 0.$$

It splits A_0 -linearly, by (c)(i), hence A_0 - G -linearly, by the Splitting Lemma (Proposition (4.1)). Let $u^0: N \rightarrow J$ be such an A_0 - G -splitting. It extends to an A_0 -algebra homomorphism $u^0: S = S_{A_0}(N) \rightarrow A$ which is still G -equivariant. The corresponding G -morphism $u: X \rightarrow Y = \text{Spec}(S)$ will be called a G -linear approximation to (X, X_0) . Its existence required only conditions (b) and (c)(i). The theorem will be proved by showing that any such G -linear approximation is an isomorphism.

PROOF OF THEOREM (9.1). Let

$$\begin{array}{ccc} X = \operatorname{Spec}(A) & \xrightarrow{u} & Y = \operatorname{Spec}(S) \\ & \searrow \quad \swarrow & \\ & X_0 = \operatorname{Spec}(A_0) & \end{array}$$

be a G -linear approximation to (X, X_0) , as in Remark 4 above. We first prove:

(1) u is étale.

The locus in X where u is étale is open and G -stable. Hence it is all of X once it contains all closed orbits. It suffices therefore to show that u is étale at each $x \in X_0 \subset X$. The maximal ideal in A of x has the form $m_x = m_0 + J$, where m_0 is maximal in A_0 . By the construction of u_0 : $S = S_{A_0}(N) \rightarrow A$, and the assumption that X_0 is a local complete intersection in X , the map of associated graded rings

$$\operatorname{gr}(u^0): \operatorname{gr}_{S_+}(S) = S \rightarrow \operatorname{gr}_J(A),$$

where $S_+ = N \cdot S$, is an isomorphism. Since $\bigcap_n S_+^n = 0$ it follows that u^0 is injective; we shall view u^0 as an inclusion, $S \subset A$. The maximal ideal in S of $y = u(x)$ is $m_y = m_x \cap S = m_0 \oplus S_+$. We want to show that $S_{m_y} \rightarrow A_{m_x}$ is étale. It suffices to show that the induced map of completed local rings $\hat{S}_{m_y} \rightarrow \hat{A}_{m_x}$ is an isomorphism. The isomorphism $\operatorname{gr}(u^0)$ of associated graded rings implies an isomorphism from the S_+ -adic completion $\bar{S}$ of S to the J -adic completion $\bar{A}$ of A , one which carries $m_y \bar{S} = m_0 \oplus \bar{S}_+$ to $m_x \bar{A} = m_0 \oplus \bar{J}$. Since $\hat{S}_{m_y}$ is the $m_y \bar{S}$ -adic completion of $\bar{S}$, and $\hat{A}_{m_x}$ is the $m_x \bar{A}$ -adic completion of $\bar{A}$, we conclude, as desired, that $\hat{S}_{m_y} \rightarrow \hat{A}_{m_x}$ is an isomorphism.

(2) The image of $X_0 = \operatorname{Spec}(A_0) \xrightarrow{\cong} \operatorname{Spec}(S/S_+) \subset Y$ contains all closed orbits of Y . Further $u: X \rightarrow Y$ carries closed orbits in X to closed orbits in Y .

We have $S_+ = S \cap J$ so $S_+^G \subset J^G$, and J^G is nilpotent (in A^G) by (5.3), hence S_+^G is nilpotent, so, by (5.3) again, $S_+ \subset G\text{-rad}(S)$, i.e. $X_0 \subset Y$ contains all closed orbits. Since $u: X \rightarrow Y$ induces the identity from $X_0 \subset X$ to $X_0 \subset Y$, closed orbits are preserved.

(3) $u/G: X/G \rightarrow Y/G$ is a finite morphism.

We have $S = A_0 \oplus S_+ \subset A = A_0 \oplus J$, and we must show that $A^G = A_0^G \oplus J^G$ is a finitely generated module over $S^G = A_0^G \oplus S_+^G$. This follows because J^G , being nilpotent, is a finitely generated A_0^G -module.

Now since an étale morphism is quasi-finite, conclusions (1), (2) and (3) above supply the hypotheses of Luna's Equivariant ZMT (Corollary (7.2)). It follows that:

(4) u is finite, hence a finite étale covering.

The degree of this covering is locally constant over Y . The locus in Y where u has degree 1 is open and G -stable. To show that u is an isomorphism, i.e. everywhere of degree 1, it therefore suffices to show that u has degree 1 over each closed orbit of Y . Let Gy ($y \in Y$) be a closed orbit; note that $Gy \subset X_0 \subset Y$, by (3). Since u is finite, $u^{-1}(Gy)$ must be a finite union of G -orbits of the same dimension as Gy , hence each of them is closed. Consequently, $u^{-1}(Gy) \subset X_0 \subset X$. Since u maps $X_0 \subset X$ isomorphically to $X_0 \subset Y$, u has degree 1 over $X_0 \subset Y$, hence u is an isomorphism. This proves Theorem (9.1).

10. Fix pointed actions. Let G be a reductive group acting on $X = \text{Spec}(A)$. Recall that $A = A^G \oplus \Delta_G(A)$.

(10.0) DEFINITION. Call the action of G on X or on A *fix pointed* if the following equivalent conditions hold.

(a) $\Delta_G(A)$ is an ideal, i.e. $\Delta_G^-(A) = \Delta_G(A) = \Delta_G^+(A)$, in the notation of §5.

(b) $\pi: X \rightarrow X/G$ induces an isomorphism $X^G \rightarrow X/G$.

Indeed the latter corresponds to the surjective homomorphism $A^G \rightarrow A/\Delta_G^+(A)$. The following equivalent conditions are an immediate consequence of (a) and (b).

(c) Every closed orbit is a point.

(d) Every nonempty closed G -stable subset of X contains a fixed point.

(e) $\Delta_G^+(A)^G$ is a nilpotent ideal in A^G .

If X/G is reduced, e.g. if X is reduced, then all of the above conditions are equivalent (Proposition (5.2)).

We call the action *one fix pointed* if it is fix pointed with a unique fixed point (so that A^G is an artin local ring).

Let Λ denote the set of types of simple k - G -modules, and, for each $\lambda \in \Lambda$, let V_λ be a simple k - G -module of type λ .

(10.1) PROPOSITION. Assume that G acts fix pointedly on X . Let P be an A - G -module that is finitely generated and projective as an A -module. Let Q denote the A^G - G -module $P/\Delta_G(A)P$.

(a) $P \cong A \otimes_{A^G} Q$ (A - G -module isomorphism).

(b) The λ -isotypic component Q_λ of Q is isomorphic as an A^G - G -module to $H_\lambda \otimes_k V_\lambda$, where $H_\lambda = \text{Hom}_{k-G}(V_\lambda, Q)$, a projective A^G -module with trivial G -action. Hence P is isomorphic as an A - G -module to $\bigoplus_\lambda A \otimes_{A^G} H_\lambda \otimes_k V_\lambda$.

(c) Suppose that projective A^G -modules are free. (This is automatic if projective A -modules are free.)

Then P is A - G -isomorphic to $W_A = A \otimes_k W$ for some k - G -module W .

Assertion (a) follows from Corollary (6.4), and (b) follows then from (a) and the discussion preceding Theorem (3.1). If projective A^G -modules are free, then we have, for each $\lambda \in \Lambda$, $H_\lambda \cong (A^G)^{n_\lambda}$ for some $n_\lambda \geq 0$. Hence $A \otimes_{A^G} H_\lambda \otimes_k V_\lambda \cong A \otimes_k V_\lambda^{n_\lambda}$, so $P \cong A \otimes_k W$, where $W = \bigoplus_\lambda V_\lambda^{n_\lambda}$. Finally, the parenthetical assertion of (c) follows from:

(10.2) REMARK. If $B \xrightarrow{i} C \xrightarrow{p} B$ is a retraction of rings, $p \circ i = \text{Id}_B$, and if projective C -modules are free, then every projective B -module P is free. Indeed $P \cong B \otimes_{C,p} (C \otimes_{B,i} P)$, and $C \otimes_{B,i} P$ is a free C -module.

Geometric interpretation of (10.1). Every G -vector bundle on X is the pullback via $\eta: X \rightarrow X/G$ of one on X/G . If vector bundles on X/G are trivial, then every G -vector bundle on X is isomorphic to $X \times W$ for some k - G -module W .

(10.3) THEOREM. Assume that:

(a) The action of G on $X = \text{Spec}(A)$ is fix pointed.

(b) X^G is a local complete intersection in X . (This is automatic if $X^G \subset X_{\text{reg}}$, by Proposition (8.4).)

Then $\pi: X/G$ admits the structure of a G -vector bundle over X/G . As such,

$$X \cong \bigoplus_{\lambda \in \Lambda} (E_\lambda \otimes_k V_\lambda)$$

(vector bundle direct sum), where E_λ for each $\lambda \in \Lambda$ is a vector bundle over X/G with trivial G -action.

The first assertion follows the Normal Linearization Theorem (9.1), with X^G ($\cong X/G$) in the role of X_0 . The second assertion follows then from the beginning of §3.

(10.4) COROLLARY. Assume (a) and (b) of Theorem (10.2) as well as:

(c) All vector bundles on X/G are trivial. (This is automatic if all vector bundles on X are trivial; Remark (10.2).)

Then X is isomorphic as a G -scheme to $(X/G) \times W$, where W is a k - G -module on which the action of G is one fix pointed.

With the notation of Theorem (10.3) we have, for each $\lambda \in \Lambda$, $E_\lambda \cong (X/G) \times k^{n_\lambda}$ for some $n_\lambda \geq 0$, so $E_\lambda \otimes_k V_\lambda \cong (X/G) \times V_\lambda^{n_\lambda}$. It follows that $X \cong (X/G) \times W$, where $W = \bigoplus_\lambda V_\lambda^{n_\lambda}$. Clearly W/G must be a single point, so the origin is the unique closed G -orbit in W , whence the last assertion.

(10.5) COROLLARY. Suppose that the reductive group G acts on the affine space $X = \mathbf{A}_k^n$. If the action is fix pointed, then X is G -isomorphic to $Y \times \mathbf{A}_k^m$, where $Y = X^G \cong X/G$, and where G acts linearly and one fix pointedly on $\mathbf{A}_k^m$. The action of G on X can be linearized iff Y is isomorphic to $\mathbf{A}_k^{n-m}$; this is automatic whenever $n - m \leq 2$.

Of the assumptions (a), (b), (c) of Corollary (10.4), (a) has been assumed, (b) follows because X is regular (see Proposition (8.4)), and (c) is just the Quillen-Suslin Theorem ("Serre Conjecture") (see [Q and S]). The first assertion thus follows from Corollary (10.4) with $m = \dim_k(W)$. If the action of G can be linearized, then $Y = X^G$ must be isomorphic to a linear space, hence $Y \cong \mathbf{A}_k^{n-m}$. Conversely any isomorphism $Y \cong \mathbf{A}_k^{n-m}$ clearly leads to a linear action of G on $Y \times \mathbf{A}_k^m \cong \mathbf{A}_k^{n-m} \times \mathbf{A}_k^m \cong \mathbf{A}_k^n$. If $n - m \leq 2$, then $Y \cong \mathbf{A}_k^{n-m}$ by the Cancellation Theorem of Fujita, Miyanishi and Sugie (cf. [Ru]).

(10.6) COROLLARY. If the action of G on X is one fix pointed and the fixed point x is regular in X , then (X, x) is G -isomorphic to the tangent space action on $(T_x(X), 0)$.

This is (10.4) in the case when X/G is a point.

(10.7) COROLLARY. If the reductive group G acts on the affine scheme $X = \text{Spec}(A)$ with a dense orbit and a regular fixed point x , then (X, x) is G -isomorphic to the tangent space action on $(T_x(X), 0)$.

By Corollary (10.6) it suffices to check that the action is one fix pointed. The hypotheses clearly imply that $\{x\}$ is the unique closed orbit, so X/G is a point, and A^G is an artin local ring. If $A^G = k$, i.e. if A^G is reduced, then $\Delta_G(A) = m_x$, and the

action is one fix pointed, as required. Suppose, on the contrary, that $a \in A^G$ is nilpotent, $\neq 0$. Then $\text{ann}_A(a)$ is a G -stable ideal $\neq A$, hence contained in the unique maximal G -stable ideal m_x . Hence a remains $\neq 0$ in A_{m_x} . But this contradicts the assumed regularity of A_{m_x} .

(10.8) REMARKS. Corollary (10.7) is contained in Luna [L, III, §1, Corollaire 2]. Note that it furnishes an equivariant characterization of affine spaces, as varieties admitting a reductive group action with a dense orbit and a regular fixed point.

11. Examples of fix pointed actions. We consider a reductive group G acting on $X = \text{Spec}(A)$, with an eye to recognizing when the action is fix pointed.

(11.1) EXAMPLE. Suppose that G is diagonalizable, i.e. isomorphic to a subgroup of a torus. Consider the character group $G^v = \text{Hom}(G, \mathbf{G}_m)$, which we shall write additively; if $\chi \in G^v$ we write t^χ in place of $\chi(t)$. The action of G on X is equivalent to the G^v -graded ring structure,

$$A = \bigoplus_{\chi \in G^v} A_\chi,$$

where A_χ consists of semi-invariants of weight χ for the (translation) action of G on A . We have $A_0 = A^G$, and $\Delta_G(A) = \bigoplus_{\chi \neq 0} A_\chi$. In order for the action to be fix pointed, i.e. in order for $\Delta_G(A)$ to be an ideal, it is clearly necessary and sufficient that $A_\chi \cdot A_{-\chi} = 0$ for all $\chi \neq 0$ in G^v .

Suppose that A is an integral domain. Then $\Sigma \stackrel{\text{def}}{=} \{\chi \in G^v \mid A_\chi \neq 0\}$ is clearly a submonoid of G^v . In this case the action is fix pointed iff

$$\chi, -\chi \in \Sigma \Rightarrow \chi = 0,$$

i.e. Σ contains no nontrivial invertible elements. This is the condition which Kambayashi and Russell [KR] call “unmixedness” (when G is a torus). Theorem (10.3) above thus becomes, when G is a torus and X is regular, a theorem of Kambayashi and Russell [KR, Theorem (2.5)], describing A as a Σ -graded symmetric algebra over $A^G = A_0$. Their Theorem (3.4) is similarly subsumed in Corollary (10.5) above.

(11.2) PROPOSITION. Let $B \subset A$ be a finitely generated G -stable subalgebra, and let $C = A/J$ where J is a G -stable ideal of A . If the action of G on A is fix pointed, then the actions of G on B and C are fix pointed.

Indeed since Δ_G is an exact functor on a k - G -module we have $\Delta_G(B) = B \cap \Delta_G(A)$, an ideal of B , and $\Delta_G(C) = \text{the image mod } J \text{ of } \Delta_G(A)$, an ideal of C .

(11.3) PROPOSITION. Let $x \in X_{\text{reg}}$ be a regular fixed point of G , and let $T = T_x(X)$, the tangent space at x . If G acts fix pointedly on X , then G acts fix pointedly on T . The converse holds if A is an integral domain.

We can write $m_x = T^* \oplus m_x^2$, where T^* is a G -submodule of m_x , isomorphic to the dual of T . Let S denote the symmetric algebra $S_k(T^*)$, and $S_+ = T^* \cdot S$. The inclusion $T^* \subset A$ defines a G -equivariant homomorphism $S \rightarrow A$ which, since x is regular, induces an isomorphism from $\text{gr}_{S_+}(S) = S$ to $\text{gr}_{m_x}(A)$. Since $\bigcap_{n \geq 0} S_+^n = 0$ it

follows that $S \rightarrow A$ is injective, so G acts fix pointedly on S if it does so on A by Proposition (11.2). Suppose that G acts fix pointedly on S . Then, again by (11.2), it does so also on $S/S_+^n \cong A/m_x^n \stackrel{\text{def}}{=} A_n$. Thus $A_n = A_n^G \oplus \Delta_G(A_n)$ with $\Delta_G(A_n)$ an ideal. Passing to the m_x -adic completion $\hat{A} = \varprojlim_n A_n$ we have $\hat{A} = \hat{A}^G \oplus \widehat{\Delta_G(A)}$, with $\widehat{\Delta_G(A)}$ an ideal. If A is an integral domain then $\bigcap_{n \geq 0} m_x^n = 0$ so $A \rightarrow \hat{A}$ is injective, and hence $\Delta_G(A) = A \cap \widehat{\Delta_G(A)}$ is an ideal of A .

In view of (11.3) it is natural to ask when a linear action is fix pointed.

(11.4) PROPOSITION. *Let W be a finite dimensional k - G -module, $W = W^G \oplus \Delta_G(W)$. The following conditions are equivalent.*

- (a) *The action of G on W is fix pointed.*
- (b) *The action of G on $\Delta_G(W)$ is (one) fix pointed.*
- (c) $S_k(\Delta_G(W)^*)^G = k$.

We have

$$S_k(W^*) = S_k((W^G)^*) \otimes_k S_k(\Delta_G(W)^*)$$

and

$$\Delta_G(S_k(W^*)) = S_k((W^G)^*) \otimes_k \Delta_G(S_k(\Delta_G(W)^*)),$$

whence (a) $\Leftrightarrow$ (b) (since $\Delta_G(W)^G = \{0\}$). Since $\Delta_G^+(S_k(\Delta_G(W)^*)) = \Delta_G(W)^* \cdot S_k(\Delta_G(W)^*)$ we have (b) $\Leftrightarrow$ (c).

(11.5) PROPOSITION. *Let W be a finite dimensional k - G -module with $W^G = 0$. Assume that G is diagonalizable, and let $\chi_1, \dots, \chi_n \in G^n$ be the weights of G in W ($n = \dim W$). The action of G on W is (one) fix pointed iff*

$$(*) \quad \left. \begin{array}{l} \sum_{i=1}^n a_i \chi_i = 0 \\ (a_i \in \mathbf{Z}, a_i \geq 0) \end{array} \right\} \Rightarrow a_i = 0 \quad \text{for } i = 1, \dots, n.$$

Let $t_1, \dots, t_n$ be a basis of W^* with t_i semi-invariant of weight $-\chi_i$. If $a = (a_1, \dots, a_n) \in \mathbf{Z}^n$, $a_i \geq 0$, then $t^a = t_1^{a_1} \cdots t_n^{a_n}$ is semi-invariant of weight $-\sum_{i=1}^n a_i \chi_i$. Hence (*) is equivalent to the condition that $S_k(W^*)^G = k$, whence the proposition, by (11.4)(c).

(11.6) COROLLARY (BIALYNICKI - BIRULA). *If an n -dimensional torus G acts effectively on $X = \mathbf{A}_k^n$, then the action can be linearized.*

According to [B-B1, Theorem 1], G has a fixed point, which we may assume to be the origin. The tangent space representation at 0 is faithful, so its weights generate G^n , hence they are linearly independent so, by (11.5) and (11.3), the action of G on X is (one) fix pointed. The corollary thus follows from Corollary (10.6).

(11.7) PROPOSITION. *Let W be a finite dimensional k - G -module with $W^G = 0$. Assume that G is semisimple. The action of G on W is (one) fix pointed iff G has a dense orbit in W .*

If G has a dense orbit in W then $S_k(W^*)^G = k$, clearly, so the action is one fix pointed by (11.4)(c). Suppose, conversely, that $S_k(W^*)^G = k$. Then $S_k(W^*)^{G^0} = k$, where G^0 is the identity component of G . Since G^0 has no rational characters and $S_k(W^*)$ is factorial it follows from Luna [L1, III, §4], that the maximum dimension of a G^0 -orbit in W is $\dim W - \dim W/G = \dim W$, whence the existence of a dense G^0 -orbit in W .

(11.8) **REMARK.** For certain semisimple groups G , the k - G -modules with a dense orbit have been explicitly determined (cf. [Se]).

(11.9) **REMARK.** Let G be a reductive group acting on $X = \text{Spec}(A)$, with A an integral domain. Put $m = \max_{x \in X} \dim Gx$ and $X_{\max} = \{x \in X | \dim Gx = m\}$. Results of Rosenlicht [Ro] imply that $X_{\max}$ is open dense in X . The action is said to be *stable* (in the sense of Popov [Po]) if there is an open dense set of closed orbits, hence an open dense part of $X_{\max}$ consisting of closed orbits.

Consider the projection $\pi: X \rightarrow X/G$. Let $\xi \in X/G$, and suppose that $\pi^{-1}(\xi)$ is a closed m -dimensional orbit. It is disjoint from the G -stable closed set $X - X_{\max}$ so there is an $f \in A^G$ such that $f = 1$ on $\pi^{-1}(\xi)$ and $f = 0$ on $X - X_{\max}$. Hence $\pi(X - X_{\max})$ is contained in the zero locus of f in X/G , so $\xi \notin \pi(X - X_{\max})$. Thus the complement of $\pi^{-1}(\pi(X - X_{\max}))$ is an open nonempty, hence dense, set consisting of closed orbits. In conclusion, *the action is stable iff there is a single closed orbit in $X_{\max}$.*

Suppose further that G has no rational characters (e.g. that G is connected and semisimple) and that X is factorial. Then Popov [Po] has shown that the action is stable iff an open dense part of $X_{\max}$ consists of affine orbits. Thus the action is not stable iff no orbit in $X_{\max}$ is closed, and most orbits in $X_{\max}$ are nonaffine.

(11.10) **PROPOSITION.** *If $G = \text{SL}_2$ acts on $X = \text{Spec}(A)$ where A is factorial, and if the action is not stable, then the action is fix pointed.*

Define $h: k^x \rightarrow G$ and $u: k \rightarrow G$ by

$$h(a) = \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \quad \text{and} \quad u(b) = \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}.$$

Put $T = h(k^x)$, $T_n = \{h(a) | a^n = 1\}$, $U = u(k)$, $B = T \cdot U$, $B_n = T_n \cdot U$, and $N = \langle T, w \rangle$, $w = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. Up to conjugacy the closed subgroups H of G are: finite, B_n ($n \geq 1$), T , N , B , G . The corresponding quotients G/H are affine except for $G/B \cong \mathbf{P}^1$, and G/B_n , which are quasi-affine.

If $m = \max_{x \in X} \dim Gx$ is 3 the action is stable by Popov's result quoted above. Suppose that the action is not stable, hence $m \leq 2$. Let $x \in X$. Then $\dim Gx = 0$ or 2. If $\dim Gx = 2$ then, since Gx is not closed (by the discussion above), $\overline{Gx}$ must contain lower dimensional orbits, hence fixed points. It follows, as claimed, that the action is fix pointed (see (10.0)).

(11.11) **COROLLARY (PANYUSHEV [P]).** *If $G = \text{SL}_2$ acts on $X = \mathbf{A}^n$ and the action is not stable, then X is G -isomorphic to $(X/G) \times V$ for some k - G -module V .*

This follows from Proposition (11.10) and Corollary (10.5).

12. An equivariant approach to the Jacobian Conjecture. An endomorphism $\varphi: \mathbf{A}_{\mathbb{C}}^n \rightarrow \mathbf{A}_{\mathbb{C}}^n$, $\varphi(X) = (\varphi_1(X), \dots, \varphi_n(X))$, defines an endomorphism of the polynomial algebra $\mathbf{C}[X] = \mathbf{C}[X_1, \dots, X_n]$, $\bar{\varphi}: X_i \mapsto \varphi_i(X)$, sending $f(X)$ to $f(\varphi(X))$. The Jacobian matrix is denoted

$$\varphi'(X) = \begin{pmatrix} D_1\varphi_1 & \cdots & D_n\varphi_1 \\ \vdots & & \vdots \\ D_1\varphi_n & \cdots & D_n\varphi_n \end{pmatrix},$$

where $D_i = \partial/\partial X_i$. The Chain Rule takes the form

$$(1) \quad (\psi \circ \varphi)'(X) = \psi'(\varphi(X)) \cdot \varphi'(X).$$

If φ is invertible and $\psi = \varphi^{-1}$, i.e. $\psi(\varphi(X)) = X$, then (1) shows that $\varphi'(X)^{-1}$ is invertible, and

$$(2) \quad \varphi'(X)^{-1} = (\varphi^{-1})'(\varphi(X)).$$

The Jacobian Conjecture asserts, conversely, that if $\varphi'(X)$ is invertible, then φ is invertible (cf. [BCW]). Suppose that $\varphi'(X)$ is invertible. Modifying φ by a translation, if necessary, we may further assume that $\varphi(0) = 0$. Then at least φ has a local analytic inverse near 0, say $\varphi^{-1}(X) = (\varphi_1^{-1}(X), \dots, \varphi_n^{-1}(X))$ with each $\varphi_i^{-1}(X)$ in the ring $\mathbf{C}\{X\} = \mathbf{C}\{X_1, \dots, X_n\}$ of convergent power series. We can view $\bar{\varphi}: X_i \mapsto \varphi_i(X)$ as an automorphism of $\mathbf{C}\{X\}$, with inverse $\bar{\varphi}^{-1}: X_i \mapsto \varphi_i^{-1}(X)$.

Let D be a derivation of $\mathbf{C}\{X\}$. Then $\bar{\varphi} \circ D \circ \bar{\varphi}^{-1}$ is another such derivation, sending X_i to

$$\begin{aligned} [(\bar{\varphi} \circ D \circ \bar{\varphi}^{-1})(X_i)](X) &= D(\varphi_i^{-1})(\varphi(X)) \\ &= \left(\sum_{j=1}^n D_j(\varphi_i^{-1}) \cdot D(X_j) \right)(\varphi(X)) \\ &= \sum_{j=1}^n D_j(\varphi_i^{-1})(\varphi(X)) \cdot D(X_j)(\varphi(X)). \end{aligned}$$

Now by (2) above, $D_j(\varphi_i^{-1})(\varphi(X))$ is the (i, j) -entry of $\varphi'(X)^{-1}$, and hence, by the assumed invertibility of $\varphi'(X)$, is a polynomial. Supposing further that D is a polynomial derivation, i.e. that each $D(X_j)$ is a polynomial, then $D(X_j)(\varphi(X))$ is likewise a polynomial. In conclusion:

$$(3) \quad \text{If } \varphi: \mathbf{A}_{\mathbb{C}}^n \rightarrow \mathbf{A}_{\mathbb{C}}^n \text{ has invertible Jacobian } \varphi'(X), \text{ then for every polynomial derivation } D \text{ of } \mathbf{C}[X], \bar{\varphi} \circ D \circ \bar{\varphi}^{-1} \text{ (which is a priori only a derivation of } \mathbf{C}\{X\}) \text{ is again a polynomial derivation.}$$

Continuing to assume that $\varphi'(X)$ is invertible and that $\varphi(0) = 0$, consider now the group $G = \mathrm{SL}_n(\mathbf{C})$ with its natural (linear) action on $\mathbf{A}_{\mathbb{C}}^n$. This is a one fix pointed action with a dense orbit. We can view $\varphi^{-1}G\varphi$ as a group of germs of automorphisms near 0 of $\mathbf{A}_{\mathbb{C}}^n$, and $\bar{\varphi}\bar{G}\bar{\varphi}^{-1}$ as a group of automorphisms of $\mathbf{C}\{X\}$. Suppose that $\varphi^{-1}G\varphi$ actually consists of polynomial automorphisms, i.e. that $\bar{\varphi}\bar{G}\bar{\varphi}^{-1}$ stabilizes $\mathbf{C}[X]$. Then $\varphi^{-1}G\varphi$ defines a new action of G on $\mathbf{A}_{\mathbb{C}}^n$, and φ becomes a

G -morphism from A^n_C with the new action to A^n_C with the natural action. As such it is a G -linear approximation to the new action, in the sense of Remark (9.2)(4), and the Normal Linearization Theorem (9.1) implies then that φ is an isomorphism.

To find reasons for $\varphi^{-1}G\varphi$ to consist of polynomial automorphisms, consider the Lie algebra $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$ of G . It acts naturally as a Lie algebra of (linear) derivations of $\mathbb{C}[X]$. The differential of the new action, $\varphi^{-1}G\varphi$, of G , represents $\mathfrak{g}$ as the algebra $\bar{\varphi}\mathfrak{g}\bar{\varphi}^{-1}$ of derivations of $\mathbb{C}\{X\}$. However observation (3) above tells us that $\bar{\varphi}\mathfrak{g}\bar{\varphi}^{-1}$ is in fact again a Lie algebra of polynomial derivations. This gives a representation of the Lie algebra $\mathfrak{g}$ as derivations of $\mathbb{C}[X]$ which, on the subalgebra $\mathbb{C}[\varphi] = \mathbb{C}[\varphi_1, \dots, \varphi_n]$, looks like the natural representation in the φ -coordinates. Then $\bar{\varphi}G\bar{\varphi}^{-1}$ is the exponential of this polynomial representation of $\mathfrak{g}$. In order to show then that $\bar{\varphi}G\bar{\varphi}^{-1}$ is polynomial, it suffices to show that the action of $\bar{\varphi}\mathfrak{g}\bar{\varphi}^{-1}$ on $\mathbb{C}[X]$ is locally finite, i.e. that $\mathbb{C}[X]$ is a union of finite dimensional subspaces stable under $\bar{\varphi}\mathfrak{g}\bar{\varphi}^{-1}$. For this will then imply that the nilpotent elements tX_iD_j ($i \neq j$) in $\mathfrak{g}$ yield locally nilpotent derivations $\bar{\varphi} \cdot tX_iD_j \cdot \bar{\varphi}^{-1}$ of $\mathbb{C}[X]$, and hence polynomial automorphisms $\exp(\bar{\varphi} \cdot tX_iD_j \cdot \bar{\varphi}^{-1}) = \bar{\varphi} \cdot \exp(tX_iD_j) \cdot \bar{\varphi}^{-1}$. The elementary matrices $\exp(tX_iD_j)$ generate $G = \mathrm{SL}_n(\mathbb{C})$, so we conclude that $\varphi^{-1}G\varphi$ is polynomial, as desired.

In conclusion, the Jacobian Conjecture (that φ is an isomorphism) would follow if one could show that the Lie algebra $\bar{\varphi}\mathfrak{sl}_n(\mathbb{C})\bar{\varphi}^{-1}$ of derivations of $\mathbb{C}[X]$ acts locally finitely on $\mathbb{C}[X]$. Hopefully the (well-studied) representation theory of $\mathfrak{sl}_n(\mathbb{C})$ may be of some help in this regard.

REFERENCES

- [BCW] H. Bass, E. H. Connell and D. Wright, *The Jacobian Conjecture: Reduction of degree and formal expansion of the inverse*, Bull. Amer. Math. Soc. (N.S.) **7** (1982), 287–330.
- [B-B1] A. Bialynicki-Birula, *Remarks on the action of an algebraic torus on k^n* , Bull. Acad. Polon. Sci. Ser. Math. Astronom. Phys. **14** (1966), 177–181.
- [B-B2] ———, *Remarks on the action of an algebraic torus on k^n* . II, Bull. Acad. Polon. Sci. Ser. Math. Astronom. Phys. **15** (1967), 123–125.
- [NB1] N. Bourbaki, *Algèbre commutative*, Chapitres 1 et 2, Hermann, Paris, 1961.
- [NB2] ———, *Algèbre commutative*, Chapitres 8 et 9, Masson, Paris, 1983.
- [EGA] A. Grothendieck, *Éléments de géométrie algébrique*. IV (Seconde partie), Publ. Inst. Hautes Etude Sci. No. 24, 1965.
- [J] H. W. E. Jung, *Über ganze birationale Transformationen der Ebene*, J. Reine Angew. Math. **184** (1942), 161–174.
- [K] T. Kambayashi, *Automorphism group of a polynomial ring and algebraic group action on an affine space*, J. Algebra **60** (1979), 439–451.
- [KR] T. Kambayashi and P. Russell, *On linearizing algebraic torus actions*, J. Pure Appl. Algebra **23** (1982), 243–250.
- [KP] H. Kraft, *Algebraic groups, invariants and representations*, International Conference, Trento, Italy, 1981, pp. 1.6–6.6.
- [L] D. Luna, *Slices étales*, Bull. Soc. Math. France Mém. **33** (1973), 81–105.
- [N] M. Nagata, *Lectures on the fourteenth problem of Hilbert*, Tata Inst. of Fundamental Research, Bombay, 1965.
- [P] D. I. Panyushev, *Semisimple automorphism groups of four dimensional affine space*, Izv. Akad. Nauk SSSR **47** (1983), 881–894. (Russian)
- [Po] V. L. Popov, *On the stability of the action of an algebraic group on an algebraic variety*, Math. USSR-Izv. **6** (1972), 367–379.
- [Q] D. Quillen, *Projective modules over polynomial rings*, Invent. Math. **36** (1976), 167–171.

- [R1] R. Richardson, *The conjugating representation of a semisimple algebraic group*, Invent. Math. **54** (1979), 229–245.
- [R2] ———, *An application of the Serre conjecture to semisimple algebraic groups*, Algebra (Carbondale 1980), Lecture Notes in Math., vol. 848, Springer-Verlag, Berlin and New York, pp. 141–151.
- [Ro] M. Rosenlicht, *Some basic theorems on algebraic groups*, Amer. J. Math. **78** (1956), 401–443.
- [Ru] P. Russell, *On affine-ruled rational surfaces*, Math. Ann. **255** (1981), 287–302.
- [Se] F. J. Servidio, *Prehomogeneous vector spaces and varieties*, Trans. Amer. Math. Soc. **176** (1973), 421–444.
- [Sh] I. Shafarevič, *Some infinite dimensional algebraic groups*, Rend. Mat. (5) **25** (1966), 208–212.
- [Sp] T. A. Springer, *Invariant theory*, Lecture Notes in Math., vol. 585, Springer-Verlag, Berlin and New York, 1977.
- [S] A. Suslin, *Projective modules over a polynomial ring*, Dokl. Akad. Nauk SSSR **26** (1976). (Russian)

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NEW YORK 10027

DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF NEW YORK AT ALBANY, ALBANY, NEW YORK 12222