AN ASYMPTOTIC FORMULA FOR HYPO-ANALYTIC PSEUDODIFFERENTIAL OPERATORS

S. BERHANU

ABSTRACT. An asymptotic expansion formula for hypo-analytic pseudodifferential operators is proved and applications are given.

INTRODUCTION

In [2] we introduced hypo-analytic pseudodifferential operators that are naturally associated with the hypo-analytic structures of [1]. In this paper we establish an asymptotic formula for these operators. Such an expansion is essential in several applications. It allows us to define, in a natural way, the symbol of a hypo-analytic pseudodifferential operator, as well as the symbols of the adjoint, transpose and composition of operators. The paper is organized as follows. In Chapter I we discuss and develop the asymptotic formula. Chapter II applies this formula to two results.

Acknowledgment. It is a pleasure to express my thanks to Professor F. Trèves for many stimulating discussions.

1. ASYMPTOTIC EXPANSION

1. Hypo-analytic structures. We will deal with structures which are a special case of the hypo-analytic structures introduced by Baouendi, Chang and Trèves in [1]. We shall summarize the relevant concepts here. Let Ω be a C^∞ manifold of dimension m . A hypo-analytic structure of maximal dimension on Ω is the data of an open covering (U_α) of Ω and for each index α , of m C^∞ functions $Z_\alpha^1, \dots, Z_\alpha^m$ satisfying the following two conditions:

- (i) $dZ_\alpha^1, \dots, dZ_\alpha^m$ are linearly independent at each point of U_α ;
- (ii) if $U_\alpha \cap U_\beta \neq \emptyset$, there are open neighborhoods $\mathcal{O}_\alpha$ of $Z_\alpha(U_\alpha \cap U_\beta)$ and $\mathcal{O}_\beta$ of $Z_\beta(U_\alpha \cap U_\beta)$ and a holomorphic map F_β^α of $\mathcal{O}_\alpha$ onto $\mathcal{O}_\beta$

such that

$$Z_\beta = F_\beta^\alpha \circ Z_\alpha \quad \text{on } U_\alpha \cap U_\beta.$$

We will use the notation $Z_\alpha = (Z_\alpha^1, \dots, Z_\alpha^m): U_\alpha \rightarrow C^m$. A distribution h defined in an open neighborhood of a point p_0 of Ω is hypo-analytic at p_0

Received by the editors August 14, 1989.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 35A20; Secondary 35S99.

©1990 American Mathematical Society
0002-9947/90 \$1.00 + \$.25 per page

if there is a chart (U_α, Z_α) of the above type whose domain contains p_0 and a holomorphic function $\tilde{h}$ defined on an open neighborhood of $Z_\alpha(p_0)$ in C^m such that $h = \tilde{h} \circ Z_\alpha$ in a neighborhood of p_0 . By a hypo-analytic local chart we mean an $m+1$ -tuple $(U, Z^1, \dots, Z^m)$ [abbreviated (U, Z)] consisting of an open subset U of Ω and of m hypo-analytic functions whose differentials are linearly independent at every point of U .

We will now reason in a hypo-analytic local chart (U, Z) of Ω . Assume that the open set U has been contracted sufficiently so that the mapping $Z = (Z^1, \dots, Z^m) : U \rightarrow C^m$ is a diffeomorphism of U onto $Z(U)$ and that U is the domain of local coordinates x_j ($1 \leq j \leq m$) all vanishing at a "central point" which will be denoted by 0. We will suppose $Z(0) = 0$ and denote by Z_x the Jacobian matrix of the Z^j with respect to the x^k . Substitution of $Z_x(0)^{-1}Z(x)$ for $Z(x)$ will allow us to assume that $Z_x(0) =$ the identity matrix. Therefore the real part of the Z^j ($j = 1, \dots, m$) can serve as coordinates and in these new coordinates

$$Z^j = x^j + \sqrt{-1}\phi^j(x), \quad j = 1, \dots, m,$$

where $\phi = (\phi^1, \dots, \phi^m)$ is real valued with 0 differential at the origin.

Moreover, the functions Z^j are selected so that all the derivatives of order two of the ϕ^j vanish at the origin. Indeed if this is not already so it suffices to replace each Z^j by

$$Z^j - \frac{\sqrt{-1}}{2} \sum_k \sum_l \frac{\partial^2 \phi^j}{\partial x^k \partial x^l}(0) Z^k Z^l.$$

We will use $\check{Z}_x$ to denote the transpose of the inverse of the matrix Z_x . Since the first and second derivatives of all the ϕ^j are zero at the origin, after contracting U if necessary, we can find a number c , $0 < c < 1$ such that for all x, y in U and for all ξ in R_m

$$\begin{aligned} |\Im \check{Z}_x(x)\xi| &\leq c |\Re \check{Z}_x(x)\xi| \quad \text{and} \\ (1.1) \quad \Re \{ \sqrt{-1} \check{Z}_x(x)\xi \cdot (Z(x) - Z(y)) - \langle \check{Z}_x(x)\xi, (Z(x) - Z(y))^2 \rangle \} \\ &\leq -c |\xi| |Z(x) - Z(y)|^2, \\ &\quad \text{where } \langle \zeta \rangle = (\zeta_1^2 + \dots + \zeta_m^2)^{\frac{1}{2}} \text{ for } |\Im \zeta| < |\Re \zeta|. \end{aligned}$$

2. Hypo-analytic pseudodifferential operators. We will continue to work in the chart (U, Z) of §1. Our aim now is to briefly describe the hypo-analytic pseudodifferential operators.

Definition 2.1. Let d be a real number. We denote by $\tilde{S}^d(U, U)$ the space of holomorphic functions $\tilde{a}(z, w, \theta)$ in a product set $\mathcal{O} \times \mathcal{O} \times \mathcal{C}$ with $\mathcal{O}$ an open neighborhood of $Z(U)$, and $\mathcal{C}$ an open cone in $C_m \setminus \{0\}$ containing $R_m \setminus \{0\}$ which have the following property :

Given any compact subset K of $\mathcal{O}$ and any closed cone $\mathcal{E}' \subset \mathcal{E}$ whose interior contains $R_m \setminus \{0\}$, there is a constant $r > 0$ such that for all z, w in K and all θ in $\mathcal{E}'$, we have

$$|\tilde{a}(z, w, \theta)| \leq r(1 + |\theta|)^d.$$

Definition 2.2. We say that a C^∞ function $a(x, y, \theta)$ in $U \times U \times R_m$ is a hypo-analytic amplitude of degree d and we write $a \in S^d(U, U)$ if there is $\tilde{a} \in \tilde{S}^d(U, U)$ such that

$$a(x, y, \theta) = \tilde{a}(Z(x), Z(y), \theta), \quad \text{for all } x \text{ in } U, y \text{ in } U, 0 \neq \theta \in R_m.$$

Let $a(x, y, \theta) = \tilde{a}(Z(x), Z(y), \theta)$ be a hypo-analytic amplitude of degree $d \in \mathbb{R}$ in $U \times U$. For any $\varepsilon > 0$ and $u \in C_c^0(U)$ we define the linear operator

$$(2.3) \quad A^\varepsilon u(x) = \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \int_U \int_{R_m} \exp(\sqrt{-1}\xi \cdot (Z(x) - Z(y) - \varepsilon|\xi|^2)) \cdot a(x, y, \xi) u(y) dZ(y) d\xi$$

We contract U sufficiently so that for every $x, y \in U$ and $\xi \in R_m$ the point $\tilde{Z}_x(x)\xi + \sqrt{-1}\langle \tilde{Z}_x(x)\xi \rangle (Z(x) - Z(y))$ will remain in the cone in which $a(x, y, \cdot)$ is defined. We observe that each $A^\varepsilon u$ is a hypo-analytic function. The results of [2] may be consolidated into:

Theorem 2.1. When $\varepsilon \rightarrow 0$, A^ε converges to a continuous linear operator $A: E'(U) \rightarrow \mathcal{D}(U)$ which maps $C_c^\infty(U)$ into $C_c^\infty(U)$ continuously. If u is hypo-analytic at 0 then Au is hypo-analytic at 0.

The first part of the theorem is proved by first deforming the path of ξ -integration from R_m to the image of R_m under the map

$$\xi \rightarrow \zeta(\xi) = \tilde{Z}_x(x)(\xi) + \sqrt{-1}\langle \tilde{Z}_x(x)\xi \rangle (Z(x) - Z(y)).$$

The second inequality in (1.1) will then force the exponential term in (2.3) to be bounded. The integral can then be treated as an oscillatory integral.

Following [2] we will call A a hypo-analytic pseudodifferential operator. When $Z(x) = x$ this specializes to the usual analytic pseudodifferential operator.

3. Formal hypo-analytic amplitudes. In this section (U, Z) will be as in §2. Our aim is to establish an asymptotic expansion formula for hypo-analytic amplitudes.

Fix a neighborhood $\mathcal{O}$ of $Z(U)$ in C^m , a cone $\mathcal{E}$ in $C^m \setminus \{0\}$ and let $R_0(z, w)$ be a positive continuous function on $\mathcal{O} \times \mathcal{O}$. For each $j = 0, 1, 2, \dots$ let $k_j(z, w, \theta)$ be a holomorphic function in the set

$$\{(z, w, \theta) \in \mathcal{O} \times \mathcal{O} \times \mathcal{E}; |\theta| > R_0(z, w) \sup(j, 1)\}.$$

Set $k_j(x, y, \theta) = \tilde{k}_j(Z(x), Z(y), \theta)$.

Definition 3.1. We will say that the series $\sum_{j=0}^{\infty} k_j(x, y, \theta)$ defines a formal hypo-analytic amplitude of degree d if there exists a continuous function $c_0(z, w) > 0$ on $\mathcal{O} \times \mathcal{O}$ such that for all (z, w) in $\mathcal{O} \times \mathcal{O}$ and all θ in $\mathcal{C}$, $|\theta| > R_0(z, w) \sup(j, 1)$,

$$|\tilde{k}_j(z, w, \theta)| \leq C_0(z, w)^{j+1} j! |\theta|^{d-j}.$$

We now show how to construct a true hypo-analytic amplitude from the formal one given above. We will work in a compact set $K \subseteq U$ and a relatively compact neighborhood $\mathcal{O}_K$ of $Z(K)$ in $\mathcal{O}$. This enables us to replace the functions $C_0(z, w)$ and $R_0(z, w)$ of the above definition by constants C_0 and R_0 . We will also assume that the cone $\mathcal{C}$ has been shrunk to satisfy: for some $\delta > 0$, whenever $\theta = \xi + \sqrt{-1}\eta \in \mathcal{C}$, then $\delta|\theta| \leq |\xi|$. Let $R > \max(R_0, C_0)$.

We will use a sequence of smooth cutoff functions $\phi_j(\xi)$ having the following properties:

$$0 \leq \phi_j(\xi) \text{ for all } \xi, \quad \text{and} \quad \phi_j(\xi) = 0 \text{ in } |\xi| < 2R \sup(j, 1),$$

$$\phi_j(\xi) = 1 \quad \text{if } |\xi| > 3R \sup(j, 1); \quad |D^\alpha \phi_j| \leq \left(\frac{C}{R}\right)^{|\alpha|} \quad \text{if } |\alpha| \leq 2j.$$

See [8] for the construction of such cutoffs. Define

$$\tilde{k}(z, w, \theta) = \sum_{j=0}^{\infty} \phi_j(\xi) \tilde{k}_j(z, w, \theta)$$

for $(z, w) \in \mathcal{O}_K \times \mathcal{O}_K$ and $\theta = \xi + \sqrt{-1}\eta \in \mathcal{C}$. $\tilde{k}$ is a C^∞ function of (z, w, θ) holomorphic in (z, w) . $\tilde{k}$ satisfies the following estimates:

$$\begin{aligned} |\tilde{k}(z, w, \theta)| &\leq \sum_{0 \leq j < d} |\tilde{k}_j(z, w, \theta)| + \sum_{j \geq d} \phi_j(\xi) |k_j(z, w, \theta)| \\ &\leq \sum_{0 \leq j < d} |\tilde{k}_j(z, w, \theta)| + \sum_{j \geq d} \phi_j(\xi) c_0^{j+1} j! |\theta|^{d-j} \\ &\leq \sum_{0 \leq j < d} |\tilde{k}_j(z, w, \theta)| + \sum_{j \geq d} \phi_j(\xi) c_0^{j+1} j! |\xi|^{d-j} \end{aligned}$$

Since for $j \geq d$ the j th term lives on the set $\{\xi : |\xi| \geq 2Rj\}$, the latter

$$\begin{aligned} &\leq \sum_{0 \leq j < d} |\tilde{k}_j(z, w, \theta)| + |\xi|^d \sum_{j \geq d} c_0^{j+1} j! \left(\frac{1}{2Rj}\right)^j \\ &\leq \sum_{0 \leq j < d} |\tilde{k}_j(z, w, \theta)| + \text{constant } |\xi|^d \\ &\leq \text{constant } |\theta|^d \end{aligned}$$

$$\begin{aligned}
\bar{\partial}_\theta \tilde{k}(z, w, \theta) &\leq \sum_{j=0}^{\infty} |\bar{\partial}_\theta \phi_j(\xi)| \tilde{k}_j(z, w, \theta) \\
&\leq \left(\sum_{j=0}^{\infty} |\bar{\partial}_\theta \phi_j(\xi)| c_0^{j+1} \frac{j!}{|\xi|^j} \right) \\
&\leq \delta^d |\xi|^d \left(\sum_{j=0}^{\infty} |\bar{\partial}_\theta \phi_j(\xi)| c_0^{j+1} \frac{j!}{|\xi|^j} \right)
\end{aligned}$$

We now use the fact that $\bar{\partial}_\theta \phi_j(\xi)$ lives in the set $\{\xi : 2Rj \leq |\xi| \leq 3Rj\}$;

$$\leq \text{constant } |\xi|^d \left(\sum_{j=0}^{\infty} c_0^{j+1} j! \left(\frac{1}{2Rj} \right)^j \right)$$

Since $j!/j^j \leq e^{-j}$, the latter $\leq \text{constant } |\xi|^d \sum_{j=0}^{\infty} \left(\frac{c_0}{2R} \right)^j e^{-j}$.

Recalling that $2Rj \leq |\xi| \leq 3Rj$, we get

$$\begin{aligned}
&\leq \text{constant } |\xi|^d \sum_{j=0}^{\infty} \left(\frac{c_0}{2R} \right)^j e^{-\frac{|\xi|}{3R}} \\
&\leq \text{constant } e^{-\frac{|\xi|}{4R}} \\
&\leq \text{constant } e^{-\frac{\delta}{4R} |\theta|}
\end{aligned}$$

Thus for $(z, w, \theta) \in \mathcal{O}_K \times \mathcal{O}_K \times \mathcal{C}$, we have: $|\tilde{k}(z, w, \theta)| \leq \text{const. } |\theta|^d$ and $|\bar{\partial}_\theta \tilde{k}(z, w, \theta)| \leq \text{const. } e^{-\frac{\delta}{4R} |\theta|}$.

We may assume that the shape of $\mathcal{C}$ has been modified to allow us to solve the Cauchy-Riemann equations in $\mathcal{C}$ (see [5]) $\bar{\partial}_\theta \tilde{k}_1 = \bar{\partial}_\theta \tilde{k}$ in such a way that the solution $\tilde{k}_1$ is holomorphic with respect to (z, w) in $\mathcal{O}_K \times \mathcal{O}_K$ and the following estimate holds on sets of the kind $K_1 \times K_2 \times \mathcal{C}_1(K_1, K_2 \subset \subset \mathcal{O}_K)$ and $\mathcal{C}_1$ a cone whose closure is contained in $\mathcal{C}$:

$$|\tilde{k}_1(z, w, \theta)| \leq \text{const. } e^{-\frac{\delta}{4R} |\theta|}.$$

Define then $h = \tilde{k} - \tilde{k}_1$. We now have, in $\mathcal{O}_K \times \mathcal{O}_K \times \mathcal{C}_1$, $\bar{\partial}_\theta \tilde{h} = 0$ and $\tilde{k} - \tilde{h}$ decays exponentially as $|\theta| \rightarrow \infty$ (uniformly, provided (z, w, θ) stays in sets like $K_1 \times K_2 \times \mathcal{C}_1$ as above).

This decay together with Theorem 2.1 of §2 imply that if for $u \in \mathcal{E}'(U)$, U sufficiently small, we define

$$\begin{aligned}
\text{op } \tilde{k}^\varepsilon u(x) &= \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \int_U \int_{R_m} e^{\sqrt{-1}\xi \cdot (Z(x) - Z(y)) - \varepsilon |\xi|^2} \\
&\quad \cdot \tilde{k}(Z(x), Z(y), \xi) u(y) dZ(y) d\xi
\end{aligned}$$

then as $\varepsilon \rightarrow 0^+$, $\text{op } \tilde{k}^\varepsilon$ will converge to an operator $\text{op } \tilde{k}$ having the properties in Theorem 2.1, §2. Moreover, for any $u \in \mathcal{E}'(U)$, $\text{op } \tilde{k}u - \text{op } \tilde{h}u$ is a hypo-analytic function. We will therefore replace $\tilde{k}$ by the hypo-analytic amplitude

$\tilde{h}$ and think of $\tilde{h}$ as being the true amplitude constructed from the formal one given by $\sum_{j=0}^{\infty} k_j(x, y, \theta)$.

4. Asymptotic expansion. Let $k(x, y, \theta)$ be a hypo-analytic amplitude of degree d say $k(x, y, \theta) = \tilde{k}(Z(x), Z(y), \theta)$ where $\tilde{k}$ is holomorphic in $\mathcal{O} \times \mathcal{O} \times \mathcal{C}$, $\mathcal{O}$ and $\mathcal{C}$ are as in §1. For each $j = 1, \dots, m$, let N_j denote the vector field $N_j Z^k = -\sqrt{-1} \delta_j^k$.

If $K \subset U$ is any compact subset, by Cauchy's inequality we have $c > 0$ such that:

$$\left| \frac{1}{\alpha!} \partial_{\xi}^{\alpha} N_y^{\alpha} k(x, x, \xi) \right| \leq c^{|\alpha|+1} \alpha! (1 + |\xi|)^{d-|\alpha|}$$

for $x \in K$, $\xi \in R_m$.

Thus if we define

$$k_j(x, \xi) = \sum_{|\alpha|=j} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} N_y^{\alpha} k(x, x, \xi)$$

then $\sum_{j=0}^{\infty} k_j(x, \xi)$ can be thought of as a formal hypo-analytic symbol. Let $(\phi_j)_j$ be the cutoff functions of the previous section. If U' is any relatively compact subset of U , we can form a true symbol by setting

$$k(x, \xi) = \sum_{j=0}^{\infty} k_j(x, \xi) \phi_j(\xi)$$

We then have two operators $\text{op } k(x, y, \xi)$ and $\text{op } \tilde{k}(x, \xi): \mathcal{E}'(U') \rightarrow D'(U')$ where for $u \in \mathcal{E}'(U')$,

$$\text{op } \tilde{k} u(x) = \lim_{\varepsilon \rightarrow 0^+} \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \int \int e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \xi - \varepsilon |\xi|^2} k(x, \xi) u(y) dZ(y) d\xi$$

and

$$\text{op } k u(x) = \lim_{\varepsilon \rightarrow 0^+} \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \int \int e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \xi - \varepsilon |\xi|^2} k(x, y, \xi) u(y) dZ(y) d\xi$$

The next theorem proves that if U' is small enough, modulo a hypo-analytic regularizing operator, $\text{op } k = \text{op } \tilde{k}$.

Theorem 4.1. *If the neighborhood U' is sufficiently small, $\text{op } k \equiv \text{op } \tilde{k}$ in the sense that for any $u \in \mathcal{D}'(U')$, $\text{op } k u - \text{op } \tilde{k} u$ is a hypo-analytic function.*

Proof. Assume U' is an open ball centered at 0, its size to be determined later. We first take $u \in C_c^0(U')$. The theorem will be proved by first establishing:

- (i) $(\text{op } k - \text{op } \tilde{k})u$ is in $C^\infty(U')$, and
- (ii) There exists $c > 0$ such that for all $\alpha \in Z_m^+$,

$$|M^\alpha (\text{op } k - \text{op } \tilde{k})u(x)| \leq c^{|\alpha|+1} \alpha! \quad \text{where } M_j = \sqrt{-1} N_j \\ \text{for each } j = 1, \dots, m.$$

Moreover, since each $(\text{op } k - K_N)u$ is hypo-analytic in a common domain, for example some neighborhood of the compact set $\bar{U}'$, we can find a constant $\tilde{c}_1 > 0$ independent of N such that for all $\alpha \in Z_m^+$,

$$|M^\alpha(\text{op } k - K_N)u(x)| \leq \tilde{c}_1^{|\alpha|+1} c_1^{N+1} \alpha!$$

(B) Estimate of $M^\alpha(S_N u)$. Write

$$s_N(x, y, \xi) = \phi_{N+1}(\xi) \sum_{|\alpha|=N+1} D_\xi^\alpha k_\alpha(x, y, \xi) = \phi_{N+1}(\xi) \tilde{s}_N(x, y, \xi).$$

For $|\alpha| = N + 1$, we have

$$\left| \frac{D_\xi^\alpha k_\alpha(x, y, \xi)}{\alpha!} \right| \leq c^{|\alpha|} \alpha! (1 + |\xi|)^{d-N-1}.$$

It follows that $|\tilde{s}_N(x, y, \xi)| \leq c_2^{N+1} N! (1 + |\xi|)^{d-N-1}$ for some $c_2 > 0$.

Let

$$I_N^\varepsilon(x) = \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \int \int e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \xi - \varepsilon |\xi|^2} \phi_{N+1}(\xi) \tilde{s}_N(x, y, \xi) u(y) dZ(y) d\xi.$$

We note that $s_N u(x) = \lim_{\varepsilon \rightarrow 0^+} I_N^\varepsilon(x)$.

We will deform the path of ξ -integration from R_m to the image of R_m under the map

$$\xi \rightarrow \theta(\xi) = \phi_{2N}(\xi) \zeta(\xi) + (1 - \phi_{2N}(\xi)) \xi$$

where

$$\zeta(\xi) = \check{Z}_x(x) \xi + \sqrt{-1} \langle \check{Z}_x(x) \xi | (Z(x) - Z(y)) \rangle.$$

The deformation is allowed since it takes place in a region where $\phi_{N+1}(\xi)$ is analytic.

We have

$$\theta(\xi) = \begin{cases} \xi, & \text{for } |\xi| \leq 4RN, \\ \zeta(\xi), & \text{for } |\xi| \geq 6RN. \end{cases}$$

$$|M^\alpha(I_N^\varepsilon(x))| \leq \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \sum_{\beta < \alpha} \binom{\alpha}{\beta} \left| \int \int \xi^{\alpha-\beta} e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \xi - \varepsilon |\xi|^2} \cdot \phi_{N+1}(\xi) M^\beta \tilde{s}_N(x, y, \xi) u(y) dZ(y) d\xi \right|$$

We use the above contour and pass to the limit to get:

$$\begin{aligned} |(M^\alpha s_N u)(x)| &\leq \left| \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \int_{2R(N+1) \leq |\xi| \leq 6RN} \int (\theta(\xi))^{\alpha-\beta} \right. \\ &\quad \cdot e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \theta(\xi)} \phi_{N+1}(\xi) M^\beta \tilde{s}_N(x, y, \xi) u(y) d\theta dZ(y) \\ &\quad + \int_{|\xi| \geq 6RN} \int (\zeta(\xi))^{\alpha-\beta} e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \zeta(\xi)} \phi_{N+1}(\xi) \\ &\quad \left. \cdot M^\beta \tilde{s}_N(x, y, \zeta(\xi)) u(y) dZ(y) d\xi \right| \end{aligned}$$

We recall that the exponential in the second integral is bounded (§1, (1.1)). By hypo-analyticity we get $\tilde{c}_3 > 0$ such that

$$\forall \beta, |M^\beta \tilde{s}_N(x, y, \xi)| \leq c_3^{|\beta|+1} \beta! c_2^{N+1} N! (1 + |\xi|)^{d-N-1}.$$

These observations imply that

$$\begin{aligned} |M^\alpha s_N u(x)| \leq \text{const} \cdot & \left(\sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \int_{2R(N+1) \leq |\xi| \leq 6RN} |\xi|^{\alpha-\beta} e^{A|\xi|} c_3^{N+|\beta|+2} \right. \\ & \cdot \beta! N! (1 + |\xi|)^{d-N-1} d\xi + \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \\ & \left. \cdot \int_{6RN \leq |\xi|} |\xi|^{\alpha-\beta} c_3^{N+|\beta|+2} \beta! N! (1 + |\xi|)^{d-N-1} d\xi \right) \end{aligned}$$

for some $c_3 \geq \max(\tilde{c}_3, c_2)$. Hence, after modifying c_3 if necessary, we get

$$\begin{aligned} |M^\alpha s_N u(x)| &\leq \alpha! \left(\sum_{\beta \leq \alpha} \frac{1}{(\alpha - \beta)!} \int_{2R(N+1) \leq |\xi|} (1 + |\xi|)^{|\alpha - \beta| + d - N - 1} N! d\xi \right) c_3^N \\ &\leq \alpha! c_3^N \left(\sum_{\beta \leq \alpha} \frac{1}{(\alpha - \beta)!} \left(\frac{1}{1 + 2RN} \right)^{N - |\alpha - \beta| - d - m + 1} N! \right) \\ &\leq \alpha! c_3^N \left(\sum_{\beta \leq \alpha} \frac{1}{(\alpha - \beta)!} (1 + 2RN)^{|\alpha - \beta| + d + m - 1} \right) \left(\frac{1}{2R} \right)^N \frac{N!}{N^N} \\ &\leq \alpha! c_3^N \left(\sum_{\beta \leq \alpha} \frac{1}{(\alpha - \beta)!} (1 + 2RN)^{|\alpha - \beta| + d + m - 1} \right) \left(\frac{1}{2Re} \right)^N Ne \\ &\leq \alpha! c_3^N \left(\sum_{\beta \leq \alpha} \frac{1}{(\alpha - \beta)!} (|\alpha - \beta| + d + m - 1)! e^{1 + 2RN} \right) \left(\frac{1}{2Re} \right)^N Ne \\ &\leq \alpha! c_3^N \left(\sum_{\beta \leq \alpha} \frac{1}{(\alpha - \beta)!} (|\alpha - \beta| + d + m - 1)! \right) \left(\frac{e^{2R}}{2Re} \right)^N Ne^2. \end{aligned}$$

Using the inequality: $(k + l)! \leq 2^{k+l} k! l!$ for any positive integers k and l , the latter is dominated by

$$\alpha! c_3^N \left(\sum_{\beta \leq \alpha} \frac{1}{(\alpha - \beta)!} |\alpha - \beta|! \right) 2^{|\alpha| + d + m - 1} \left(\frac{e^{2R}}{2Re} \right)^N Ne^2.$$

For $|\alpha| \leq N$, we can find another constant which we will still call c_3 such that the above quantity $\leq \alpha! c_3^N$.

(C) **Estimate of $M^\alpha(\text{op } \tilde{k} - \tilde{K}_N)u$.** Let

$$J^\varepsilon u(x) = \left(\frac{1}{4\pi^3}\right)^{\frac{m}{2}} \int \int e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \xi - \varepsilon|\xi|^2} \cdot \left(\sum_{j>N} \phi_j(\xi) k_j(x, \xi)\right) u(y) dZ(y) d\xi.$$

For each $j > N$, we will use the contour

$$\theta_j(\xi) = \phi_{2j}(\xi)\zeta(\xi) + (1 - \phi_{2j}(\xi))\xi = \begin{cases} \xi, & \text{when } |\xi| \leq 4Rj, \\ \zeta(\xi), & \text{when } |\xi| \geq 6Rj. \end{cases}$$

In the quantity

$$M^\alpha(J^\varepsilon u)(x) = \left(\frac{1}{4\pi^3}\right)^{\frac{m}{2}} \sum_{j>N} \left[\sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \int \int \xi^{\alpha-\beta} e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \xi - \varepsilon|\xi|^2} \cdot \phi_j(\xi) M^\beta k_j(x, \xi) u(y) dZ d\xi \right]$$

we use the contours θ^j in each term of the sum and take limits to get

$$M^\alpha(\text{op } \tilde{k} - \tilde{K}_N)u(x) = \sum_{j>N} (I_1^j(x) + I_2^j(x))$$

where

$$I_1^j(x) = \left(\frac{1}{4\pi^3}\right)^{\frac{m}{2}} \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \int_{2Rj \leq |\xi| \leq 6Rj} \int \theta^j(\xi) e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \theta^j(\xi)} \cdot \phi_j(\xi) M^\beta k_j(x, \theta^j(\xi)) u(y) dZ d\theta^j$$

while $I_2^j(x)$ is a similar expression except that the integration in ξ is carried out over the region $\{\xi : |\xi| \geq 6Rj\}$.

Assuming that $|\alpha| \leq N - d - m$, we have

$$\begin{aligned} |I_1^j(x)| &\leq \text{const.} \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \int_{2Rj \leq |\xi| \leq 6Rj} (1 + |\xi|)^{d-j+|\alpha|-|\beta|} (e^{6RA})^j c_0^{|\beta|+j+1} j! \beta! d\xi \\ &\leq \text{const.} (c_0 e^{6RA})^j \alpha! \left| \sum_{\beta \leq \alpha} \frac{1}{(\alpha - \beta)!} \int_{2Rj \leq |\xi| \leq 6Rj} (1 + |\xi|)^{d-j+|\alpha|-|\beta|} j! \right| c_0^N \\ &\leq \text{const.} (c_0 e^{6RA})^j \alpha! \left| \sum_{\beta \leq \alpha} \frac{1}{(\alpha - \beta)!} \int_{0 \leq \rho \leq 6Rj} \rho^{d-j+|\alpha|-|\beta|+m-1} j! d\rho \right| c_0^N \end{aligned}$$

(We have used the fact that $d - j + |\alpha| \leq 0$.)

$$\leq \text{const.} \alpha! \left(\frac{c_0 e^{6RA}}{6R}\right)^j (6R)^{d+m+N} c_0^N \left(\sum_{\beta \leq \alpha} \frac{1}{(\alpha - \beta)!} \frac{j!}{j^{j-d-m-|\alpha|+|\beta|}} \right).$$

Therefore, for some $\tilde{c}_4 > 0$ independent of j and N ,

$$|I_1^j(x)| \leq \alpha! c_4^{N+1} \left(\frac{c_0 e^{6RA}}{6R} \right)^j$$

Similarly, after modifying the constant $\tilde{c}_4$ if necessary,

$$\begin{aligned} |I_2^j(x)| &\leq \text{const.} \cdot \alpha! \sum_{\beta \leq \alpha} \frac{1}{(\alpha - \beta)!} \left(\frac{1}{1 + 6Rj} \right)^{j-d-m-|\alpha|+|\beta|} c_0^{|\beta|+j+1} j! \\ &\leq \alpha! c_4^{N+1} \left(\frac{c_0}{6R} \right)^j \end{aligned}$$

We recall that $c_0 \leq R$. At this point we choose U' so small that if $A = \sup_{x, y \in U'} |Z(x) - Z(y)|$, then $c_0 e^{6RA} < 6R$.

We then get a constant $c_4 > 0$ such that: $|M^\alpha(\text{op } \tilde{k} - \tilde{K}_N)u(x)| \leq \alpha! c_4^{N+1}$ for $|\alpha| \leq N - d - m$.

(D) Estimate of $M^\alpha(R_N u)$.

$$\begin{aligned} R_N u(x) &= \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \int \int e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \xi} u(y) \\ &\quad \cdot \left(\sum_{j \leq N} (\phi_{N+1}(\xi) - \phi_j(\xi)) k_j(x, \xi) \right) dZ(y) d\xi \end{aligned}$$

is hypo-analytic since each $\phi_{N+1} - \phi_j$ is supported in $2Rj \leq |\xi| \leq 3R(N+1)$.

We estimate

$$\begin{aligned} \left| \sum_{j \leq N} (\phi_{N+1}(\xi) - \phi_j(\xi)) k_j(x, \xi) \right| &\leq \left(\sum_{j \leq N} c_0^{j+1} j! |\xi|^{-j} \right) |\xi|^d \\ &\leq \left(\sum_{j \leq N} c_0^{j+1} j! \left(\frac{1}{2Rj} \right)^j \right) |\xi|^d \quad (\text{since } 2Rj \leq |\xi|) \\ &\leq \left(\sum_{j \leq N} \left(\frac{c_0}{2Re} \right)^j j \right) c_0 e |\xi|^d \quad \text{since } \frac{j!}{j^j} \leq j e^{-j+1}. \end{aligned}$$

It follows that

$$|R_N u(x)| \leq \text{constant} \int_{|\xi| \leq 3R(N+1)} |\xi|^d d\xi \leq \text{const.} \cdot 3R(N+1)^{d+m}$$

which in turn implies that there is a constant $\tilde{c}_5 > 0$ such that $|R_N u(x)| \leq \tilde{c}_5^{N+1}$. Moreover, by hypo-analyticity, we get $c_5 > 0$ satisfying $|M^\alpha R_N u(x)| \leq \alpha! c_5^{N+1}$ for all $|\alpha| \leq N$.

(E) Estimate of $M^\alpha(T_N u)$.

$$T_N u(x) = \lim_{\varepsilon \rightarrow 0} \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \int \int e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \xi - \varepsilon |\xi|^2} t_N(x, y, \xi) u(y) dZ(y) d\xi$$

where

$$t_N(x, y, \xi) = \sum_{|\alpha| \leq N+1} \frac{1}{\alpha!} \{ (D_\xi^\alpha (\phi_{N+1}(\xi) k_\alpha(x, y, \xi)) - \phi_{N+1}(\xi) D_\xi^\alpha k_\alpha(x, y, \xi)) \}.$$

We can therefore take the limit under the integral sign and write

$$T_N u(x) = \sum_{|\alpha| \leq N+1} A_\alpha(x)$$

where for each α , $|\alpha| \leq N+1$,

$$A_\alpha(x) = \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \sum_{0 \neq \beta \leq N} \int_{2R(N+1) \leq |\xi| \leq 3R(N+1)} \int e^{\sqrt{-1}Z((x)-Z(y)) \cdot \xi} \frac{1}{\beta!} \\ \cdot (D_\xi^\beta \phi_{N+1}(\xi)) \frac{D_\xi^{\alpha-\beta} k_\alpha(x, y, \xi)}{(\alpha-\beta)!} u(y) dZ(y) d\xi$$

Therefore

$$|A_\alpha(x)| \leq \text{const.} \cdot \alpha! c_0^{|\alpha|+1} (e^{3RA})^{N+1} \left[\sum_{0 \neq \beta \leq \alpha} \frac{1}{\beta!} \left(\frac{[3R(N+1)]^{d+m+1}}{[2R(N+1)]^{|\alpha-\beta|}} \right) \left(\frac{c_0}{R} \right)^{|\beta|} \right] \\ \leq \text{const.} \cdot \frac{\alpha!}{[2R(N+1)]^{|\alpha|}} (e^{3RA})^{N+1} c_0^{|\alpha|+1} \\ \cdot [3R(N+1)]^{d+m+1} \left(\sum_{0 \leq \beta \leq \alpha} \frac{[2(N+1)c_0]^{|\beta|}}{\beta!} \right).$$

Since $|\alpha| \leq N$ and R may be taken to be larger than 1, we know that the factor $\frac{\alpha!}{[2R(N+1)]^{|\alpha|}} \leq 1$. Therefore, we conclude that there is a constant $c_6 \geq 0$ for which $|M^\alpha(T_N u)| \leq c_6^{N+1} N!$ whenever $|\alpha| \leq N$.

From (a)-(e) we conclude that there is a positive number c such that

$$|M^\alpha(\text{op } k - \text{op } \tilde{k})u(x)| \leq c^{N+1} N!$$

for all α , $|\alpha| \leq N - m - d$.

If we take $|\alpha| = N - m - d$, we can get a constant $\tilde{c} \geq c$ satisfying:

$$\forall \alpha, |M^\alpha(\text{op } k - \text{op } \tilde{k})u(x)| \leq \tilde{c}^{|\alpha|+1} \alpha! \quad \text{for every } x \in U'.$$

By using integration by parts we also reach the same conclusion for $u \in \mathcal{E}'(U')$. Indeed all we need is a representation of the form $u = \sum_{|\alpha| \leq N} M^\alpha u_\alpha$ where each $u_\alpha \in C_c^0(U')$ which is always possible. We have thus shown that $(\text{op } k - \text{op } \tilde{k})u$ is in $C^\infty(U')$ and that there is $c > 0$ such that for all $\alpha \in Z_m^+$,

$$|M^\alpha(\text{op } k - \text{op } \tilde{k})u(x)| \leq c^{|\alpha|+1} \alpha!.$$

By Theorem 3.1 of [1] it follows that $\text{op } k u - \text{op } \tilde{k} u$ is a hypo-analytic function.

II. APPLICATIONS

1. Parametrix for an elliptic operator. As an application of Theorem 4.1 we consider here the construction of a parametrix for an elliptic hypo-analytic differential operator. We will begin by composing a hypo-analytic differential operator A with a hypo-analytic pseudodifferential operator B . In [3] we introduced hypo-analytic differential operators. In the local chart (U, Z) , the operator A is given by $A = \sum_{|\alpha| \leq n} a_\alpha(x) N^\alpha$ where each $a_\alpha(x)$ is a hypo-analytic function and $N_j = -\sqrt{-1} M_j$ for $j = 1, \dots, m$.

Theorem 4.1 of the previous chapter allows us to represent the operator B by a symbol $b(x, \theta)$. From §2, Theorem 2.1 we know that both $B \circ A$ and $A \circ B$ are continuous linear maps from $\mathcal{E}'(U)$ to $\mathcal{D}'(U)$. We first assume that the operator $A = a(x) N^\beta$ for some hypo-analytic function $a(x)$ and some index β . Then $B(Au)(x)$ is by definition the limit as $\varepsilon \rightarrow +0$ of

$$B^\varepsilon(Au)(x) = \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \iint e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \xi - \varepsilon |\xi|^2} b(x, \xi) a(y) N^\beta u(y) dZ(y) d\xi.$$

On the other hand, $\lim_{\varepsilon \rightarrow 0^+} B^\varepsilon(Au)(x) = C \circ (N^\beta u)(x)$ where C is a hypo-analytic pseudodifferential operator with amplitude given by $b(x, \xi) a(y)$. Therefore, Theorem 4.1 tells us that C can be represented by the symbol $c(x, \xi) = \sum_\alpha \frac{\partial_\xi^\alpha b N^\alpha a(x)}{\alpha!}$. It follows that modulo a hypo-analytic function, we can write

$$B(Au)(x) = \lim_{\varepsilon \rightarrow 0^+} \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \iint e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \xi - \varepsilon |\xi|^2} \xi^\beta c(x, \xi) u(y) dZ(y) d\xi.$$

The latter says that a symbol of $B \circ A$ is given by

$$\xi^\beta c(x, \xi) = \sum_\alpha \frac{\partial_\xi^\alpha b(x, \xi) N^\alpha (a(x) \xi^\beta)}{\alpha!}.$$

On the other hand, applying the operator A to

$$B^\varepsilon u(x) = \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \iint e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \xi - \varepsilon |\xi|^2} b(x, \xi) u(y) dZ(y) d\xi$$

gives

$$\begin{aligned} A(B^\varepsilon u(x)) &= \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \iint e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \xi - \varepsilon |\xi|^2} \\ &\quad \cdot \left(\sum_{\gamma \leq \beta} \binom{\beta}{\gamma} \xi^{\beta-\gamma} a(x) N^\beta b(x, \xi) \right) u(y) dZ d\xi \\ &= \left(\frac{1}{4\pi^3} \right)^{\frac{m}{2}} \iint e^{\sqrt{-1}(Z(x)-Z(y)) \cdot \xi - \varepsilon |\xi|^2} \\ &\quad \cdot \left(\sum_\alpha \left[\frac{\partial_\xi^\alpha (a(x) \xi^\beta) N^\alpha b(x, \xi)}{\alpha!} \right] \right) u(y) dZ d\xi. \end{aligned}$$

This means that $A \circ B$ has a symbol given by

$$\sum_{\alpha} \frac{\partial_{\xi}^{\alpha} (a(x) \xi^{\beta}) N^{\alpha} b(x, \xi)}{\alpha!}.$$

By linearity, we will have the same formulas for the symbol of $B \circ A$ and $A \circ B$ when A is also given by $A = \sum_{|\alpha| \leq n} a_{\alpha}(x) N^{\alpha}$.

We have thus shown that if either A or B is hypo-analytic differential operator, the composition $A \circ B$ is hypo-analytic pseudodifferential operator with symbol

$$\sum_{\alpha} \frac{\partial_{\xi}^{\alpha} a(x, \xi) N^{\alpha} b(x, \xi)}{\alpha!}.$$

Definition 1.1. Let $P = \sum_{|\alpha| \leq k} a_{\alpha}(Z(x)) M^{\alpha}$ where the $a_{\alpha}(z)$ are holomorphic in a neighborhood of $Z(U)$ in C^m . We say a point $(x, \xi) \in T^*U \setminus \{0\}$ is in the characteristic set of P if the point $(Z(x), \check{Z}_x(x)\xi)$ is in the characteristic set of $P^Z = \sum_{|\alpha| \leq k} a_{\alpha}(z) (\frac{\partial}{\partial z})^{\alpha}$.

Notation. Char P = the characteristic set of P as given by Definition 1.1.

Definition 1.2. A hypo-analytic differential operator P is said to be elliptic at a point $x \in U$ if for every $(x, \xi) \in T^*U$, $(x, \xi) \notin \text{Char } P$.

Now suppose $P = \sum_{|\alpha| \leq k} a_{\alpha}(Z(x)) M^{\alpha}$ is a hypo-analytic differential operator that is elliptic at our central point $0 \in U$. Since $Z(0) = 0$ and $dZ(0) = \text{Id}$, we can find a neighborhood $\mathcal{O}$ of 0 in C^m , a cone $\mathcal{C}$ in C_m containing $R_m \setminus \{0\}$ and constants $c, R > 0$ such that: when $z \in \mathcal{O}$ and $\zeta \in \mathcal{C}$, $|\zeta| \geq R$ we have $|\sum_{|\alpha| \leq k} a_{\alpha}(z) \zeta^{\alpha}| \geq c|\zeta|^k$.

We now have all the ingredients we need to state

Theorem 1.1. Let A be hypo-analytic differential operator in Ω that is elliptic of order d . Given any relatively compact open subset $\tilde{\Omega}$ of Ω , there is a hypo-analytic pseudodifferential operator B in $\tilde{\Omega}$ of order $-d$ such that $AB - I$ and $BA - I$ are hypo-analytic regularizing in $\tilde{\Omega}$.

The proof of this theorem is a simple adaptation of that of the corresponding theorem for analytic pseudodifferential operators as given by Treves [8]. Therefore we omit it.

2. Propagation of hypo-analyticity. In [3] it was shown that hypo-analytic singularities for solutions propagate along the bicharacteristics of hypo-analytic differential operators. Here we extend this result to what may be called classical hypo-analytic pseudodifferential operator. This result may also be viewed as an extension of a theorem of Hanges [4].

We will work in the hypo-analytic local chart (U, Z) of Chapter I. Let P be a classical hypo-analytic pseudodifferential operator with principal symbol p . Let $t \rightarrow (x(t), \xi(t)) = \gamma(t)$ be a curve in $T^*U \setminus \{0\}$ and set $\hat{\gamma}(t) = (\hat{x}(t), \hat{\xi}(t)) = (Z(x(t)), \check{Z}_x(x(t))\xi(t))$.

Definition 2.1. The curve $\gamma(t)$ is said to be a bicharacteristic for P if the equations

$$\frac{d\tilde{x}}{dt} = \frac{\partial p}{\partial \tilde{\xi}}(\tilde{x}(t), \tilde{x}(x)), \quad \frac{d\tilde{\xi}}{dt} = \frac{-\partial p}{\partial z}(\tilde{x}(t), \tilde{\xi}(t))$$

hold.

We can now state the theorem of this section.

Theorem 2.1. Assume $p(0, \xi_0) = 0$ and P is of principal type at $(0, \xi_0)$. Suppose $\gamma = \{(x(t), \xi(t))\}$ is a bicharacteristic for P through $(x(0), \xi(0)) = (0, \xi_0)$ and that Pu is hypo-analytic on γ . Then either u is hypo-analytic at every point of γ or u is not hypo-analytic at any point of γ .

The proof will use a version of the FBI transform as developed by Sjöstrand in [7]. We will therefore first discuss Sjöstrand's FBI transformations adapted to our situation here.

Let H be a totally real submanifold of C^m of maximal dimension with defining functions $h_1, \dots, h_m$.

Define

$$\Lambda_H = \left\{ \left(x, \frac{2}{i} \partial h(x) \right) : h \in C^\infty(C^m, \mathbb{R}), h \equiv 0 \text{ on } H \right\}.$$

Note that if $x_0 \in H$, then $(x_0, \xi_0) \in \Lambda_H$ iff $\exists$ real numbers $t_1, \dots, t_m \ni$

$$\xi_0 = \frac{2}{i} \sum_{j=1}^m t_j \partial h_j(x_0).$$

Fix a point $(y_0, \eta_0) \in \Lambda_H$. Let φ be a holomorphic function defined near $(x_0, y_0) \ni$

$$(2.1) \quad \frac{\partial \varphi}{\partial y}(x_0, y_0) = -\eta_0,$$

$$(2.2) \quad \det \frac{\partial^2 \varphi}{\partial x \partial y}(x_0, y_0) \neq 0,$$

$$(2.3) \quad \Im \varphi_{yy}(x_0, y_0)|_{T_{y_0} H \times T_{y_0} H} > 0.$$

Here $\Im \varphi$ is considered as a function on $C^n \times H$, defined locally.

Set

$$\varphi_1(x, y) = -\Im \varphi(x, y).$$

Condition (2.1) implies that $H \ni y \mapsto \varphi_1(x_0, y)$ has a critical point at y_0 since $\frac{2}{i} \frac{\partial \varphi_1}{\partial y}(x_0, y_0) = \frac{\partial \varphi}{\partial y}(x_0, y_0) = -\eta_0$ and that therefore $d_y \varphi_1(x_0, y_0) = dh(y_0)$ for some h vanishing on H . This together with condition (2.3) and the implicit function theorem give us neighborhoods $N(x_0)$ of x_0 in C^m , $N(y_0)$ of y_0 in H and a unique C^∞ function $y = y(x) : N(x_0) \rightarrow N(y_0)$ such that $y(x)$ is the unique critical point for $H \ni y \mapsto \varphi_1(x, y)$, $x \in N(x_0)$. We next note that for $x \in N(x_0)$, $(y(x), \frac{-2}{i} \frac{\partial \varphi_1}{\partial y}(x, y(x))) \in \Lambda_H$. Indeed, this follows from the fact that $H \ni y \mapsto \varphi_1(x, y)$ has a critical point at $y(x)$ and that $h_1, \dots, h_m$ are the defining functions for H .

For $x \in N(x_0)$, let $\eta(x) = \frac{-2}{i} \frac{\partial \varphi_1}{\partial y}(x, y(x))$. Then

$$(y(x), \eta(x)) = \left(y(x), \frac{-2}{i} \frac{\partial \varphi_1}{\partial y}(x, y(x)) \right) \in \Lambda_H.$$

Moreover, for x in $N(x_0)$, $y(x)$ is the unique point in $N(y_0)$ such that

$$\frac{-\partial \varphi}{\partial y}(x, y(x)) = \frac{-2}{i} \frac{\partial \varphi_1}{\partial y}(x, y(x)) \in (\Lambda_H)_{y(x)}.$$

This is due to the uniqueness of the critical point.

Let $\Phi(x) = \varphi_1(x, y(x))$. Let $a(x, y, \lambda)$ be a classical analytic symbol defined near (x_0, y_0) and elliptic at this point. For Ψ a real-valued function defined on an open set W in C^m , we define the space $H_\Psi^{\text{loc}}(W) = \{v : W \times R_+ \rightarrow C : v(z, \lambda) \text{ is holomorphic in } z \text{ and for any } K \subset\subset W \text{ and } \varepsilon > 0 \exists c \ni |v(z, \lambda)| \leq ce^{\lambda(\Psi(z)+\varepsilon)} \text{ for all } z \in K, \lambda \geq 1\}$.

Let $u \in D'(N(y_0))$, and for z in $N(x_0)$ set

$$Tu(z, \lambda) = \int_H e^{i\lambda\varphi(z, y)} a(z, y, \lambda) \chi(y) u(y) dy$$

where $\chi \in C_0^\infty(N(y_0))$, $\chi \equiv 1$ near y_0 .

Here we are assuming that the neighborhoods $N(y_0)$ and $N(x_0)$ have been contracted so that the symbol a and the phase function φ are defined. It is easily checked that

$$T : D'(N(y_0)) \longrightarrow H_\Phi^{\text{loc}}(N(x_0)).$$

In the sequel, $WF_{ha}u$ denotes the hypo-analytic wave front set of Baouendi-Chang-Treves [1]. Our proof of Theorem 2.1 will use the following proposition of Sjöstrand [7].

Proposition 2.1. *Let $z_1 \in N(y_0)$. Then $(y(z_1), \eta(z_1)) \notin WF_{ha}u$ iff $Tu \in H_{\Phi-\varepsilon_0}^{\text{loc}}(W)$ for some $\varepsilon_0 > 0$ and some neighborhood W of z_1 .*

Proof of Theorem 2.1. In order to obtain a suitable phase function, we will need the following two lemmas from [6]. For notational convenience we will use y_0 for $0 \in Z(U) = H$.

Lemma 2.1. *Set $z_0 = (y_0' - i\xi_0', 0) \in C^{n-1} \times C$. There exists a holomorphic function φ defined near (z_0, y_0) which solves*

$$\frac{\partial \varphi}{\partial z_n}(z, y) = p \left(y, \frac{-\partial \varphi}{\partial y}(z, y) \right)$$

and satisfies (2.1)–(2.3) with $\eta_0 = \xi_0$.

We remark that the lemma is proved by using the Cauchy-Kovalevska theorem, which guarantees the existence of a holomorphic φ that solves the initial value problem

$$\frac{\partial \varphi}{\partial z_n} = p \left(y, \frac{-\partial \varphi}{\partial y} \right)$$

and

$$\varphi(z, 0, y) = \frac{i}{2} \sum_{j=1}^{n-1} (z_j - y_j)^2 - (\xi_0)_n y_n + iC(y_n - (y_0)_n)^2$$

where $\Re C$ is chosen sufficiently large. In the sequel, the neighborhoods $N(z_0)$, $N(y_0)$ and the function Φ are related to the φ of Lemma 2.1 as before.

Lemma 2.2. *There is an elliptic analytic symbol $a(z, y, \lambda)$ such that the FBI transformation T with phase φ and symbol a satisfies $D_{z_n} T = TP$ in H_{Φ, z_0} .*

That is, if $Y \subseteq Z(U) = H$ is a small neighborhood of y_0 , then for z in $W \subseteq C^m$ a small neighborhood of $z_0 = (y'_0 - i\xi'_0, 0)$ and $u \in \mathcal{E}'(Y)$ we have

$$D_{z_n} Tu - TPu \in H_{\Phi-\varepsilon}^{\text{loc}}(W)$$

for some $\varepsilon > 0$.

The symbol $a(z, y, \lambda)$ is constructed by solving the transport equations at each degree of homogeneity.

We recall now that

$$\hat{y}(t) = (\hat{x}(t), \hat{\xi}(t))$$

and

$$\hat{y}(0) = (y_0, \xi_0) = (Z(x(0)), \check{Z}_x(x(0))\xi_0).$$

Write $y_0 = (y'_0, (y_0)_n)$ and $\xi_0 = (\xi'_0, (\xi_0)_n)$.

We will use the equations

$$(2.4) \quad \begin{cases} \frac{\partial \varphi}{\partial z_n}(z, y) = p\left(y, \frac{-\partial \varphi}{\partial y}(z, y)\right), \\ \frac{\partial \varphi}{\partial y}(z_0, y_0) = -\xi^0 \end{cases}$$

to prove that $\hat{\xi}(t) = -\frac{\partial \varphi}{\partial y}(y'_0 - i\xi'_0, t, \hat{x}(t))$.

We recall that

$$\begin{cases} \frac{d\hat{x}}{dt} = \frac{\partial p}{\partial \xi}(\hat{x}(t), \hat{\xi}(t)) \quad \text{and} \\ \frac{d\hat{\xi}}{dt} = \frac{-\partial p}{\partial z}(\hat{x}(t), \hat{\xi}(t)). \end{cases}$$

Hence

$$\begin{aligned} & \frac{d}{dt} \left[\frac{\partial \varphi}{\partial y}(y'_0 - i\xi'_0, t, \hat{x}(t)) \right] \\ &= \varphi_{yz_n}(y'_0 - i\xi'_0, t, \hat{x}(t)) + \varphi_{yy}(y'_0 - i\xi'_0, t, \hat{x}(t)) \frac{d\hat{x}}{dt} \\ &= \varphi_{yz_n}(y'_0 - i\xi'_0, t, \hat{x}(t)) + \varphi_{yy}(y'_0 - i\xi'_0, t, \hat{x}(t)) \frac{\partial p}{\partial \xi}(\hat{x}(t), \hat{\xi}(t)) \end{aligned}$$

Now (2.4) implies that

$$\varphi_{yz_n}(z, y) = \frac{\partial p}{\partial y}\left(y, \frac{-\partial \varphi}{\partial y}\right) - \frac{\partial p}{\partial \xi}\left(y, \frac{-\partial \varphi}{\partial y}\right) \varphi_{yy}(z, y).$$

It follows that

$$\begin{aligned}
 & \frac{d}{dt} \left[\frac{-\partial \varphi}{\partial y} (y'_0 - i\xi'_0, t, \tilde{x}(t)) \right] \\
 (2.5) \quad &= \frac{-\partial p}{\partial y} \left(\tilde{x}(t), -\frac{\partial \varphi}{\partial y} (y'_0 - \xi'_0, t, \tilde{x}(t)) \right) \\
 &+ \frac{\partial p}{\partial \xi} \left(\tilde{x}(t), -\frac{\partial \varphi}{\partial y} (y'_0 - i\xi'_0, t, \tilde{x}(t)) \right) \varphi_{yy} (y'_0 - i\xi'_0, t, \tilde{x}(t)) \\
 &- \varphi_{yy} (y'_0 - i\xi'_0, t, \tilde{x}(t)) \frac{\partial p}{\partial \xi} (\tilde{x}(t), \tilde{\xi}(t)).
 \end{aligned}$$

But $\tilde{\xi}(t)$ also satisfies (2.5) since

$$\begin{aligned}
 \frac{d\tilde{\xi}}{dt} &= -\frac{\partial p}{\partial y} (\tilde{x}(t), \tilde{\xi}(t)) + \frac{\partial p}{\partial \xi} (\tilde{x}(t), \tilde{\xi}(t)) \varphi_{yy} (y'_0 - i\xi'_0, t, \tilde{x}(t)) \\
 &- \varphi_{yy} (y'_0 - i\xi'_0, t, \tilde{x}(t)) \frac{\partial p}{\partial \xi} (\tilde{x}(t), \tilde{\xi}(t)) \\
 &= -\frac{\partial p}{\partial y} (\tilde{x}(t), \tilde{\xi}(t)).
 \end{aligned}$$

Moreover, by 2.4, $\frac{-\partial p}{\partial y} (y'_0 - i\xi'_0, 0, y_0) = \xi_0 = \tilde{\xi}(0)$.

We conclude that

$$(2.6) \quad \tilde{\xi}(t) = \frac{-\partial \varphi}{\partial y} (y'_0 - i\xi'_0, t, \tilde{x}(t)).$$

For $t \in [0, 1]$, let

$$z(t) = z_0 + (0', t) = (y'_0 - i\xi'_0, t) \in C^{n-1} \times R.$$

We now recall that for z near z_0 , $y(z)$ is the unique point in $N(y_0) \subseteq H$ such that

$$y(z_0) = y_0 \quad \text{and} \quad \frac{-\partial \varphi}{\partial y} (z, y(z)) \in (\Lambda_H)_{y(z)}.$$

But by (2.6), $\tilde{\xi}(t) = \frac{-\partial \varphi}{\partial y} (z(t), \tilde{x}(t))$ and since the forms $\frac{\partial}{\partial t} h_1, \dots, \frac{\partial}{\partial t} h_n$ are real on $H = Z(U)$ and span all of T^*H , we know that

$$\tilde{\xi}(t) = \tilde{Z}_x(x(t))\xi(t) \in (\Lambda_H)_{\tilde{x}(t)}.$$

It therefore follows that

$$y(z(t)) = \tilde{x}(t).$$

In our previous notation,

$$\eta(z(t)) = \frac{-\partial \varphi}{\partial y} (z(t), y(z(t))) = \frac{-\partial \varphi}{\partial y} (z(t), \tilde{x}(t)) = \tilde{\xi}(t).$$

Thus

$$(2.7) \quad (\tilde{x}(t), \tilde{\xi}(t)) = (y(z(t)), \eta(z(t))).$$

Since $WF_{ha}(Pu) \cap \gamma = \emptyset$ and γ is compact, (2.7) and Proposition (2.1) tell us that

$$T(Pu) \in H_{\Phi-\varepsilon_0}^{\text{loc}}(N)$$

for some $\varepsilon_0 > 0$ and a neighborhood N of $\{z(t) = 0 \leq t \leq 1\}$ in C^m . If W is chosen as in Lemma 2.2, then

$$D_{z_n} Tu \in H_{\Phi-\varepsilon_0}^{\text{loc}}(N \cap W).$$

This may require a modification of ε_0 .

Now $z(0) = z_0 \in N \cap W$. Therefore, $\exists t_1 > 0$ such that $N \cap W$ is a neighborhood of $\{z(t) : 0 \leq t \leq t_1\}$. It is crucial to note that the size of t_1 is independent of the distribution u .

If now K is a compact neighborhood of $\{z(t) : 0 \leq t \leq t_1\}$, then $\exists c > 0$ such that

$$(2.8) \quad |D_{z_n} Tu(z, \lambda)| \leq ce^{\lambda(\Phi(z) - \frac{\varepsilon_0}{2})} \quad \forall z \in K \text{ and } \lambda \geq 1.$$

If $(y_0, \xi_0) = (y(z(0)), \eta(z(0))) \notin WF_{ha}u$, we know that, after modifying c and ε_0 ,

$$(2.9) \quad |Tu(z, \lambda)| \leq ce^{\lambda(\Phi(z) - \frac{\varepsilon_0}{2})} \quad \forall \lambda \geq 1 \text{ and } \forall z \text{ near } z_0.$$

From (2.7), (2.8) and (2.9), it follows that

$$WF_{ha}(u) \cap \{(y(t), \xi(t)) : 0 \leq t \leq t_1\} = \emptyset.$$

REFERENCES

1. M. S. Baouendi, C. H. Chang and F. Trèves, *Microlocal hypo-analyticity and extensions of CR functions*, J. Differential Geometry **18** (1983), 331–391.
2. S. Berhanu, *Hypo-analytic pseudodifferential operators*, Proc. Amer. Math. Soc. **105** (1989), 582–588.
3. ———, *Propagation of hypo-analyticity along bicharacteristics*, Pacific J. Math **138** (1989), 221–232.
4. N. Hanges, *Propagation of analyticity along real bicharacteristics*, Duke Math. J. **49** (1981), 269–277.
5. L. Hörmander, *An introduction to complex analysis in several variables*, Van Nostrand, Princeton, N.J., 1966.
6. J. Sjöstrand, *Singularités analytiques microlocales*, Astérisque **95** (1982).
7. ———, *The FBI transform for CR submanifolds of C^N* , (preprint).
8. F. Trèves, *Introduction to pseudodifferential and Fourier integral operators*, Volume I, Plenum Press, 1980.

DEPARTMENT OF MATHEMATICS, TEMPLE UNIVERSITY, PHILADELPHIA, PENNSYLVANIA 19122