ON CONJUGACY SEPARABILITY OF FUNDAMENTAL GROUPS OF GRAPHS OF GROUPS

M. SHIRVANI

ABSTRACT. A complete determination of when the elements of a fundamental group of a (countable) graph of profinite groups are conjugacy distinguished is given. By embedding an arbitrary fundamental group G into one with profinite vertex groups and making use of the above result, questions on conjugacy separability of G can be reduced to the solution of equations in the vertex groups of G .

1. INTRODUCTION

An element g of a group G is said to be *conjugacy distinguished* (or conjugacy separable) in G if for every element h of G not conjugate to g there exists $N \triangleleft_f G$ such that gN and hN are nonconjugate in G/N . A group is *conjugacy separable* if all its elements are conjugacy distinguished. The best-known classes of conjugacy separable groups are polycyclic-by-finite groups [10], profinite groups [11], free-by-finite groups [5], and certain Fuchsian groups [19]. Results are also known on conjugacy separability of certain amalgamated free products (e.g., of free groups [18]), and some one-relator groups with torsion [1]. In the last mentioned paper the authors ask whether every one-relator group with torsion is conjugacy separable. More generally, one can ask when the fundamental group of a graph of groups is conjugacy separable.

In order to investigate these problems one might adopt the following strategy: let G be a residually finite group, with $\widehat{G}$ its profinite completion (i.e., $\widehat{G}$ is the inverse limit of the system of finite images G/N of G). It is easily shown that if $g, h \in G$ are conjugate in every finite image of G , then they are conjugate in $\widehat{G}$, and so G is conjugacy separable if and only if whenever $g, h \in G$ are conjugate in $\widehat{G}$, then they are conjugate in G . Unfortunately, if G is a non-trivial fundamental group of a graph of groups, then $\widehat{G}$ is still not sufficiently well understood (cf. [7, 20]) to allow for the successful completion of the last step. A complete answer can be given when the vertex groups are themselves profinite groups. This, in turn, allows necessary and sufficient conditions to be derived for the general case.

Let X be a connected graph, let $\mathcal{G} = \{A_x, x \in VX, H_e, e \in EX\}$ be a graph of groups over X , and let $G = \pi_1(\mathcal{G}, X)$ be the fundamental group of

Received by the editors September 21, 1990.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 20E26; Secondary 20E06, 20E18.

this graph of groups (see [2, 4], or [14] for the relevant definitions and properties). We assume that G is residually finite, and denote by $\widehat{A}_x$ the profinite completion of the vertex group A_x with respect to the induced topology $\{A_x \cap M : M \triangleleft_f G\}$. Let $\overline{H}_e$ denote the topological closure of H_e in $\widehat{A}_{i(e)}$. The edge isomorphisms can be extended to these closures (cf. Lemma 3.1). The result is a graph of groups over X with fundamental group G^+ (for details see §3). The obvious maps $A_x \rightarrow \widehat{A}_x$ extend to an embedding of G into G^+ . If we denote the conjugacy class of g in G by g^G and $\bigcap_{M \triangleleft_f G} g^G M$ by $\text{cl}_G(g^G)$, then we have:

Theorem 3.4. *For every $g \in G$ we have $\text{cl}_G(g^G) = \text{cl}_{G^+}(g^{G^+}) \cap G$.*

The point of this result is that most elements of G^+ turn out to be conjugacy distinguished (i.e. $\text{cl}_{G^+}(g^{G^+}) = g^{G^+}$). If $g \in G$ is such an element, then g is conjugacy distinguished in G if and only if $g^{G^+} \cap G = g^G$. This condition is quite tractable, and amounts to whether certain equations hold in the vertex groups A_x and $\widehat{A}_x$. Assuming that we have enough information about the vertex groups and their completions, the conjugacy separability of G can be decided.

To state the results for G^+ , let Y be a maximal subtree of X , and write $A^+ = \langle \widehat{A}_x : x \in VX \rangle \leq G^+$. Then A^+ is the fundamental group of the graph of groups $\mathcal{G}$ restricted to Y , and G^+ is an HNN-extension with base group A^+ . As a consequence of 4.4, 4.6, and 4.9 below we get

Theorem A. *Let $a \in A^+$.*

(i) *If either no conjugate of a belongs to a vertex group, or X is locally finite, $a \in \widehat{A}_x$, and no $\widehat{A}_x$ -conjugate of a belongs to an edge subgroup, then a is conjugacy distinguished in A^+ .*

(ii) *Assume that X is locally finite, let $D = \bigcup_{e \in EX} \overline{H}_e$, and suppose $a \in D$ belongs to only a finite number of edge subgroups. Then a is conjugacy distinguished in A^+ if and only if there exist vertices $x_1, \dots, x_m$ of Y such that $D \cap a^{A^+} = D \cap \{a^{a_1 \cdots a_m} : a_i \in \widehat{A}_{x_i}\}$. $\square$*

For elements of G^+ we obtain from 5.1, 5.3, 5.6, and 5.8

Theorem B. (i) *All elements of $G^+ \setminus A^+$, and all elements of A^+ satisfying condition (i) of Theorem A, are conjugacy distinguished in G^+ .*

(ii) *Assume that X is locally finite, and let $D' = \bigcup_{e \in EX \setminus EY} \overline{H}_e$ and $C = \bigcup_{x \in VX} \widehat{A}_x$.*

Then $h \in D'$ is conjugacy distinguished in G^+ if and only if $h^{G^+} \cap C$ is a union of a finite number of sets of the form $C \cap \{h^g : g \in \widehat{A}_{x_1} t_{e_1}^{\varepsilon_1} \cdots \widehat{A}_{x_n} t_{e_n}^{\varepsilon_n} \widehat{A}_{x_{n+1}}, \text{ for some } e_i \in EX \setminus EY, \varepsilon_i = \pm 1, \text{ and } x_i \in VX\}$.

This in particular settles the problem for the case when the vertex groups are profinite, since such a G can be residually finite if and only if $G \cong G^+$, by 3.3 below.

2. NOTATION

Let X be a directed connected graph, $\mathcal{G}$ a graph of groups $\{A_x : x \in VX; H_e : e \in EX \text{ with isomorphisms } \theta_e \text{ from } H_e \leq A_{i(e)} \text{ to } H_{\bar{e}} \leq A_{l(e)}\}$ over

X , and $G = \pi_1(\mathcal{G}, X)$ the fundamental group of $(\mathcal{G}, X)$ relative to the choice of a fixed maximal subtree Y (see [2, 4], or [14] for details). It is well known that if $A = \pi_1(\mathcal{G}|_Y, Y)$ then the obvious map of A to $\langle A_x : x \in VY \rangle \subseteq G$ is an isomorphism, and G is an HNN-extension with base group A and stable letters t_e corresponding to the edges $e \in EX \setminus EY$:

$$G = \langle A, t_e : t_e^{-1} h t_e = h \theta_e \text{ for all } h \in H_e, e \in EX \setminus EY \rangle.$$

In the notation of [16] let $I = I(\mathcal{G}, X)$ denote the set of all sequences $(P_x)_{x \in VX}$ satisfying the following conditions: (a) $P_x \triangleleft A_x$ and there exists an integer $m = m(P)$ such that $|A_x : P_x| \leq m$ for all $x \in VX$, (b) $(P_{i(e)} \cap H_e) \theta_e = P_{i(e)} \cap H_{\bar{e}}$ for all $e \in EX$. For $P, Q \in I$ we write $P \leq Q$ if $P_x \subseteq Q_x$ for all x . For $P \in I$ let $\mathcal{G}_P$ denote the graph of groups $\{A_x/P_x, H_e P_{i(e)}/P_{i(e)}, \text{ induced isomorphisms } \theta_{e,P}\}$, and let $G_P = \pi_1(\mathcal{G}_P, X)$. The projections $A_x \rightarrow A_x/P_x$ clearly extend to an epimorphism $\pi_P : G \rightarrow G_P$. It is well known that G_P is free-by-finite (e.g., [14, Exercise 2 on p. 123]). Also note that if $M \triangleleft_f G$ then $P = (M \cap A_x)_{x \in VX} \in I$, and G/M is a homomorphic image of G_P . In particular, this implies that if G is residually finite then $\bigcap_{P \in I} P_x = \langle 1 \rangle$ for all $x \in VX$. We assume this from now on. For $g \in G$ write g^G for the conjugacy class of g in G .

Proposition 2.1. *Let $g \in G$. Then $\bigcap_{M \triangleleft_f G} g^G M = \bigcap_{P \in I} g^G \ker \pi_P$. In particular, $\bigcap_{M \triangleleft_f G} M = \bigcap_{P \in I} \ker \pi_P$.*

Proof. If $M \triangleleft_f G$ induces $P \in I$ then $M \supseteq \ker \pi_P$, so $g^G M \supseteq \bigcap_P g^G \ker \pi_P$. Conversely, given $P \in I$ let $P \uparrow = \{M \triangleleft_f G : M \cap A_x = P_x \text{ for all } x \in VX\}$. Now by a theorem of Dyer [5], G_P is conjugacy separable, which implies that

$$(g \pi_P)^{G_P} = \bigcap_{M \in P \uparrow} (g \pi_P)^{G_P} M \pi_P = \bigcap_{M \in P \uparrow} (g^G M) \pi_P,$$

whence $g^G \ker \pi_P = \bigcap_{M \in P \uparrow} g^G M$. The result follows. (For the final part take $g = 1$.) $\square$

3. PROFINITE CLOSURES

It is evident that if $P, Q \in I$, then $P \cap Q = (P_x \cap Q_x)_{x \in VX}$ also belongs to I . We refer to the topology on A_x with $\{P_x : P \in I\}$ as a fundamental system of open neighbourhoods of the identity as the I -topology. For each x let $\hat{A}_x = \varprojlim_{P \in I} (A_x/P_x)$, the inverse limit being formed relative to the partial order $\leq$ of I introduced above. Let $\bar{H}_e$ denote the topological closure of H_e in $\hat{A}_{i(e)}$. We have

Lemma 3.1. *For every $e \in EX$, the isomorphism $\theta_e : H_e \rightarrow H_{\bar{e}}$ extends to an isomorphism $\bar{\theta}_e : \bar{H}_e \rightarrow \bar{H}_{\bar{e}}$ such that $(\bar{H}_e \cap \bar{P}_{i(e)}) \bar{\theta}_e = \bar{H}_{\bar{e}} \cap \bar{P}_{i(e)}$.*

Proof. For each $P \in I$ there are canonical isomorphisms

$$\begin{aligned} \bar{H}_e / (\bar{H}_e \cap \bar{P}_{i(e)}) &\cong \bar{H}_e \bar{P}_{i(e)} / \bar{P}_{i(e)} = H_e \bar{P}_{i(e)} / \bar{P}_{i(e)} \\ &\cong H_e / (H_e \cap \bar{P}_{i(e)}) = H_e / H_e \cap P_{i(e)}, \end{aligned}$$

and $\bar{H}_e$ is naturally isomorphic to the inverse limit $\varprojlim_{P \in I} (H_e / H_e \cap P_{i(e)})$ [12]. The isomorphisms induced by θ_e on the quotients $H_e / H_e \cap P_{i(e)}$ are compatible

with the inverse limit structure, so the existence of $\bar{\theta}_e$ follows. (If one thinks of the inverse limit $\hat{A}_x$ as a subgroup of the cartesian product $\prod_{x \in VX} (A_x/P_x)$, then the image of the element $h = (h_P \pi_P) \in \bar{H}_e$ is $h\bar{\theta}_e = (h_P \theta_e \pi_P) \in \bar{H}_{\bar{e}}$.) The intersection property of $\bar{\theta}_e$ is also trivial from the construction. $\square$

Thus we have a graph of groups $\mathcal{G}^+ = \{\hat{A}_x, \bar{H}_e, \bar{\theta}_e\}$ over X . Let $G^+ = \pi_1(\mathcal{G}^+, X)$. The natural embeddings $A_x \rightarrow \hat{A}_x$ extend to a homomorphism $\mu: G \rightarrow G^+$. The next few results exploit the relationship between G^+ and the residual properties of G .

Lemma 3.2. *Let G , G^+ , and μ be as above. Then G^+ is residually finite and $\ker \mu = \bigcap_{M \triangleleft_f G} M$.*

Proof. The edge subgroups of G^+ are closed in their vertex groups by definition, and 3.1 implies that if $P \in I(\mathcal{G})$ then $\bar{P} = (\bar{P}_x) \in I(\mathcal{G}^+)$. The residual finiteness of G^+ is therefore a consequence of the theorem of [16]. For the second part let $P \in I$, and consider the following diagram with exact top row:

$$\begin{array}{ccc} \ker \mu \hookrightarrow G & \xrightarrow{\mu} & G^+ \\ \pi_P \downarrow & & \downarrow \pi_P^+ \\ G_P & \xrightarrow{\mu_P} & G_P^+ \end{array}$$

The diagram is easily seen to commute, and $A_x/P_x \cong \hat{A}_x/\bar{P}_x$ implies that μ_P is an isomorphism. Then $1 = (\ker \mu)\mu\pi_P^+ = (\ker \mu)\pi_P\mu_P$ implies that $\ker \mu \subseteq \ker \pi_P$ for all P , so $\ker \mu \subseteq \bigcap_P \ker \pi_P = \bigcap_{M \triangleleft_f G} M$ by 2.1. On the other hand, $G/\ker \mu$ is a subgroup of G^+ ; being residually finite, the reverse inclusion follows. $\square$

Before proceeding further we mention that, in general, profinite closedness of the edge subgroups in their vertex groups is not necessary for the residual finiteness of G (cf. [15, 17]). In the case of profinite vertex groups, however, we have the following result.

Corollary 3.3. *Let $G = \pi_1(\mathcal{G}, X)$, and assume that $A_x = \varprojlim_{P \in I} (A_x/P_x)$ for all $x \in VX$. Then G is residually finite if and only if every edge subgroup is profinitely closed in its vertex group.*

Proof. Suppose there exists an element $a \in \bar{H}_e \setminus H_e$ for some $e \in EX$. If e is an edge in the maximal subtree Y let $g = a^{-1}(a\bar{\theta}_e)$; otherwise let $g = a^{-1}t_e(a\bar{\theta}_e)t_e$. Then $g \neq 1$ in G , while $g\mu = 1$, so G is not residually finite by 3.2. The converse follows from 3.2 and the fact that when the edge subgroups are closed we have $G = G^+$. $\square$

If S is a subset of a group H , write $\text{cl}_H(S)$ for $\bigcap_{M \triangleleft_f H} SM$.

Theorem 3.4. *Let G , G^+ , and μ be as above. If $g \in G$ then $\text{cl}_G(g^G)\mu = \text{cl}_{G^+}((g\mu)^{G^+}) \cap G\mu$.*

We are not assuming that μ is injective.

Proof. Let $S = \text{cl}_G(g^G) = \bigcap_{P \in I} g^G \ker \pi_P$ (by 2.1). Then using $\ker \mu \subseteq \ker \pi_P$ (cf. the proof of 3.2) we have

$$S\mu = \bigcap_{P \in I} (g^G \ker \pi_P)\mu = \bigcap_{P \in I} (g\mu)^{G\mu}(\ker \pi_P)\mu.$$

Now it is easy to see from the commutative diagram in the proof of 3.2 and the fact that μ_P is an isomorphism, that $(\ker \pi_P)\mu = \ker \pi_P^+ \cap G\mu$. The proof of 2.1 applies without change to show that $\text{cl}_{G^+}(y^{G^+}) = \bigcap_{P \in I} y^{G^+} \ker \pi_P^+$ for all $y \in G^+$. Thus

$$S\mu = \bigcap_{P \in I} (g\mu)^{G\mu}(\ker \pi_P^+ \cap G\mu) = \bigcap_{P \in I} (g\mu)^{G\mu} \ker \pi_P^+ \cap G\mu,$$

which is certainly contained in $\bigcap_{P \in I} (g\mu)^{G^+} \ker \pi_P^+ \cap G\mu$. Conversely, suppose we have an element $z\mu \in \bigcap_{P \in I} (g\mu)^{G^+} \ker \pi_P^+$, where $z \in G$. Then for each P there exists $w_P \in G^+$ such that $z\mu\pi_P^+ = (w_P^{-1}(g\mu)\pi_P)\pi_P^+$. Since μ_P is an isomorphism, there exists $u_P \in G$ such that $u_P\pi_P\mu_P = w_P\pi_P^+$. The above equation now becomes $z\pi_P\mu_P = (u_P^{-1}gu_P)\pi_P\mu_P$, and since μ_P is injective we obtain $z \in \bigcap_{P \in I} g^G \ker \pi_P = S$. We have now shown that

$$S\mu = \bigcap_{P \in I} (g\mu)^{G\mu} \ker \pi_P^+ \cap G\mu \subseteq \bigcap_{P \in I} (g\mu)^{G^+} \ker \pi_P^+ \cap G\mu \subseteq S\mu,$$

as required. $\square$

Since conjugacy separable groups are residually finite, we henceforth assume that $\ker \mu = \langle 1 \rangle$, and identify G with $G\mu$. The statement of 3.4 can then be written more simply as $\text{cl}_G(g^G) = \text{cl}_{G^+}(g^{G^+}) \cap G$. What is the point of this result? Let g and g' be elements of G which are conjugate in every finite image of G . By 2.1 this means that for every $P \in I$ there exists $w_P \in G$ such that $g'\pi_P = (w_P^{-1}gw_P)\pi_P$, and at first glance it is conceivable that the “length” of the elements w_P (in whatever sense) might be unbounded. Now if we have $\text{cl}_{G^+}(g^{G^+}) = g^{G^+}$, then 3.4 implies that $g' \in \text{cl}_G(g^G) = g^{G^+} \cap G$, so $g' = w^{-1}gw$, where $w \in G^+$. This, of course, means that the above w_P can be chosen to have bounded length. It turns out that conjugacy separability is commonplace for elements of G^+ . Before we proceed with this, we state the following, which does not require $\ker \mu = \langle 1 \rangle$.

Corollary 3.5. *If g is conjugacy distinguished in G , then*

$$(g\mu)^{G^+} \cap G\mu = g^G\mu.$$

Proof. For if $(g\mu)^{G^+} \cap G\mu = S\mu$, where $S \neq g^G$, then

$$g^G\mu = \text{cl}_G(g^G)\mu = \text{cl}_{G^+}((g\mu)^{G^+}) \cap G\mu \supseteq (g\mu)^{G^+} \cap G\mu = S\mu,$$

and so $\text{cl}_G(g^G) \supseteq S \supseteq g^G$. $\square$

It is necessary to study conjugacy separability of the base group of G^+ first. The next section is devoted to the study of $A^+ = \pi_1(\mathcal{G}|_Y, Y)$, where Y is a maximal subtree of X .

4. CONJUGACY SEPARABILITY OF A^+

We begin with the following general fact.

Proposition 4.1. *Let $\mathcal{G}$ be a graph of groups over X , and assume that*

$$(1) \quad \bigcap_{P \in I} H_e P_{i(e)} = H_e \quad \text{for all } e \in EX,$$

with $G = \pi_1(\mathcal{G}, X)$. Let Y be a connected subgraph of X , and put $G(Y) = \langle A_x : x \in VY \rangle \leq G$. Then $G(Y) = \bigcap_{P \in I} G(Y) \ker \pi_P$. If Y is finite then $G(Y) = \bigcap_{M \trianglelefteq G} G(Y)M$.

Proof. Fix a vertex $x_0 \in VY$, and for each edge $e \in EX$ let $T_e \ni 1$ be a left transversal of H_e in $A_{i(e)}$. Then every element of G is uniquely represented by a normal word $t_1 \cdots t_n a$, where $a \in A_{x_0}$, $t_i \in T_{e_i}$ for $1 \leq i \leq n$, $(e_1, \dots, e_n)$ is a closed path at x_0 , and if $e_i = \bar{e}_{i-1}$ then $t_i \neq 1$ [8, Corollary 1]. Moreover, the proof shows that the elements of $G(Y)$ are represented by paths that entirely belong to Y . Now condition (1) means that, given any element $g = t_1 \cdots t_n a$ in normal form in G , we can find $P \in I$ such that $(t_1 \pi_P) \cdots (t_n \pi_P)(a \pi_P)$ is the normal form of $g \pi_P$ in G_P (relative to suitable transversals of the $H_e \pi_P$ in the $A_{i(e)} \pi_P$). In particular, if $g \in G \setminus G(Y)$ then g is represented by a path that goes outside Y , and since $g \pi_P$ is represented by the same path, we have $g \pi_P \notin G_P(Y) = \langle A_x \pi_P : x \in VY \rangle = G(Y) \pi_P$. The first part follows.

Now suppose that Y is finite. Then for any $P \in I$ we have the finitely generated subgroup $G(Y) \pi_P$ of the free-by-finite group G_P . Such subgroups are profinitely closed (e.g., [2, p. 229]). This evidently means that $G(Y) \ker \pi_P = \bigcap_{M \in P\uparrow} G(Y)M$, and the second part follows (cf. the proof of 2.1). $\square$

To simplify the notation let Y be a tree, $\mathcal{G}$ a graph of groups over Y , and $A = \pi_1(\mathcal{G}, Y)$. Also write $C = \bigcup_{x \in VY} A_x$ and $D = \bigcup_{e \in EY} H_e$, the subsets of *vertex elements* and *edge elements* respectively in A .

Proposition 4.2. *Let g be an element of A . Then the following assertions are true:*

(i) *If $g \in D$, then there exists a subtree Y_g of Y such that $g \in H_e$ if and only if e is an edge of Y_g .*

(ii) *If $g \notin D$, then there exists a FINITE subtree Y_g of Y such that $g \in A(Y_g)$, and if Z is any subtree of Y with $g \in A(Z)$, then $Y_g \subseteq Z$.*

Proof. (i) Let Y_g consist of all vertices x of Y such that $g \in A_x$, and all geodesics between them. If $g \in A_x \cap A_y$, and p is the Y -geodesic from x to y , then g belongs to every vertex group of p (since adjacent vertex groups in A generate their amalgamated free product). In other words, g belongs to every vertex group of Y_g . Moreover, $H_e = A_{i(e)} \cap A_{t(e)}$ for any edge e , so $g \in H_e$ if and only if $i(e)$ and $t(e)$ belong to Y_g .

(ii) Among all subtrees Z with $g \in A(Z)$ pick one with the fewest number of vertices, and call it Y_g . Fix a vertex x_0 in Y_g . Then the normal form $t_1 \cdots t_n a$ of g can be represented by a closed path p at x_0 , with p entirely contained in Y_g (since $g \in A(Y_g)$). Now let Z be a subtree such that $g \in A(Z)$. Suppose first that $Y_g = \{x_0\}$. If $x_0 \notin VZ$ then $g \in A_{x_0} \cap A(Z)$ must be an edge element (consider the geodesic from x_0 to $A(Z)$), a contradiction. So in this case $Y_g \subseteq Z$.

There remains the case where Y_g has more than one vertex. If Z contains every end vertex of Y_g then $Y_g \subseteq Z$, so suppose Z does not contain some end vertex x of Y_g , and let e be the edge of Y_g that ends at x . Now the path p must visit x at least once (since otherwise $g \in A(Y_g \setminus \{e, x\})$), and if $t_i \in A_x$ then $t_i \neq 1$ (since p looks like $\dots, e, x, \bar{e}, \dots$ at x , and we have a normal form). If Z and Y_g were disjoint then $g \in A(Z) \cap A(Y_g)$ would be an edge element, which is not the case. Thus $Z' = Z \cup Y_g \setminus \{e, x\}$ is a subtree of Y such that $g \in A(Z')$. If x_0 is on Z' then we have the immediate contradiction that g can be represented in normal form by a path q in Z' , and $q \neq p$ since x is on p but not on q . So we must have $x_0 = x$. But since Z and Y_g are not disjoint we can always choose x_0 to be a common vertex. This final contradiction shows that $Z \supseteq Y_g$, as required. $\square$

To define the notion of a reduced form for elements of A we need a definition. Let Δ' be a nonempty subtree of a finite tree Δ . By a *reduction process* R from Δ to Δ' we mean a sequence $\Delta_1, \dots, \Delta_m$ of subtrees of Δ such that $\Delta_1 = \Delta$, $\Delta_m = \Delta'$, and Δ_{i+1} is obtained from Δ_i by deleting an end vertex x_i (for $i = 1, \dots, m-1$). We let y_i be the vertex of Δ_i adjacent to x_i , so when $\Delta \subseteq Y$ we have

$$A(\Delta_i) = \langle A(\Delta_{i+1}) * A_{x_i} : H_{y_i x_i} = H_{x_i y_i} \text{ via } \theta_{y_i x_i} \rangle, \quad i = 1, \dots, m-1.$$

Let $\Delta \supseteq \Delta'$ be finite subtrees of Y , and $R = \{\Delta_1, \dots, \Delta_m\}$ a reduction process from Δ to Δ' . Say g is reduced (resp. cyclically reduced) relative to R if either $g \in A(\Delta')$ or $g \in A(\Delta_i) \setminus A(\Delta_{i+1})$ for some $i \leq m-1$, and then g is reduced (resp. cyclically reduced) in the amalgamated free product $A(\Delta_i)$ of $A(\Delta_{i+1})$ and A_{x_i} . Every element of $A(\Delta)$ can be written in reduced form, and is conjugate, in $A(\Delta)$, to a cyclically reduced element of $A(\Delta)$. Note that an element may be cyclically reduced relative to one reduction process, but not another.

Lemma 4.3. *Assume that (1) holds. Let g be a nonvertex element of A such that g is cyclically reduced relative to some reduction process from Y_g to a single vertex. Then $\text{cl}_A(g^A) = \bigcup_{\alpha} g_{\alpha}^A$, where each $g_{\alpha} \in A(Y_g)$.*

Proof. There exists $P_0 \in I$ such that $Y_{g\pi_P} = Y_g$ for all $P \subseteq P_0$. To begin with, we can choose P_1 such that $g\pi_P$ is cyclically reduced in $A(Y_g)\pi_P$ relative to R , for all $P \subseteq P_1$. This is because (1) can be used to ensure that the finitely many vertex elements in the cyclically reduced form of g are excluded from the finitely many edge subgroups encountered by R . In particular, if $P \subseteq P_1$ then $g\pi_P$ is not an edge element of $A\pi_P$. For such P we have $Y_{g\pi_P} \subseteq Y_g$ (since $g\pi_P \in A(Y_g)\pi_P$), and if $Q \subseteq P$ then $Y_{g\pi_Q} \supseteq Y_{g\pi_P}$. Since Y_g is finite, there exists a subtree $\Delta \subseteq Y_g$ and $P_0 \subseteq P_1$ such that for all $P \subseteq P_0$ we have $Y_{g\pi_P} = \Delta$. Then $g \in \bigcap_{P \subseteq P_0} A(\Delta) \ker \pi_P = A(\Delta)$ by 4.1, and then $\Delta = Y_g$ by definition of Y_g .

Now suppose $P \subseteq P_0$. Then $g\pi_P$ is a cyclically reduced nonvertex element of $A(Y_g)\pi_P$, so the conjugacy theorem for amalgamated free products [9] implies that the conjugacy class of $g\pi_P$ in $A(Y_g)\pi_P$ contains no vertex elements. If $\Delta \supseteq Y_g$ is a finite subtree then the conjugacy class of $g\pi_P$ in $A(\Delta)\pi_P$ contains no vertex elements (consider a reduction process from Δ to Y_g). Now

let $g' \in \text{cl}_A(g^A)$, and let $\Delta \supseteq Y_g$ be a finite subtree with $g' \in A(\Delta)$. Let $R = \{\Delta_1, \dots, \Delta_m\}$ be a reduction process from Δ to Y_g . Replacing g' by a conjugate in $A(\Delta)$ we may assume that g' is cyclically reduced relative to R , and we may choose P_0 such that $g'\pi_P$ is cyclically reduced for all $P \subseteq P_0$. If $P \subseteq P_0$ then $g\pi_P$ has minimal length in its $A(\Delta)\pi_P$ -conjugacy class, and belongs to $A(\Delta_2)\pi_P \setminus H_{y_1x_1}\pi_P$. The conjugacy theorem for amalgamated free products [9] implies that $g'\pi_P \in A(\Delta_2)\pi_P$. Continuing in this way we finally obtain $g'\pi_P \in A(Y_g)\pi_P$, whence $g' \in \bigcap_{P \subseteq P_0} A(Y_g) \ker \pi_P = A(Y_g)$, as required. $\square$

We can now prove

Theorem 4.4. *Let g be a nonvertex element of A such that g is cyclically reduced relative to some reduction process R from Y_g to a vertex. Assume that (1) holds and that $H_{x_1y_1}$ is compact in the I -topology (where x_1y_1 is the first edge deleted in the reduction process). Then g is conjugacy distinguished in A .*

Proof. By 4.3 we have $\text{cl}_A(g^A) = \bigcup g_\alpha^A$, where each $g_\alpha \in A(Y_g)$, and we may assume that each g_α is cyclically reduced relative to R . Fix α , and choose P_0 such that for all $P \subseteq P_0$, $g\pi_P$, and $g_\alpha\pi_P$ are cyclically reduced in $A(Y_g)\pi_P$ relative to R . The conjugacy theorem for amalgamated free products can now be applied to deduce that $g_\alpha\pi_P$ is conjugate to some cyclic permutation of $g\pi_P$, via an element of $H_{x_1y_1}\pi_P$. Replacing g by a cyclic permutation if necessary we have $g_\alpha\pi_P = (h_P^{-1}gh_P)\pi_P$ for all $P \subseteq P_0$, where $h_P \in H_{x_1y_1}$. Consider the function $f: H_{x_1y_1} \rightarrow A(Y_g)$ given by $f(h) = g_\alpha^{-1}h^{-1}gh$. With the appropriate profinitely topologies it is easy to see that f is continuous and $\ker \pi_P \cap A(Y_g)$ is closed in $A(Y_g)$. Thus for each $P \subseteq P_0$, $f^{-1}(\ker \pi_P)$ is a nonempty closed subset of $H_{x_1y_1}$, and $P \subseteq Q$ implies that $f^{-1}(\ker \pi_P) \subseteq f^{-1}(\ker \pi_Q)$, so we have the finite intersection property. If

$$h \in \bigcap_{P \subseteq P_0} f^{-1}(\ker \pi_P) = f^{-1} \left(\bigcap_{P \subseteq P_0} \ker \pi_P \right) = f^{-1}(\langle 1 \rangle),$$

then $g_\alpha^{-1}h^{-1}gh = 1$, so $g_\alpha \in g^A$. The result follows. $\square$

The next lemma collects the information we need on conjugacy of edge and vertex elements.

Lemma 4.5. *Let $C = \bigcup_{x \in VY} A_x$ and $D = \bigcup_{e \in EY} H_e$.*

(i) *Let $h \in D$, and let $a \in C$ be such that $h^a \in C$. Then $a \in A_x$ for some $x \in VY_h$.*

(ii) *Let $a \in A_x$, $b \in A_y$, and $a^b \in C$. Then there exists a vertex z such that $a, b \in A_z$.*

(iii) *Let $h \in D$, and w be a nonvertex element of A such that $h^w \in C$. Let $w = a_1 \cdots a_r$ be the reduced form of w (relative to some reduction process R of Y_w), where the $a_i \in C$. Then $h^{a_1 \cdots a_i} \in D$ for $1 \leq i \leq r-1$.*

(iv) *Let $a \in A_x$ be such that $a^{A_x} \cap D = \emptyset$. Then $a^A \cap D = \emptyset$, and $a^A \cap C = a^{A_x}$.*

Proof. (i) Choose x as close to Y_h as possible subject to $a \in A_x$, and get a contradiction to $h^a \in C$ by assuming that $x \notin VY_h$ and considering the geodesic from x to Y_h .

(ii) Similar to (i): choose x_1 and y_1 as close to each other as possible (on the geodesic joining x and y) subject to $a \in A_{x_1}$ and $b \in A_{y_1}$.

(iii) First note that $Y_h \cap Y_w \neq \emptyset$, for otherwise a consideration of the geodesic from Y_h to Y_w gives the contradiction $h^w \notin C$. Thus $a = h^w \in A(Y_w)$. The proof is by induction on r , where $w = a_1 \cdots a_r$. Consider the first step in the reduction process: if h^{a_1} is not an edge element then h^w is reduced as written (in the amalgamated free product of $A(\Delta_2)$ and A_{x_1}) and so cannot be a vertex element. Therefore $h_1 = h^{a_1} \in D$, and $a = h_1^{a_2 \cdots a_r}$. The result follows by induction on r (using part (i)).

(iv) Suppose $a^w = h \in D$ for some nonvertex element $w = a_1 \cdots a_r$, in reduced form. Then $h^{a_r^{-1} \cdots a_1^{-1}} = a \in C$, and by part (iii) we get $a^{a_1} = h^{a_r^{-1} \cdots a_2^{-1}} \in D$. So $a^A \cap D \neq \emptyset$ implies that $a^b \in D$ for some vertex element b . By part (ii), a and b must belong to the same vertex group, which has to be A_x since a belongs to no other vertex group. But now we have $a^{A_x} \cap D \neq \emptyset$, a contradiction. It follows that $a^A \cap D = \emptyset$.

For the second part let $b = a^w \in A_y$, and choose y as close to x as possible subject to $b \in A_y$. Claim that $y = x$. For if not, then $Y_w \neq \{x\}$, and we have two cases to consider:

Case 1. $x \notin VY_w$. Then there exists a finite subtree $\Delta \supseteq Y_w$ and an edge yx with $y \in V\Delta$. Now $a \in A_x \setminus H_{xy}$, $w \in A(\Delta)$, and $b = w^{-1}aw \in A(\Delta)$ (as Δ has to contain y because $b = a^w \in A(\Delta \cup \{x\})$ and $y \neq x$). This forces $w \in H_{yx} = H_{xy}$, and $a^w \in A(\Delta) \cap A_x = H_{xy}$, contradicting $a^A \cap D = \emptyset$.

Case 2. $x \in VY_w$. Let $R = \{\Delta_1, \dots, \Delta_m\}$ be a reduction process from Y_w to a point, such that the first vertex deleted is $x_1 \neq x$ (this can be done since $Y_w \neq \{x\}$). Write $w = a_1 \cdots a_r$ in reduced form relative to R . Then $b = a_r^{-1} \cdots a_1^{-1} a a_1 \cdots a_r \in A_y \subseteq A(\Delta_2) \cup A(x_1)$, so we must have $a_1 \in A_x$ and $a^{a_1} \in H_{y_1 x_1}$ (since $r \geq 2$). This is a contradiction.

We have now shown that $b = a^w \in A_x$. An argument similar to that in Case 2 now shows that $w \in A_x$, so $a^A \cap C \subseteq a^{A_x}$, are required. $\square$

We can now prove

Theorem 4.6. *Let $g \in A_x$ be such that $g^{A_x} \cap D = \emptyset$. Assume that X is locally finite, that every edge subgroup of A is compact in the I -topology, and that $\bigcap_{P \in I} g^{A_x} P_x = g^{A_x}$. Then g is conjugacy distinguished in A .*

Proof. If $\text{cl}_A(g^A)$ contains a nonvertex cyclically reduced element g_0 , then by 4.4 we have $\text{cl}_A(g^A) = \text{cl}_A(g_0^A) = g_0^A$, so $g^A = g_0^A = \text{cl}_A(g^A)$ and we are done. (Note that condition (1) is a consequence of the compactness assumption, so 4.4 is applicable.) We may therefore suppose that $\text{cl}_A(g^A) = \bigcup_i A_i^A$, where each A_i is a vertex element of A . Let $S = \bigcup_{i(e)=x} H_e$, and note that S is compact since X is locally finite. Put $T_P = g^{A_x} \pi_P \cap S \pi_P$. We claim that there exists P_0 such that $T_P = \emptyset$ for all $P \subseteq P_0$. Suppose not. Note that $Q \subseteq P$ implies that T_Q maps into T_P under the obvious map (from A_Q to A_P), so if $T_P \neq \emptyset$ then we have an inverse system of nonempty finite sets. Let $(h_P \pi_P) \in \varprojlim T_P$, where each $h_P \in S$. Since S is compact there exists $h \in S$ such that $h \pi_P = h_P \pi_P$ for all P . This clearly means that $h \in \bigcap_{P \in I} g^{A_x} P_x = g^{A_x}$, contradicting $g^{A_x} \cap D = \emptyset$. Hence there exists P_0 such that $g^{A_x} \pi_P \cap D \pi_P = \emptyset$ for all $P \subset P_0$. By 4.5(iv)

we have $g^A \pi_P \cap D \pi_P = \emptyset$, and $a_i \pi_P \in g^A \pi_P \cap C \pi_P = g^{A_x} \pi_P$. Thus first of all $a_i \in \bigcap_{P \subseteq P_0} A_x \ker \pi_P = A_x$, and then $a_i \in \bigcap_{P \subseteq P_0} g^{A_x} P_x = g^{A_x}$. Thus $\text{cl}_A(g^A) = g^A$, as required. $\square$

Remark 1. For g as in 4.6, the condition $\bigcap_P g^{A_x} P_x = g^{A_x}$ is implied by $\text{cl}_A(g^A) = g^A$. For if $b \in \bigcap g^{A_x} P_x$, then $b \in \text{cl}_A(g^A) = g^A$, and so $b \in g^A \cap C = g^{A_x}$ by 4.5(iv).

Remark 2. The usual proof of the conjugacy separability of the profinite group $\widehat{A_x}$ in fact shows that for $g \in \widehat{A_x}$ we have $\bigcap_{P \in I} g^{\widehat{A_x}} \overline{P}_x = g^{\widehat{A_x}}$.

The next two results give partial information about conjugacy separability of edge elements.

Lemma 4.7. *Assume that (1) holds. Let h be an edge element of A , and assume that there exists $P_0 \in I$ such that $Y_{h\pi_{P_0}} = Y_h$. Then $\text{cl}_A(h^A) \cap D = \text{cl}_{A(Y_h)}(h^{A(Y_h)}) \cap D$.*

Proof. The condition implies that $h\pi_P \notin H_e \pi_P$ for all $P \subseteq P_0$, where e is an edge leading out of Y_h . Let $z \in \text{cl}_A(h^A) \cap D$, so $z\pi_P = h^w \pi_P$ for some $w \in A$. We claim that w can be chosen in $A(Y_h)$. For if not, let Δ be the smallest tree containing $h\pi_P$ and $w\pi_P$, and consider a reduction process $\{\Delta_1, \dots, \Delta_m\}$ from Δ to $\Delta \cap Y_h$. Now the first vertex $x_1 \notin VY_h$, and $w\pi_P \notin A(\Delta_2)\pi_P$ by the minimal choice of Δ , and yet $(h\pi_P)^{w\pi_P}$ belongs to a vertex group. By the conjugacy theorem for amalgamated free product, we must have $h\pi_P \in H_{y_1 x_1} \pi_P$, contradicting the choice of P . We have therefore shown that

$$h_0 \in \bigcap_{P \subseteq P_0} h^{A(Y_h)} \ker \pi_P \subseteq \bigcap_{P \subseteq P_0} A(Y_h) \ker \pi_P = A(Y_h),$$

and so

$$\begin{aligned} h_0 &\in \bigcap_{P \subseteq P_0} h^{A(Y_h)} \ker \pi_P \cap A(Y_h) \\ &= \bigcap_{P \subseteq P_0} h^{A(Y_h)} (\ker \pi_P \cap A(Y_h)) = \text{cl}_{A(Y_h)}(h^{A(Y_h)}). \quad \square \end{aligned}$$

Let h be an edge element of A . For any sequence $\sigma = (x_1, \dots, x_r)$ of vertices of A put $D_\sigma(h) = D \cap \{h^{a_1 \dots a_r} : a_i \in A_{x_i}\}$. We have

Theorem 4.8. *Assume that X is locally finite and D is compact. Let h be an edge element of A^+ . Then h is conjugacy distinguished in A^+ if and only if $h^{A^+} \cap D = D_\sigma(h)$ for some finite sequence σ of vertices.*

Proof. We know that $\text{cl}_{A^+}(h^{A^+}) = \bigcup h_\alpha^{A^+}$ with each $h_\alpha \in D$. Let $K = h^{A^+} \cap D = \bigcup_\sigma D_\sigma(h)$, the union being over all finite sequences σ . If $z \in \text{cl}_{A^+}(h^{A^+}) \cap D$ and $P \in I$ then there exists a sequence $\sigma(P)$ such that $z\pi_P \in D_{\sigma(P)}(h)\pi_P \subseteq K\pi_P$, so $z \in K \ker \pi_P$. Thus

$$\text{cl}_{A^+}(h^{A^+}) \cap D \subseteq \bigcap_P K(\ker \pi_P) = \text{cl}_{A^+}(K).$$

Now D is a compact subset of A^+ , and so is closed. Therefore $\text{cl}_{A^+}(K) = \text{cl}_D(K)$. Conversely if $z \in \text{cl}_D(K)$ and $P \in I$ then $z\pi_P \in K\pi_P$, so $z\pi_P \in$

$D_\sigma(h)\pi_P$ for some sequence σ (depending on P). In other words, $z \in \text{cl}_{A^+}(h^{A^+})$, and we have shown that $\text{cl}_{A^+}(h^{A^+}) \cap D = \text{cl}_D(h^{A^+} \cap D)$. We also note that for a fixed sequence σ , the set $D_\sigma(h)$ is a complete subspace of the compact set D (since each vertex group $\hat{A}_x$ is compact), and is therefore compact. So if $h^{A^+} \cap D = D_\sigma(h)$ then $\text{cl}_{A^+}(h^{A^+}) \cap D = \text{cl}_D(K) = K = h^{A^+} \cap D$, which implies that every $h_\alpha \in h^{A^+}$, and so h is conjugacy distinguished in A^+ . Conversely, suppose $\text{cl}_{A^+}(h^{A^+}) = h^{A^+}$. Then $K = h^{A^+} \cap D = \text{cl}_D(K)$, so the compact (since closed) set K is the (countable) union of the closed sets $D_\sigma(h)$. Enumerate the finite sequences $\{\sigma_1, \sigma_2, \dots\}$, and put $T_i = \bigcup_{1 \leq j \leq i} D_{\sigma_j}(h)$. Then $K = \bigcup_{i=1}^\infty T_i$, and each T_i is closed. By the Baire Category Theorem (or rather its proof, using compactness), some T_{i_0} has nonempty interior, and so contains a set of the form $hP_x \cap D$, where $h \in D$ and $P \in I$. Since $P_x \triangleleft_f A_x$ and K is compact, it is easy to see that K is the union of finitely many T_i , whence $K = T_j$ for some j . Let σ be the concatenation of the sequences $\sigma_1, \dots, \sigma_j$. Then it is clear that $D_{\sigma_i}(h) \subseteq D_\sigma(h)$ for $1 \leq i \leq j$, and so $h^{A^+} \cap D = K = T_j \subseteq D_\sigma(h)$, as required. $\square$

The following are immediate consequences of 4.7 and 4.8.

Corollary 4.9. *Let X be locally finite, and let h be an edge element of A^+ such that Y_h is finite. Then h is conjugacy distinguished in A^+ if and only if for some finite sequence σ of vertices of Y_h we have $h^{A^+} \cap D = D_\sigma(h)$. $\square$*

Corollary 4.10. *Let A^+ be the amalgamated free product of countably many groups. Then an element h of the associated subgroup H is conjugacy distinguished in A^+ if and only if $h^{A^+} \cap H = D_\sigma(h)$ for some finite sequence σ of vertices. $\square$*

5. CONJUGACY SEPARABILITY OF G^+

If X is not a tree then G^+ is a nontrivial HNN-extension over the base group $A^+ = \langle \hat{A}_x : x \in VX \rangle$. By the length $|g|$ of an element $g \in G^+$ we mean the number of t -symbols appearing in an HNN-reduced form for g . Thus $g = a_1 t_{e_1}^{\epsilon_1} \cdots t_{e_n}^{\epsilon_n} a_{n+1}$, where all $a_i \in A^+$, a_{n+1} is arbitrary, each $\epsilon_i = \pm 1$, each $e_i \in EX \setminus EY$ (where Y is a fixed maximal subtree of X), and if $e_{i-1} = e_i$ and $\epsilon_{i-1} + \epsilon_i = 0$ then $a_i \notin H_{e_i e_i}$ (we write H_{-e} instead of $H_{\bar{e}}$ for ease of notation). The reduced element g is cyclically reduced if $a_{n+1} = 1$ and $t_{e_n}^{\epsilon_n} a_1 t_{e_1}^{\epsilon_1} \cdots a_n$ is also reduced.

Theorem 5.1. *Let $g \in G^+$ be cyclically reduced and of length at least one. Then g is conjugacy distinguished in G^+ .*

Proof. Let z be a cyclically reduced element in $\text{cl}_{G^+}(g^{G^+})$. It is easy to see that the edge subgroups $\overline{H}_{ee}$ are closed in A^+ , so we can find $P_0 \in I$ such that $|g| = |g\pi_P|$, $|z| = |z\pi_P|$, and both $g\pi_P$ and $z\pi_P$ are cyclically reduced in G_P for all $P \subseteq P_0$. The conjugacy theorem for HNN-extensions (e.g. [3]) now implies that $|g\pi_P| = |z\pi_P|$, so $|g| = |z|$. Modulo the same P we know that $g\pi_P$ and some cyclic permutation of $z\pi_P$ have the same sequence of t -symbols with the same exponents. Replacing z by a suitable cyclic permutation we may assume that $g = a_1 t_{e_1}^{\epsilon_1} \cdots a_n t_{e_n}^{\epsilon_n}$ and $z = b_1 t_{e_1}^{\epsilon_1} \cdots b_n t_{e_n}^{\epsilon_n}$. For $P \subseteq P_0$ the conjugacy theorem in G_P implies the existence of elements $h_{0,P}, \dots, h_{n,P}$ such that, modulo P

$$\begin{aligned}
h_{0,P}b_1 &\equiv a_1h_{1,P}, & h_{1,P} &\in \overline{H}_{\varepsilon_1e_1}, \\
(h_{1,P}\overline{\theta}_{e_1}^{\varepsilon_1})b_2 &\equiv a_2h_{2,P}, & h_{2,P} &\in \overline{H}_{\varepsilon_2e_2}, \\
&\vdots \\
(h_{n-1,P}\overline{\theta}_{e_{n-1}}^{\varepsilon_{n-1}})b_n &\equiv a_nh_{n,P}, & h_{n,P} &\in \overline{H}_{\varepsilon_ne_n}, \\
(h_{n,P}\overline{\theta}_{e_n}^{\varepsilon_n}) &\equiv h_{0,P}, & \text{so } h_{0,P} &\in \overline{H}_{-\varepsilon_ne_n} = \overline{H}_{\varepsilon_0e_0}.
\end{aligned}$$

Since $\bigcup_{i=0}^n \overline{H}_{\varepsilon_ie_i}$ is compact, there exist elements $h_i \in \overline{H}_{\varepsilon_ie_i}$ such that $h_i\pi_P = h_{i,P}\pi_P$ for all P . Then we have the equations $h_0b_1 = a_1h_1$, $(h_1\overline{\theta}_{e_1}^{\varepsilon_1})b_2 = a_2h_2$, $\dots$, $h_n\overline{\theta}_{e_n}^{\varepsilon_n} = h_0$, and so $z = h_0^{-1}gh_0$. The result follows. $\square$

To deal with conjugacy separability of elements of A^+ in G^+ , we begin with the following trivial observation, which can be proved by using a length argument (as in the proof of 5.1) and the conjugacy theorem for HNN-extensions.

Lemma 5.2. *If $a \in A^+$ then $\text{cl}_{G^+}(a^{G^+}) = \bigcup a_\alpha^{G^+}$, where each $a_\alpha \in A^+$. In particular, a is conjugacy distinguished in G^+ if and only if $\text{cl}_{G^+}(a^{G^+}) \cap A^+ = a^{G^+} \cap A^+$. $\square$*

The conjugacy theorem for elements of the base of an HNN-extension states that if $a, b \in A^+$ are conjugate in G^+ , then there exist elements $h_1, \dots, h_m \in \bigcup_{e \in EX \setminus EY} \overline{H}_e$, such that $a \sim h_1$ in A^+ (i.e., conjugate in A^+), $b \sim h_m$ in A^+ , and for $1 \leq i \leq m-1$, either $h_{i+1} \sim h_i$ in A^+ or $h_{i+1} = h_i\theta_e$ for some $e \in EX \setminus EY$. Now we have

Proposition 5.3. *Let $a \in A^+$ be a nonvertex cyclically reduced element (relative to some reduction process of Y_a). Then $\text{cl}_{G^+}(a^{G^+}) \cap A^+ = a^{A^+}$, and so a is conjugacy distinguished in G^+ .*

Proof. Choose P_0 such that for all $P \subseteq P_0$ we have $Y_{a\pi_P} = Y_a$ and $a\pi_P$ is cyclically reduced (cf. the proof of 4.4). Then $(a\pi_P)^{A^+\pi_P} \cap D\pi_P = \emptyset$ by 4.5(iii) and using the fact that no nonvertex conjugate of an edge element can be cyclically reduced. Now let $b \in \text{cl}_{G^+}(a^{G^+}) \cap A^+$. Then $b\pi_P$ is conjugate to $a\pi_P$ in G_P . Since no $A^+\pi_P$ -conjugate of $a\pi_P$ can belong to an edge subgroup in G_P , we must have $b\pi_P$ conjugate to $a\pi_P$ in $A^+\pi_P$, and so $b \in \text{cl}_{A^+}(a^{A^+}) = a^{A^+}$, the latter by 4.4. The final part follows from 5.2. $\square$

For vertex elements of A^+ we have the following (recall that $C = \bigcup_{x \in VX} \widehat{A}_x$):

Proposition 5.4. *Assume that X is locally finite. Let a be a vertex element of A^+ , and let $b \in \text{cl}_{G^+}(a^{G^+}) \cap C$. Then there exists a sequence $c_0, c_1, \dots$ of elements of C such that (i) $c_0 = a$; (ii) for each $i \geq 1$, either $c_i \sim c_{i-1}$ in C , or $c_i = c_{i-1}\theta_{e_i}$ for some $e_i \in EX \setminus EY$; and (iii) $c_i \rightarrow b$ as $i \rightarrow \infty$.*

Proof. For $P \in I$ let $S_P(a, b)$ denote the set of all sequences $\sigma = \{a_{i,P}\pi_P : i \geq 0\}$, where each $a_{i,P} \in C$, $a_{0,P}\pi_P = a\pi_P$, $a_{i,P}\pi_P = b\pi_P$ for all sufficiently large i , and each $a_{i,P}\pi_P$ is either $C\pi_P$ -conjugate to, or the θ_e - p -image of, $a_{i-1,P}\pi_P$ (in other words, each $\sigma \in S_P(a, b)$ represents a possible conjugacy $a\pi_P \sim b\pi_P$ in G_P^+). For fixed i and P write $T_i(P)$ for the set of i th components of all $\sigma \in S_P(a, b)$. Suppose $a \in \widehat{A}_x$, and let Δ_i denote the (finite)

subtree of radius i with centre x . A simple inductive argument shows that $T_i(P) \subseteq \bigcup_{y \in V_{\Delta_i}} \hat{A}_y \pi_P$, so each $T_i(P)$ is finite. If $Q \subseteq P$ then every sequence in $S_Q(a, b)$ maps to a sequence in $S_P(a, b)$ under the induced map $G_Q^+ \rightarrow G_P^+$, and so $T_i(Q)$ maps into $T_i(P)$.

We construct the elements c_i inductively, beginning with $c_0 = a$. Assume that $c_1, \dots, c_{i-1}$ have been chosen subject to (ii) and $c_j \pi_P \in T_j(P)$ for all P , and $1 \leq j \leq i-1$. Let $K_P = \{c' \pi_P \in T_i(P) : \text{there exists } \sigma \in S_P(a, b) \text{ such that } \sigma_{i-1} = c_{i-1} \pi_P \text{ and } \sigma_i = c' \pi_P\}$. Then each $K_P \neq \emptyset$, and if $Q \subseteq P$ then K_Q maps into K_P under the induced map $G_Q \rightarrow G_P$. Let $(d_P \pi_P)$ belong to the inverse limit of the sets K_P . Then each $d_P \in \bigcup_{y \in V_{\Delta_i}} \hat{A}_y$, and the latter set is compact, so there exists $c_i \in \bigcup_{y \in V_{\Delta_i}} \hat{A}_y$ such that $c_i \pi_P = d_P \pi_P$ for all P . Therefore, for every P there exists $\sigma \in S_P(a, b)$ such that $\sigma_{i-1} = c_{i-1} \pi_P$ and $\sigma_i = c_i \pi_P$. Also $c_i \pi_P \in T_i(P)$ for all P . To verify (ii), note that for each P the element $c_i \pi_P$ is either conjugate to $c_{i-1} \pi_P$ by some element of $\bigcup_{y \in V_{\Delta_{i-1}}} \hat{A}_y \pi_P$, or is the image of $c_{i-1} \pi_P$ under some $\theta_{e,P}$, where $e \in E\Delta_i$. The number of possibilities is therefore finite and independent of P , and so at least one must hold for all P . If $c_i \pi_P = c_{i-1} \pi_P \theta_{e,P}$ for all P , then $c_{i-1} \in \overline{H}_e$ and we get $c_i \pi_P = (c_{i-1} \theta_e) \pi_P$ for all P , whence $c_i = c_{i-1} \theta_e$. If $c_i \pi_P \sim c_{i-1} \pi_P$ in $\bigcup_{y \in V_{\Delta_i}} \hat{A}_y \pi_P$ for all P , then $c_i \sim c_{i-1}$ in $\bigcup_{y \in V_{\Delta_i}} \hat{A}_y$, using compactness. This proves (ii) for c_i , and so the inductive construction goes through. Finally given P it is clear that $c_i \pi_P = b \pi_P$ for all sufficiently large i , so $c_i \rightarrow b$ as $i \rightarrow \infty$. $\square$

Corollary 5.5. *Let $a \in A^+$ be a vertex element. Then a is conjugacy distinguished in G^+ if and only if $a^{G^+} \cap C$ is a closed subset of A^+ .*

Proof. By 5.2 and the proof of 5.3 we see that $\text{cl}_{G^+}(a^{G^+}) = \bigcup a_\alpha^{G^+}$, where each $a_\alpha \in C$. By 5.4, for each a_α there exists a sequence $c_i \in C$ such that $c_i \sim a$ in G^+ and $c_i \rightarrow a_\alpha$ as $i \rightarrow \infty$. Thus each $a_\alpha \in \text{cl}_{A^+}(a^{G^+} \cap C)$, and so $\text{cl}_{G^+}(a^{G^+}) \cap C \subseteq \text{cl}_{A^+}(a^{G^+} \cap C)$. The reverse inclusion being trivial, we have $\text{cl}_{G^+}(a^{G^+}) \cap C = \text{cl}_{A^+}(a^{G^+} \cap C)$. The result follows. $\square$

Corollary 5.6. *Let a be a vertex element of A^+ , and let $D' = \bigcup_{e \in EX \setminus EY} \overline{H}_e$. If $a^{A^+} \cap D' = \emptyset$, then a is conjugacy distinguished in G^+ if and only if a is conjugacy distinguished in A^+ .*

Proof. Since no A^+ -conjugate of a belongs to an associated subgroup, we have $a^{G^+} \cap A^+ = a^{A^+}$ by the conjugacy theorem for HNN-extensions, and so $a^{G^+} \cap C = a^{A^+} \cap C$. The result follows from 5.5 and the fact that $\text{cl}_{A^+}(a^{A^+}) = \bigcup a_\alpha^{A^+}$, with all $a_\alpha \in C$ (this is a consequence of 4.4). $\square$

To deal with elements of $D' = \bigcup_{e \in EX \setminus EY} \overline{H}_e$ we need the following:

Lemma 5.7. *If D is compact, then so is $C = \bigcup_{x \in V_X} \hat{A}_x$.*

Proof. It is sufficient to prove that C is complete, since we can always regard C as a subspace of the profinite completion $\hat{A}^+$. So let $\{a_P : P \in I\}$ be a subset of C such that for all $Q \subseteq P$ we have $a_Q \pi_P = a_P \pi_P$. If every $a_P \in D$, then the compactness of D implies the existence of $h \in D$ such that $a_P \pi_P = h \pi_P$ for all P , and we are done. Now suppose that $a_{P_0} \pi_{P_0} \notin D \pi_{P_0}$ for some P_0 . Then $a_P \pi_P \notin D \pi_P$ for all $P \subseteq P_0$, for if $a_P \in \hat{A}_y \neq \hat{A}_x$ for some $P \subseteq P_0$,

then $a_{P_0}\pi_{P_0} = a_P\pi_{P_0} \in \widehat{A}_Y\pi_{P_0} \cap \widehat{A}_X\pi_{P_0} \subseteq D\pi_{P_0}$, contrary to assumption. But now the compactness of $\widehat{A}_X$ gives the result. $\square$

Consider the set of all finite sequences $\sigma = (x_1, e_1^{\varepsilon_1}, \dots, e_n^{\varepsilon_n}, x_{n+1})$, where $x_i \in VX$, $e_i \in EX \setminus EY$, and $\varepsilon_i = \pm 1$. For such a sequence let

$$B_\sigma(h) = C \cap \{h^g : g = a_1 t_{e_1^{\varepsilon_1}} \cdots t_{e_n^{\varepsilon_n}} a_{n+1}, a_i \in \widehat{A}_{x_i} \text{ for all } i\}.$$

Theorem 5.8. *Assume that X is locally finite and D is compact. Then $h \in D'$ is conjugacy distinguished in G^+ if and only if there exist finite sequences $\sigma_1, \dots, \sigma_m$ such that $h^{G^+} \cap C = \bigcup_{i=1}^m B_{\sigma_i}(h)$.*

Proof. By 5.5 the element h is conjugacy distinguished in G^+ if and only if $K = h^{G^+} \cap C$ is closed in A^+ . Since C is compact (by 5.7) and therefore closed, this is equivalent to K being closed in C , i.e., to K being compact. Now $K = \bigcup_\sigma B_\sigma(h)$, the union being over the countably many sequences σ . An application of Baire's Category Theorem (cf. the proof of 4.8) shows that K is compact if and only if it is the union of finitely many $B_{\sigma_i}(h)$. (One needs to show that each $B_\sigma(h)$ is closed. For this, use the compactness of the $\widehat{A}_X$ to show that $B_\sigma(h)$ is complete.) $\square$

REFERENCES

1. R. B. J. T. Allenby and C. Y. Tang, *Conjugacy separability of certain 1-relator groups with torsion*, J. Algebra **103** (1986), 619–637.
2. D. E. Cohen, *Combinatorial group theory: A topological approach*, Cambridge Univ. Press, 1989.
3. D. J. Collins, *Recursively enumerable degrees and the conjugacy problem*, Acta Math. **122** (1969), 115–160.
4. W. Dicks and M. J. Dunwoody, *Groups acting on graphs*, Cambridge Univ. Press, 1989.
5. J. L. Dyer, *Separating conjugates in free-by-finite groups*, J. London Math. Soc. (2) **20** (1979), 215–221.
6. —, *Separating conjugates in amalgamated free products and HNN extensions*, J. Austral. Math. Soc. Ser. A **29** (1980), 35–51.
7. D. Gildenhuys and L. Ribes, *Profinite groups and Boolean graphs*, J. Pure Appl. Algebra **12** (1978), 21–47.
8. P. J. Higgins, *The fundamental groupoid of a graph of groups*, J. London Math. Soc. (2) **13** (1976), 145–149.
9. W. Magnus, A. Karrass, and D. Solitar, *Combinatorial group theory*, Wiley-Interscience, 1966.
10. V. N. Remeslennikov, *Conjugacy in polycyclic groups*, Algebra and Logic **8** (1969), 404–411.
11. —, *Groups that are residually finite with respect to conjugacy*, Siberian Math. J. **12** (1971), 783–792.
12. L. Ribes, *Introduction to pro-finite groups and Galois cohomology*, Queen's Papers in Pure and Appl. Math., vol. 24, Queen's University, Kingston, Ontario, 1970.
13. J.-P. Serre, *Cohomologie Galoisienne*, Springer, Berlin, 1964.
14. —, *Trees*, Springer, Berlin, 1980.
15. M. Shirvani, *On residually finite HNN-extensions*, Arch. Math. **44** (1985), 110–115.
16. —, *On residually finite graph products*, J. Pure Appl. Algebra **49** (1987), 281–282.
17. —, *A converse to a residual finiteness theorem of G. Baumslag*, Proc. Amer. Math. Soc. **104** (1988), 703–706.
18. P. F. Stebe, *Conjugacy separability of certain free products with amalgamation*, Trans. Amer. Math. Soc. **156** (1971), 119–129.

19. —, *Conjugacy separability of certain Fuchsian groups*, Trans. Amer. Math. Soc. **163** (1972), 173–188.
20. P. A. Zalesskii and O. V. Mel'nikov, *Subgroups of profinite groups acting on trees*, Math. USSR-Sb. **63** (1989), 405–424.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA, T6G 2G1
CANADA