

WAVELETS IN WANDERING SUBSPACES

T. N. T. GOODMAN, S. L. LEE, AND W. S. TANG

ABSTRACT. Mallat's construction, via a multiresolution approximation, of orthonormal wavelets generated by a single function is extended to wavelets generated by a finite set of functions. The connection between multiresolution approximation and the concept of wandering subspaces of unitary operators in Hilbert space is exploited in the general setting. An example of multiresolution approximation generated by cardinal Hermite B -splines is constructed.

1. INTRODUCTION

Wavelets are functions generated by translating and dilating a function or a finite set of functions. They are useful in many areas of mathematics and theoretical physics [1, 2] and also in practical applications such as image and signal processing [13, 14]. Orthonormal bases of wavelets generated by one function have been constructed for various function spaces [11, 15, 16]. Recently, Mallat [12] has unified the construction of these bases for $L^2(\mathbf{R})$ via the multiresolution approximation which is defined to be a sequence of closed subspaces $(V_m)_{m \in \mathbf{Z}}$ of $L^2(\mathbf{R})$ satisfying the following properties:

$$(1.1) \quad V_m \subset V_{m+1}, \quad m \in \mathbf{Z},$$

$$(1.2) \quad \bigcup_{m \in \mathbf{Z}} V_m \text{ is dense in } L^2(\mathbf{R}) \quad \text{and} \quad \bigcap_{m \in \mathbf{Z}} V_m = \{0\},$$

$$(1.3) \quad f \in V_m \Leftrightarrow D_2 f \in V_{m+1}, \quad m \in \mathbf{Z},$$

where $D_a f(x) := f(ax)$, $x \in \mathbf{R}$, for any positive number a ,

$$(1.4) \quad f \in V_m \Leftrightarrow T_{2^{-m}n} f \in V_m, \quad (m, n) \in \mathbf{Z}^2,$$

where $T_\tau(x) := f(x - \tau)$, $x \in \mathbf{R}$, for any $\tau \in \mathbf{R}$,

$$(1.5) \quad \text{there exists an isomorphism } \mathcal{J}: V_0 \rightarrow l^2(\mathbf{Z}) \text{ which commutes with the action of } \mathbf{Z}.$$

For a multiresolution approximation $(V_m)_{m \in \mathbf{Z}}$, Mallat [12] has shown the existence of $\phi \in V_0$ such that $(T_n \phi)_{n \in \mathbf{Z}}$ is an orthonormal basis of V_0 . Furthermore, if

$$\phi_{m,n}(x) := \sqrt{2^m} \phi(2^m x - n), \quad (m, n) \in \mathbf{Z}^2,$$

Received by the editors April 30, 1991.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 41A15, 42C15, 47B37.

Key words and phrases. Wavelets, Hilbert space, wandering subspaces, unitary operators, Hermite B -splines.

then for each $m \in \mathbf{Z}$, $(\phi_{m,n})_{n \in \mathbf{Z}}$ is an orthonormal basis of V_m . Let W_m be the orthogonal complement of V_m in V_{m+1} . Mallat has also shown the existence of $\psi \in W_0$ such that $(T_n \psi)_{n \in \mathbf{Z}}$ is an orthonormal basis of W_0 , and if

$$\psi_{m,n}(x) := \sqrt{2^m} \psi(2^m x - n), \quad (m, n) \in \mathbf{Z}^2,$$

then $(\psi_{m,n})_{n \in \mathbf{Z}}$ is a complete orthonormal set in W_m .

Properties (1.1) and (1.2) of the multiresolution approximation imply that

$$W_j \perp W_k, \quad j \neq k,$$

and

$$L^2(\mathbf{R}) = \sum_{m \in \mathbf{Z}} \oplus W_m.$$

Since $(\psi_{m,n})_{n \in \mathbf{Z}}$ is an orthonormal basis of W_m , it follows that $(\psi_{m,n})_{(m,n) \in \mathbf{Z}^2}$ is an orthonormal basis of $L^2(\mathbf{R})$.

Our object is to extend Mallat's results to wavelets generated by a finite number of functions. In this connection we observe the relationship between wavelets and the concept of wandering subspaces in operator theory (see Halmos [6, Problem 155], and Robertson [17]). This provides a general setting to wavelets in Hilbert space where the translation and dilation operators are replaced by unitary operators. We prove the existence of orthonormal wavelet bases for V_m generated by a finite number of vectors, and using a result of Robertson [17], we also prove the existence of bases of orthonormal wavelets for W_m and derive other results which are reminiscent of multiresolution approximation. This is done in §2. A more detailed analysis of the corresponding results in $L^2(\mathbf{R})$ is given in §3. An example of multiresolution approximation generated by cardinal Hermite B -splines (see Schoenberg [18, Chapter 5]) is constructed in §4. In §5 we construct compactly supported Hermite spline wavelets which extends the results of Chui and Wang [4].

2. EXISTENCE OF ORTHONORMAL WAVELETS AND WANDERING SUBSPACES

Let T be a unitary operator on a complex Hilbert space \mathcal{H} . A subspace \mathcal{S} of \mathcal{H} is called a *wandering subspace* for T if $T^m(\mathcal{S}) \perp T^n(\mathcal{S})$ for all $m \neq n$ (see [6, 17]). Further, if $V = \sum_{n \in \mathbf{Z}} \oplus T^n(\mathcal{S})$, then we say that \mathcal{S} is a complete wandering subspace of V for T . We shall denote by E the forward shift operator on $l^2(\mathbf{Z})$.

We begin by stating the Fuglede-Putnam Theorem in the form which will be needed later.

Lemma 2.1. *Let \mathcal{H}_i , $i = 1, 2$, be complex Hilbert spaces, and $A_i: \mathcal{H}_i \rightarrow \mathcal{H}_i$, $i = 1, 2$, be normal operators. If $B: \mathcal{H}_2 \rightarrow \mathcal{H}_1$, is a bounded operator such that $A_1 B = B A_2$, then $A_1^* B = B A_2^*$. If in addition B is invertible, then there exists a unitary operator $U: \mathcal{H}_2 \rightarrow \mathcal{H}_1$ such that $A_1 U = U A_2$.*

When $\mathcal{H}_1 = \mathcal{H}_2$, a proof of Fuglede-Putnam Theorem can be found in [6, Solution 192]. The proof with obvious modifications works for the general case.

Theorem 2.1. *Let V_0 be a closed subspace of a complex Hilbert space \mathcal{H} , and T a unitary operator on \mathcal{H} such that*

$$(2.1) \quad T V_0 = V_0.$$

Suppose that there exists a bounded invertible operator $J: V_0 \rightarrow l^2(\mathbf{Z})$ such that

$$(2.2) \quad JT = E^r J$$

for some $r \in \mathbf{Z} \setminus \{0\}$, and a unitary operator D on \mathcal{H} such that

$$(2.3) \quad V_0 \subset DV_0,$$

and

$$(2.4) \quad TD = DT^k,$$

for some $k \in \mathbf{Z}$, $|k| > 1$. Let

$$(2.5) \quad V_m := D^m V_0, \quad m \in \mathbf{Z}.$$

Then the following results hold:

(1) For every $m \in \mathbf{Z}$, $V_m \subset V_{m+1}$.

(2) There exist $\phi_j \in \mathcal{H}$, $1 \leq j \leq |r|$, such that

$$\{T^n \phi_j: n \in \mathbf{Z}, 1 \leq j \leq |r|\}$$

is an orthonormal basis of V_0 .

(3) The set

$$\{T^n DT^l \phi_j: n \in \mathbf{Z}, 0 \leq l \leq |k| - 1, 1 \leq j \leq |r|\}$$

is an orthonormal basis of V_1 .

(4) There exist ψ_j , $1 \leq j \leq |r|(|k| - 1)$, such that

$$\{T^n \psi_j: n \in \mathbf{Z}, 1 \leq j \leq |r|(|k| - 1)\}$$

is an orthonormal basis of the orthogonal complement W_0 of V_0 in V_1 .

(5) The collection

$$\{D^m T^n \psi_j: (m, n) \in \mathbf{Z}^2, 1 \leq j \leq |r|(|k| - 1)\}$$

is an orthonormal basis of $\overline{\bigcup_{m \in \mathbf{Z}} V_m} \ominus \bigcap_{m \in \mathbf{Z}} V_m$.

Remark 1. The result (2) of Theorem 2.1 is equivalent to the existence of a complete $|r|$ -dimensional wandering subspace of V_0 for T . Similarly, (4) is equivalent to the existence of a complete $|r|(|k| - 1)$ -dimensional wandering subspace of W_0 for T .

Remark 2. For a nonzero integer r , condition (2.2) is equivalent to the existence of a bounded invertible operator $\tilde{J}: V_0 \rightarrow l^2(\mathbf{Z})^{|r|}$ such that $\tilde{J}T = \tilde{E}\tilde{J}$, where $\tilde{E}(s_1, \dots, s_{|r|}) = (E^\varepsilon s_1, \dots, E^\varepsilon s_{|r|})$, $(s_1, \dots, s_{|r|}) \in l^2(\mathbf{Z})^{|r|}$, $\varepsilon := r/|r|$.

Remark 3. If $\mathcal{H} = L^2(\mathbf{R})$, and $T = T_1$ and $D = D_2$ are the translation and dilation operators respectively on $L^2(\mathbf{R})$, then T and D satisfy condition (2.4) with $k = 2$. In this case Theorem 2.1, with $r = 1$, reduces to the results of Mallat [12] on the existence of orthonormal wavelet bases for V_m and its orthogonal complement W_m in V_{m+1} corresponding to a multiresolution approximation $(V_m)_{m \in \mathbf{Z}}$ of $L^2(\mathbf{R})$.

Proof of Theorem 2.1. It is clear from the definition that $V_m \subset V_{m+1}$, $m \in \mathbf{Z}$.

Let $e_j \in l^2(\mathbf{Z})$ such that $e_j(n) = \delta_{jn}$, $j, n \in \mathbf{Z}$. Then, $\{(E^r)^n e_j: n \in \mathbf{Z}, 1 \leq j \leq |r|\}$ is an orthonormal basis of $l^2(\mathbf{Z})$. Since T and E^r are normal (indeed unitary) operators on V_0 and $l^2(\mathbf{Z})$ respectively, and they are similar by

(2.2), it follows from Fuglede-Putnam Theorem (Lemma 2.1), that there exists a unitary operator $U: V_0 \rightarrow l^2(\mathbf{Z})$ such that $UT = E^r U$. If $\phi_j := U^{-1}e_j$, $1 \leq j \leq |r|$, then $\{T^n \phi_j: n \in \mathbf{Z}, 1 \leq j \leq |r|\}$ is an orthonormal basis of V_0 . Hence $\mathcal{X} := \text{span}\{\phi_j: 1 \leq j \leq |r|\}$ is an $|r|$ -dimensional complete wandering subspace of V_0 for T . This proves (2).

Since D is a unitary operator on \mathcal{X} , and $V_m := D^m V_0$, $\{D^m T^n \phi_j: n \in \mathbf{Z}, 1 \leq j \leq |r|\}$ is an orthonormal basis of V_m . In particular, since $T^n D = D T^{nk}$, $n \in \mathbf{Z}$, the result (3) follows.

Let $\mathcal{Y} := \text{span}\{DT^l \phi_j: 0 \leq l \leq |k| - 1, 1 \leq j \leq |r|\}$. Then by (3), \mathcal{Y} is an $|rk|$ -dimensional complete wandering subspace of V_1 for T . By (2.3),

$$\sum_{n \in \mathbf{Z}} \oplus T^n(\mathcal{X}) \subset \sum_{n \in \mathbf{Z}} \oplus T^n(\mathcal{Y}).$$

By a theorem of Robertson [17, Theorem 2, p. 235], there exist vectors ψ_j , $1 \leq j \leq |r|(|k| - 1)$, such that $\{T^n \psi_j: n \in \mathbf{Z}, 1 \leq j \leq |r|(|k| - 1)\}$ is an orthonormal basis of the orthogonal complement W_0 of V_0 in V_1 .

Finally, letting W_m be the orthogonal complement of V_m in V_{m+1} , a standard argument shows that $W_m = D^m(W_0)$, $m \in \mathbf{Z}$, and

$$\overline{\bigcup_{m \in \mathbf{Z}} V_m} \ominus \bigcap_{m \in \mathbf{Z}} V_m = \sum_{m \in \mathbf{Z}} \oplus W_m.$$

Hence, for each m , $\{D^m T^n \psi_j: n \in \mathbf{Z}, 1 \leq j \leq |r|(|k| - 1)\}$ is an orthonormal basis of W_m and the collection $\{D^m T^n \psi_j: (m, n) \in \mathbf{Z}^2, 1 \leq j \leq |r|(|k| - 1)\}$ is an orthonormal basis of $\overline{\bigcup_{m \in \mathbf{Z}} V_m} \ominus \bigcap_{m \in \mathbf{Z}} V_m$. \square

Remark 4. In Theorem 2.1, it was assumed that (2.4) holds with $|k| \neq 1$. Suppose that it holds with $|k| = 1$, and all the other hypotheses of the theorem are satisfied. Then as in the proof of the theorem, \mathcal{X} and \mathcal{Y} are both $|r|$ -dimensional wandering subspaces for T , and

$$V_0 = \sum_{n \in \mathbf{Z}} \oplus T^n(\mathcal{X}) \subset \sum_{n \in \mathbf{Z}} \oplus T^n(\mathcal{Y}) = V_1.$$

By Theorem 1 of [17], this implies that $V_0 = V_1$. Hence $W_0 = \{0\}$, and $V_m = V_0$ for all $m \in \mathbf{Z}$.

3. WAVELETS GENERATED BY A FINITE SET OF FUNCTIONS IN $L^2(\mathbf{R})$

Let r be a positive integer and

$$l^2(\mathbf{Z})^r := \{(s_1, \dots, s_r): s_j \in l^2(\mathbf{Z}), j = 1, \dots, r\}.$$

For $s = (s_1, \dots, s_r) \in l^2(\mathbf{Z})^r$, its norm is given by

$$\|s\| := \left(\sum_{j=1}^r \|s_j\|^2 \right)^{1/2}.$$

The space of 2π -periodic square integrable functions and the space of 2π -periodic continuous functions will be denoted by $\tilde{L}^2(0, 2\pi)$ and $\tilde{C}(0, 2\pi)$ respectively. The Fourier transform of $f \in L^2(\mathbf{R})$ will be denoted by \hat{f} .

Let $g_j \in L^2(\mathbf{R})$, $j = 1, \dots, r$, such that $\sum_{n \in \mathbf{Z}} s(n) T_n g_j \in L^2(\mathbf{R})$, for all $s = (s(n))_{n \in \mathbf{Z}} \in l^2(\mathbf{Z})$. Then $\sum_{n \in \mathbf{Z}} |\hat{g}_j(u + 2\pi n)|^2$ is integrable, and for almost all $u \in \mathbf{R}$, $(\hat{g}_j(u + 2\pi n))_{n \in \mathbf{Z}} \in l^2(\mathbf{Z})$. Let

$$(3.1) \quad \hat{g}_{ij}(u) := \sum_{n \in \mathbf{Z}} \hat{g}_i(u + 2\pi n) \overline{\hat{g}_j(u + 2\pi n)}, \quad i, j = 1, \dots, r,$$

and

$$(3.2) \quad \hat{G}(u) := (\hat{g}_{ij}(u))_{i,j=1}^r.$$

Then \hat{G} is Hermitian, and therefore there is a unitary matrix U such that

$$(3.3) \quad \hat{G} = U^* D U,$$

where U^* denotes the conjugate transpose of U ,

$$D(u) := \text{diag}(\lambda_1(u), \dots, \lambda_r(u)),$$

and λ_j are the eigenvalues of \hat{G} .

Theorem 3.1. Let $K: l^2(\mathbf{Z})^r \rightarrow L^2(\mathbf{R})$ be defined by

$$(3.4) \quad K(s) := \sum_{j=1}^r \sum_{n \in \mathbf{Z}} s_j(n) T_n g_j, \quad s := (s_1, \dots, s_r) \in l^2(\mathbf{Z})^r.$$

Then K is an isomorphism onto a subspace of $L^2(\mathbf{R})$ if and only if there exist positive constants A and B such that

$$(3.5) \quad A \leq \lambda_j(u) \leq B, \quad j = 1, \dots, r, \text{ almost everywhere.}$$

Furthermore, $\{T_n g_j: n \in \mathbf{Z}, j = 1, \dots, r\}$ is an orthonormal subset of $L^2(\mathbf{R})$ if and only if K is an isometry, and these hold if and only if the matrix $\hat{G}(u)$ is the identity for almost all u .

Proof. For $s = (s_1, \dots, s_r) \in l^2(\mathbf{Z})^r$,

$$\widehat{K(s)}(u) = \sum_{j=1}^r h_j(u) g_j(u)$$

where

$$(3.6) \quad h_j \in \tilde{L}^2(0, 2\pi) \quad \text{with} \quad \|h_j\| = \|s_j\|.$$

Therefore,

$$\|\widehat{K(s)}\|^2 = \int_{-\infty}^{\infty} \sum_{i=1}^r \sum_{j=1}^r h_i(u) \overline{h_j(u)} \hat{g}_i(u) \overline{\hat{g}_j(u)} du.$$

Decomposing the integral over the intervals $[2\pi n, 2\pi(n+1))$, $n \in \mathbf{Z}$, and using (3.1) leads to

$$\|\widehat{K(s)}\|^2 = \int_0^{2\pi} \sum_{i=1}^r \sum_{j=1}^r h_i(u) \overline{h_j(u)} \hat{g}_{ij}(u) du.$$

The integrand of the last integral is a quadratic form. Using the Parseval identity and letting $H(u) := (h_1(u), \dots, h_r(u))^*$, it can be written as

$$(3.7) \quad \|K(s)\|^2 = \frac{1}{2\pi} \int_0^{2\pi} H^* \hat{G} H(u) du.$$

By (3.3), $H^* \widehat{G} H = Q^* D Q$, where

$$Q = (q_1, \dots, q_r)^* := UH,$$

with

$$(3.8) \quad \sum_{j=1}^r \|q_j\|^2 = \sum_{j=1}^r \|h_j\|^2,$$

since U is unitary. Therefore (3.7) becomes

$$\|K(s)\|^2 = \frac{1}{2\pi} \int_0^{2\pi} Q^* D Q(u) du = \frac{1}{2\pi} \int_0^{2\pi} \sum_{j=1}^r |q_j(u)|^2 \lambda_j(u) du.$$

Now, K is an isomorphism if and only if there exist positive constants A and B such that

$$A\|s\|^2 \leq \|K(s)\|^2 \leq B\|s\|^2, \quad s \in l^2(\mathbf{Z})^r.$$

By (3.6) and (3.8), this is equivalent to

$$\begin{aligned} A \sum_{j=1}^r \int_0^{2\pi} |q_j(u)|^2 du &\leq \sum_{j=1}^r \int_0^{2\pi} |q_j(u)|^2 \lambda_j(u) du \\ &\leq B \sum_{j=1}^r \int_0^{2\pi} |q_j(u)|^2 du \end{aligned}$$

for all $q_j \in L^2(0, 2\pi)$, $j = 1, \dots, r$, which in turn is equivalent to (3.5).

The last assertion of the theorem is an easy consequence of the above relations. \square

A function $f \in L^2(\mathbf{R})$ is said to be *regular* if \hat{f} is continuous and satisfies $\hat{f}(u) = O(u^{-1})$ as $|u| \rightarrow \infty$.

Corollary 3.1. *Suppose that g_j , $j = 1, \dots, r$, are regular. Then K is an isomorphism onto a subspace of $L^2(\mathbf{R})$ if and only if $\widehat{G}(u)$ is positive definite for all u .*

Proof. The assumption that g_j , $j = 1, \dots, r$, are regular implies that \hat{g}_{ij} , $i, j = 1, \dots, r$, are continuous. If $\widehat{G}(u)$ is positive definite for all u , its eigenvalues $\lambda_j(u)$ are positive for all u and $j = 1, \dots, r$. Since λ_j are 2π -periodic and continuous, there exist positive constants A and B such that (3.5) hold for all u . The converse is immediate. \square

Theorem 3.2. *Suppose that g_j , $j = 1, \dots, r$, are regular. Then K is an isomorphism onto a subspace of $L^2(\mathbf{R})$ if and only if for each u , the infinite matrix*

$$(3.9) \quad M(u) := (\hat{g}_i(u + 2\pi j))_{i=1, j \in \mathbf{Z}}^r$$

has rank r .

Proof. By (3.2) and (3.9), $\widehat{G} = M M^*$. For any $H = (h_1, \dots, h_r)^* \in \mathbf{C}^r$,

$$H^* \widehat{G} H = (H^* M)(H^* M)^* = \sum_{j \in \mathbf{Z}} \left| \sum_{i=1}^r h_i \hat{g}_i(u + 2\pi j) \right|^2.$$

Therefore, for any $H \neq 0$, $H^* \widehat{G} H > 0$ if and only if $\sum_{i=1}^r h_i \hat{g}_i(u + 2\pi j)$ for some j , if and only if $\text{rank}(M) = r$. \square

In the next theorem, the notation C_j means the j th column of the matrix C .

Theorem 3.3. *Suppose that K is an isomorphism. Let*

$$D^{1/2}(u) := \text{diag}(\lambda_1(u)^{1/2}, \dots, \lambda_r(u)^{1/2}),$$

and

$$(3.10) \quad (\hat{\phi}_1(u), \dots, \hat{\phi}_r(u))^T := ((U^* D^{1/2})^{-1} M)_0.$$

Then $\{T_n \phi_j : n \in \mathbf{Z}, j = 1, \dots, r\}$ is an orthonormal set.

Proof. Since U^* and $D^{1/2}$ are 2π -periodic, by (3.10)

$$(\hat{\phi}_i(u + 2\pi j))_{i=1, j \in \mathbf{Z}}^r = (U^* D^{1/2})^{-1} M.$$

Let

$$(3.11) \quad \hat{\phi}_{ij}(u) := \sum_{n \in \mathbf{Z}} \hat{\phi}_i(u + 2\pi n) \overline{\hat{\phi}_j(u + 2\pi n)},$$

and $\hat{\Phi} := (\hat{\phi}_{ij})_{i,j=1}^r$. Then

$$\hat{\Phi} = (U^* D^{1/2})^{-1} M M^* (D^{1/2} U)^{-1}.$$

Since $M M^* = \widehat{G} = (U^* D^{1/2})(D^{1/2} U)$, it follows that $\hat{\Phi}(u)$ equals the identity matrix for almost all u . By Theorem 3.1, the map $K_1: l^2(\mathbf{Z})^r \rightarrow L^2(\mathbf{R})$ defined by

$$(3.12) \quad K_1(s) := \sum_{j=1}^r \sum_{n \in \mathbf{Z}} s_j(n) T_n \phi_j, \quad s \in l^2(\mathbf{Z})^r,$$

is an isometry onto a subspace of $L^2(\mathbf{R})$. Therefore $\{T_n \phi_j : n \in \mathbf{Z}, j = 1, \dots, r\}$, is an orthonormal set. \square

Remark 5. Suppose that K is an isomorphism onto a subspace V_0 of $L^2(\mathbf{R})$. Since $T_1 V_0 = V_0$ and $T_1 K = K E^r$, Theorem 2.1 also yields the existence of ϕ_j , $j = 1, \dots, r$, such that $\{T_n \phi_j : n \in \mathbf{Z}, j = 1, \dots, r\}$ is an orthonormal basis of V_0 .

Equation (3.10) is more of theoretical interest since the eigenvalues and eigenvectors of \widehat{G} are, in general, difficult to compute. We shall now give an inductive construction of the orthonormal wavelets which could be more useful in practice. Suppose that $\{\phi_j : 1 \leq j \leq r-1\}$, is a set of regular functions in $L^2(\mathbf{R})$ such that $\{T_n \phi_j : n \in \mathbf{Z}, 1 \leq j \leq r-1\}$ is an orthonormal set. Then by Theorem 3.1 the mapping $K_1: l^2(\mathbf{Z})^{r-1} \rightarrow L^2(\mathbf{R})$ defined by (3.12) with $r-1$ instead of r , is an isometry onto a subspace of $L^2(\mathbf{R})$, and $\hat{\Phi} := (\hat{\phi}_{ij})_{i,j=1}^{r-1}$, where $\hat{\phi}_{ij}$ are defined in (3.11), is the identity matrix.

Theorem 3.4. Suppose $g \in L^2(\mathbf{R})$ is regular. Then the map $K: l^2(\mathbf{Z})^r \rightarrow L^2(\mathbf{R})$, such that

$$(3.13) \quad K(s) := \sum_{j=1}^{r-1} \sum_{n \in \mathbf{Z}} s_j(n) T_n \phi_j + \sum_{n \in \mathbf{Z}} s_r(n) T_n g,$$

is an isomorphism onto a subspace V_0 of $L^2(\mathbf{R})$ if and only if for all $u \in \mathbf{R}$,

$$(3.14) \quad |\langle \hat{g}, \hat{g} \rangle(u)| > \sum_{j=1}^{r-1} |\langle \hat{g}, \hat{\phi}_j \rangle(u)|^2,$$

where

$$\langle \hat{g}, \hat{g} \rangle(u) := \sum_{n \in \mathbf{Z}} |\hat{g}(u + 2\pi n)|^2,$$

and

$$\langle \hat{g}, \hat{\phi}_j \rangle(u) := \sum_{n \in \mathbf{Z}} \hat{g}(u + 2\pi n) \overline{\hat{\phi}_j(u + 2\pi n)}.$$

Further, if (3.14) holds and

$$\begin{aligned} \tau(u) &:= (\langle \hat{g}, \hat{g} \rangle(u) - \sum_{j=1}^{r-1} |\langle \hat{g}, \hat{\phi}_j \rangle(u)|^2)^{-1/2}, \\ t_j(u) &:= -\tau(u) \langle \hat{g}, \hat{\phi}_j \rangle(u), \quad j = 1, \dots, r-1, \end{aligned}$$

and

$$(3.15) \quad \hat{\phi}_r := \sum_{j=1}^{r-1} t_j \hat{\phi}_j + \tau \hat{g},$$

then $\{T_n \phi_j: n \in \mathbf{Z}, j = 1, \dots, r\}$ is an orthonormal basis of V_0 .

Proof. Let $g_j := \phi_j$, $j = 1, \dots, r-1$, and $g_r := g$. Then \hat{G} is a matrix of order r whose last row is $(\hat{g}_{r1}, \dots, \hat{g}_{rr})$ and last column is $(\hat{g}_{r1}, \dots, \hat{g}_{rr})^*$, and the remaining submatrix is the identity. Clearly, \hat{G} is positive definite if and only if $|\hat{G}| > 0$ which is equivalent to (3.14).

If (3.14) holds then

$$V_0 = \left\{ f: f = \sum_{j=1}^{r-1} \sum_{n \in \mathbf{Z}} s_j(n) T_n \phi_j + \sum_{n \in \mathbf{Z}} s_r(n) T_n g \right\}.$$

Now $f \in V_0$ if and only if there are unique $h_j \in \tilde{L}^2(0, 2\pi)$, $j = 1, \dots, r$, with

$$\hat{f} = \sum_{j=1}^{r-1} h_j \hat{\phi}_j + h_r \hat{g}.$$

Therefore, we want to find $\tau, t_1, \dots, t_{r-1}$ in $\tilde{L}^2(0, 2\pi)$ such that, with

$$(3.16) \quad \hat{\phi}_r := \sum_{j=1}^{r-1} t_j \hat{\phi}_j + \tau \hat{g},$$

$\{T_n\phi_j: n \in \mathbf{Z}, j = 1, \dots, r\}$ is an orthonormal set. This is equivalent to

$$(3.17) \quad \langle \hat{\phi}_r, \hat{\phi}_j \rangle = \delta_{r,j}, \quad j = 1, \dots, r.$$

Since $\langle \hat{\phi}_i, \hat{\phi}_j \rangle = \delta_{ij}$, $i, j = 1, \dots, r-1$, it follows from (3.16) that (3.17) is equivalent to

$$(3.18) \quad t_j + \tau \langle \hat{g}, \hat{\phi}_j \rangle = 0, \quad j = 1, \dots, r-1,$$

and

$$(3.19) \quad \sum_{j=1}^{r-1} |t_j|^2 + |\tau|^2 \langle \hat{g}, \hat{g} \rangle + \bar{\tau} \sum_{j=1}^{r-1} t_j \langle \hat{\phi}_j, \hat{g} \rangle + \tau \sum_{j=1}^{r-1} \bar{t}_j \langle \hat{g}, \hat{\phi}_j \rangle = 1.$$

Substituting for t_1, \dots, t_{r-1} from (3.18) into (3.19) leads to

$$|\tau|^2 = (\langle \hat{g}, \hat{g} \rangle - \sum_{j=1}^{r-1} |\langle \hat{g}, \hat{\phi}_j \rangle|^2)^{-1}.$$

By (3.14), the function

$$(3.20) \quad \tau(u) := (\langle \hat{g}, \hat{g} \rangle(u) - \sum_{j=1}^{r-1} |\langle \hat{g}, \hat{\phi}_j \rangle(u)|^2)^{-1/2}$$

is 2π -periodic and continuous, and therefore belongs to $\tilde{L}^2(0, 2\pi)$. We then define

$$(3.21) \quad t_j(u) := -\tau(u) \langle \hat{g}, \hat{\phi}_j \rangle(u), \quad j = 1, \dots, r-1,$$

which belong to $\tilde{L}^2(0, 2\pi)$ and satisfy (3.18) and (3.19). So with ϕ_r defined by (3.16), (3.20) and (3.21), $\{T_n\phi_j: n \in \mathbf{Z}, j = 1, \dots, r\}$ is an orthonormal set in V_0 .

We denote the Fourier transform operator by \mathcal{F} and define $U: \mathcal{F}V_0 \rightarrow \mathcal{F}V_0$ such that

$$U\hat{f} := \sum_{j=1}^{r-1} h_j \hat{\phi}_j = h_r \hat{\phi}_r, \quad \hat{f} = \sum_{j=1}^{r-1} h_j \hat{\phi}_j + h_r \hat{g} \in V_0,$$

where $h_j \in \tilde{L}^2(0, 2\pi)$, $j = 1, \dots, r$. By (3.16) we see that U is an isomorphism of $\mathcal{F}V_0$ onto itself. Since $\{T_n\phi_j: n \in \mathbf{Z}, j = 1, \dots, r\}$ is orthonormal, it follows that the map $\tilde{K}: l^2(\mathbf{Z})^r \rightarrow V_0$ such that

$$\tilde{K}(s) := \sum_{j=1}^r \sum_{n \in \mathbf{Z}} s_j(n) T_n \phi_j, \quad s = (s_1, \dots, s_r) \in l^2(\mathbf{Z})^r,$$

is an isometry. But then $\tilde{K} = \mathcal{F}^{-1}U\mathcal{F}K$, so that \tilde{K} maps $l^2(\mathbf{Z})^r$ onto V_0 . Thus $\{T_n\phi_j: n \in \mathbf{Z}, j = 1, \dots, r\}$ is complete in V_0 . \square

Now suppose that $\phi_j \in L^2(\mathbf{R})$, $j = 1, \dots, r$, such that $\{T_n\phi_j: n \in \mathbf{Z}, j = 1, \dots, r\}$ is an orthonormal set, and let V_0 be its closed linear span in $L^2(\mathbf{R})$. Then

$$V_0 = \left\{ f \in L^2(\mathbf{R}): \hat{f} = \sum_{j=1}^r \hat{\phi}_j h_j, h_j \in \tilde{L}^2(0, 2\pi) \right\}.$$

Let $V_1 := D_2 V_0$. Then

$$V_1 = \left\{ f \in L^2(\mathbf{R}): \hat{f}(2u) = \sum_{j=1}^r \hat{\phi}_j(u) h_j(u), h_j \in \tilde{L}^2(0, 2\pi) \right\}.$$

Clearly, $T_n V_1 = V_1$, $n \in \mathbf{Z}$, and the following results are easy consequences.

Proposition 3.1. *The following conditions are equivalent:*

- (1) $V_0 \subset V_1$,
- (2) $\{\phi_1, \dots, \phi_r\} \subset V_1$,
- (3) *There exist $a_{ij} \in \tilde{L}^2(0, 2\pi)$, $i, j = 1, \dots, r$, such that*

$$(3.22) \quad \hat{\phi}_i(2u) = \sum_{j=1}^r \hat{\phi}_j(u) a_{ij}(u), \quad i = 1, \dots, r.$$

We shall henceforth assume that the conditions in Proposition 3.1 hold.

Proposition 3.2. *Suppose that $f_i \in V_1$, $i = 1, \dots, r$, and*

$$\hat{f}_i(2u) = \sum_{j=1}^r \hat{\phi}_j(u) h_{ij}(u), \quad h_{ij} \in \tilde{L}^2(0, 2\pi), \quad i, j = 1, \dots, r.$$

Let $H := (h_{ij})_{i,j=1}^r$. Then $\{T_n f_i: n \in \mathbf{Z}, i = 1, \dots, r\}$ is orthonormal if and only if

$$H(u)H(u)^* + H(u + \pi)H(u + \pi)^* = I_r,$$

where I_r denotes the identity matrix of order r .

Proof. For $i, j = 1, \dots, r$,

$$\begin{aligned} & \sum_{n \in \mathbf{Z}} \hat{f}_i(u + 2\pi n) \overline{\hat{f}_j(u + 2\pi n)} \\ &= \sum_{k=1}^r \sum_{l=1}^r h_{ik} \left(\frac{u}{2} \right) \overline{h_{jl} \left(\frac{u}{2} \right)} \sum_{n \in \mathbf{Z}} \hat{\phi}_k \left(\frac{u}{2} + 2\pi n \right) \overline{\hat{\phi}_l \left(\frac{u}{2} + 2\pi n \right)} \\ &+ \sum_{k=1}^r \sum_{l=1}^r h_{ik} \left(\frac{u}{2} + \pi \right) \overline{h_{jl} \left(\frac{u}{2} + \pi \right)} \sum_{n \in \mathbf{Z}} \hat{\phi}_k \left(\frac{u}{2} + \pi + 2\pi n \right) \overline{\hat{\phi}_l \left(\frac{u}{2} + \pi + 2\pi n \right)} \\ &= \sum_{k=1}^r h_{ik} \left(\frac{u}{2} \right) \overline{h_{jk} \left(\frac{u}{2} \right)} + \sum_{k=1}^r h_{ik} \left(\frac{u}{2} + \pi \right) \overline{h_{jk} \left(\frac{u}{2} + \pi \right)}, \end{aligned}$$

by Theorem 3.1 since $\{T_n \phi_j: n \in \mathbf{Z}, j = 1, \dots, r\}$ is an orthonormal set.

Therefore, $\{T_n f_i: n \in \mathbf{Z}, i = 1, \dots, r\}$ is orthonormal if and only if

$$\sum_{n \in \mathbf{Z}} \hat{f}_i(u + 2\pi n) \overline{\hat{f}_j(u + 2\pi n)} = \delta_{ij}, \quad i, j = 1, \dots, r,$$

if and only if

$$\sum_{k=1}^r h_{ik} \left(\frac{u}{2} \right) \overline{h_{jk} \left(\frac{u}{2} \right)} + \sum_{k=1}^r h_{ik} \left(\frac{u}{2} + \pi \right) \overline{h_{jk} \left(\frac{u}{2} + \pi \right)} = \delta_{ij},$$

$$i, j = 1, \dots, r, u \in \mathbf{R}. \quad \square$$

Let a_{ij} be defined as in (3.22), and $A := (a_{ij})_{i,j=1}^r$. By Proposition 3.2,

$$(3.23) \quad A(u)A(u)^* + A(u + \pi)A(u + \pi)^* = I_r.$$

Let W_0 be the orthogonal complement of V_0 in V_1 , and let $\psi_i \in V_1$, $i = 1, \dots, r$, such that

$$(3.24) \quad \hat{\psi}_i(2u) = \sum_{j=1}^r \hat{\phi}_j(u)b_{ij}(u)$$

where $b_{ij} \in \tilde{L}^2(0, 2\pi)$, $i, j = 1, \dots, r$. Let $B := (b_{ij})_{i,j=1}^r$. A similar argument as in the proof of Proposition 3.2 gives

Proposition 3.3. *The functions ψ_i , $i = 1, \dots, r$, are in W_0 if and only if*

$$(3.25) \quad A(u)B(u)^* + A(u + \pi)B(u + \pi)^* = 0.$$

Furthermore, $\{T_n\psi_i: n \in \mathbf{Z}, i = 1, \dots, r\}$ is an orthonormal set if and only if

$$(3.26) \quad B(u)B(u)^* + B(u + \pi)B(u + \pi)^* = I_r.$$

Given $A = (a_{ij})_{i,j=1}^r$ satisfying (3.23), the functions $\hat{\psi}_i$, $i = 1, \dots, r$, can be obtained by constructing $B = (b_{ij})_{i,j=1}^r$ so that (3.25) and (3.26) are satisfied. Indeed (3.23) is equivalent to the orthonormality of the vectors

$$(a_{i1}(u), \dots, a_{ir}(u), a_{i1}(u + \pi), \dots, a_{ir}(u + \pi)), \quad i = 1, \dots, r,$$

for $u \in [0, \pi)$. By Gram-Schmidt process they can be extended to an orthonormal basis of \mathbf{C}^{2r} . Let the new vectors in the basis be denoted by

$$(b_{i1}(u), \dots, b_{ir}(u), b_{i1}(u + \pi), \dots, b_{ir}(u + \pi)), \quad i = 1, \dots, r,$$

and extend b_{ij} , $i, j = 1, \dots, r$, periodically with period 2π . Then $B := (b_{ij})_{i,j=1}^r$ satisfies (3.25) and (3.26). Let ψ_i be defined as in (3.24).

Theorem 3.5. *The set $\{T_n\psi_i: n \in \mathbf{Z}, 1 \leq i \leq r\}$ is an orthonormal basis of W_0 . Furthermore,*

$$(3.27) \quad W_0 = \left\{ f \in L^2(\mathbf{R}): \hat{f} = \sum_{j=1}^r \hat{\psi}_j h_j, h_j \in \tilde{L}^2(0, 2\pi) \right\}.$$

Proof. By Proposition 3.3, $\{T_n\psi_i: n \in \mathbf{Z}, 1 \leq i \leq r\}$ is an orthonormal subset of W_0 . We have only to show that it is complete.

Take $\psi \in W_0$ and orthogonal to $\{T_n\psi_i: n \in \mathbf{Z}, 1 \leq i \leq r\}$. Then there exist g_1, \dots, g_r , in $\tilde{L}^2(0, 2\pi)$ such that

$$\hat{\psi}(2u) = \sum_{j=1}^r \hat{\phi}_j(u)g_j(u),$$

and by a similar argument as in the proof of Proposition 3.2,

$$\sum_{k=1}^r (a_{ik}(u)\overline{g_k(u)} + a_{ik}(u + \pi)\overline{g_k(u + \pi)}) = 0$$

and

$$\sum_{k=1}^r (b_{ik}(u)\overline{g_k(u)} + b_{ik}(u + \pi)\overline{g_k(u + \pi)}) = 0$$

for $i = 1, \dots, r$. This means that the vector

$$(g_1(u), \dots, g_r(u), g_1(u + \pi), \dots, g_r(u + \pi))$$

is orthogonal to the orthonormal basis of \mathbf{C}^{2r} . Therefore, all g_j , $j = 1, \dots, r$, and hence ψ are zero. This implies that $\{T_n \psi_i : n \in \mathbf{Z}, 1 \leq i \leq r\}$ is an orthonormal basis of W_0 .

Functions in W_0 can be written uniquely in the form

$$f = \sum_{j=1}^r \sum_{n \in \mathbf{Z}} c_{jn} T_n \psi_j, \quad (c_{jn})_{n \in \mathbf{Z}} \in l^2(\mathbf{Z}).$$

Taking Fourier transform leads to (3.27). \square

4. MULTIREOLUTION APPROXIMATION GENERATED BY CARDINAL HERMITE B -SPLINES

Let n and r be positive integers, n even, such that $2r \leq n$, and

$$(4.1) \quad \mathcal{S}_n^r := \{f \in C^{n-r-1}(\mathbf{R}) : f|_{[\nu, \nu+1]} \in \Pi_{n-1}, \nu \in \mathbf{Z}\},$$

where Π_{n-1} is the class of polynomials of degree $\leq n-1$. Functions in \mathcal{S}_n^r are called cardinal Hermite splines of degree $n-1$. For $j = 0, \dots, r-1$, let

$$(4.2) \quad \mathcal{S}_{n,j}^r := \{f \in \mathcal{S}_n^r : f^{(k)}(\nu) = 0, \nu \in \mathbf{Z}, k = 0, \dots, r-1, k \neq j\}.$$

The space $\mathcal{S}_{n,j}^r$ has a basis consisting of integer translates of a function $N_{n,j} = N_{n,j}^r \in \mathcal{S}_{n,j}^r$, $j = 0, \dots, r-1$, with minimal support $[-\frac{n}{2} - 1 + r, \frac{n}{2} + 1 - r]$, in the sense that every $f \in \mathcal{S}_{n,j}^r$ has a unique representation of the form

$$(4.3) \quad f(x) = \sum_{\nu \in \mathbf{Z}} c_\nu N_{n,j}(x - \nu), \quad x \in \mathbf{R}$$

(see [18, 8]). The functions $N_{n,j}$ are called cardinal Hermite B -splines.

For computational purposes, the Fourier transforms $\hat{N}_{n,j}$ of the cardinal Hermite B -splines are very useful. They can be expressed in closed forms with the help of the Hankel determinant $|H_r(a_n)|$ where $H_r(a_n)$ is the matrix of order r whose j th column is $(a_{n-j+1}, \dots, a_{n-j-r+2})^T$ (see [9]). Indeed if

$$(4.4) \quad \alpha_k(u) := \sum_{\nu \in \mathbf{Z}} (u + 2\pi\nu)^{-k}, \quad k = 2, \dots, n,$$

and $H_{r,j}(\alpha_n(u))$ denotes the matrix obtained from $H_r(\alpha_n(u))$ by replacing the $(j+1)$ th column by $(u^n, u^{n-1}, \dots, u^{n-r+1})^T$, $j = 0, \dots, r-1$, then

$$(4.5) \quad \hat{N}_{n,j}(u) = \left(2 \sin \frac{u}{2}\right)^n |H_{r,j}(\alpha_n(u))|.$$

For $j = 0, \dots, r-1$, $N_{n,j}$ are regular. Indeed, $\hat{N}_{n,j}$ are continuous and $\hat{N}_{n,j}(u) = O(u^{-n+r-1})$ as $n \rightarrow \infty$. Therefore the map $K : l^2(\mathbf{Z})^r \rightarrow L^2(\mathbf{R})$ such that

$$(4.6) \quad K(s) := \sum_{j=0}^{r-1} \sum_{\nu \in \mathbf{Z}} s_j(\nu) T^\nu N_{n,j},$$

is well defined.

Theorem 4.1. *The map K defined by (4.6) is an isomorphism of $l^2(\mathbf{Z})^r$ onto a subspace of $L^2(\mathbf{R})$.*

Proof. Let $\hat{N}(u) := (\hat{N}_{n,i}(u + 2\pi j))_{i=0, j \in \mathbf{Z}}^{r-1}$. Suppose that $0 < u < 2\pi$. By a result of [10], $|H_r(\alpha_n(u))| \neq 0$. Therefore, equation (4.5) can be expressed in the form

$$(\hat{N}_{n,i}(u))_{i=0}^{r-1} = (2 \sin u/2)^n |H_r(\alpha_n(u))| H_r(\alpha_n(u))^{-1} (u^{-n}, u^{-n+1}, \dots, u^{-n+r-1})^T.$$

Since $H_r(\alpha_n(u))$ is 2π -periodic, it follows that

$$(4.7) \quad (\hat{N}_{n,i}(u + 2\pi j))_{i=0}^{r-1, r-1} = (2 \sin u/2)^n |H_r(\alpha_n(u))| H_r(\alpha_n(u))^{-1} ((u + 2\pi j)^{-n+i})_{i=0, j=0}^{r-1, r-1}.$$

Since the last matrix on the right of (4.7) is a Vandermonde, we see that both sides are nonsingular. In particular $\hat{N}(u)$ is of rank r , if $0 < u < 2\pi$.

For $u = 0$, a straightforward computation shows that

$$(4.8) \quad (\hat{N}_{n,i}(2\pi j))_{i=0, j=1}^{r-1, r} = |H_{r-1}(\beta_{n-2})| H_{r-1}(\beta_{n-2})^{-1} ((2\pi j)^{-n+i})_{i=1, j=1}^{r-1, r-1},$$

where

$$\beta_k := \sum_{\nu \neq 0} (2\pi\nu)^{-k}.$$

A similar argument as above shows that $\hat{N}(0)$ is of rank r . The result now follows from Theorem 3.2. \square

Let $\widetilde{\mathcal{S}}_n^r$ denote the range of K . This is a closed subspace of $L^2(\mathbf{R})$. Furthermore, $\widetilde{\mathcal{S}}_n^r = \mathcal{S}_n^r \cap L^2(\mathbf{R})$. Therefore, if $Df(x) := f(2x)$,

$$\bigcup_{m \in \mathbf{Z}} D^m \widetilde{\mathcal{S}}_n^r \supseteq \overline{\bigcup_{m \in \mathbf{Z}} D^m \widetilde{\mathcal{S}}_n^1} = L^2(\mathbf{R}).$$

Also $\bigcap_{m \in \mathbf{Z}} D^m \widetilde{\mathcal{S}}_n^r = \{0\}$, since it comprises polynomials in $L^2(\mathbf{R})$.

Let $V_m := D^m \widetilde{\mathcal{S}}_n^r$, $m \in \mathbf{Z}$. Then $(V_m)_{m \in \mathbf{Z}}$ is a multiresolution approximation of $L^2(\mathbf{R})$ generated by the cardinal Hermite B -splines $N_{n,j}$, $j = 0, \dots, r-1$. We shall call $(V_m)_{m \in \mathbf{Z}}$, a Hermite spline multiresolution approximation of $L^2(\mathbf{R})$.

5. COMPACTLY SUPPORTED HERMITE SPLINE WAVELETS FOR THE ORTHOGONAL COMPLEMENT

Let W_0 be the orthogonal complement of V_0 in V_1 , so that $W_m := D^m W_0$ is the orthogonal complement of V_m in V_{m+1} , $m \in \mathbf{Z}$. For convenience, we shall assume that the B -spline $N_{n,j}$ is supported on $[0, n - 2r + 2]$. This can be easily achieved by translation.

Take $j = 0, \dots, r-1$, and consider functions of the form

$$f_j(x) = \sum_{\nu=0}^m c_j N_{2n,j}(2x - \nu), \quad \text{with } f_j^{(j)}(k) = 0, \quad k \in \mathbf{Z},$$

where m is a positive integer. Then $f_j^{(n)} \in V_1$ and integrating by parts, in the sense of distributions, shows that $f_j^{(n)}$ is orthogonal to V_0 , and hence belongs

to W_0 . Such a function f_j with minimal support is given by

$$(5.1) \quad f_j(x) = \sum_{\nu=0}^{2n-2r} (-1)^\nu N_{2n,j}^{(j)}(\nu+1) N_{2n,j}(2x-\nu).$$

For $j = 0, \dots, r-1$, let

$$\widetilde{\mathcal{F}}_{2n,j}^r := \{f \in \widetilde{\mathcal{F}}_{2n}^r : f^{(k)}(\nu) = 0, \nu \in \mathbf{Z}, k = 0, \dots, r-1, k \neq j\}.$$

Lemma 5.1. *Let $f \in D\widetilde{\mathcal{F}}_{2n,j}^r$ such that $f(x)$ decays exponentially as $|x| \rightarrow \infty$ and $f^{(j)}(\nu) = 0, \nu \in \mathbf{Z}$. Then*

$$(5.2) \quad f(x) = - \sum_{\nu \in \mathbf{Z}} a_\nu f_j(x-\nu),$$

where $(a_\nu)_{\nu \in \mathbf{Z}}$ decays exponentially as $|\nu| \rightarrow \infty$.

Proof. Since $f \in D\widetilde{\mathcal{F}}_{2n,j}^r$,

$$(5.3) \quad f(x) = \sum_{\nu \in \mathbf{Z}} c_\nu N_{2n,j}(2x-\nu),$$

where $(c_\nu)_{\nu \in \mathbf{Z}}$ decays exponentially as $|\nu| \rightarrow \infty$, and the condition $f^{(j)}(\nu) = 0, \nu \in \mathbf{Z}$, leads to

$$(5.4) \quad \sum_{\nu \in \mathbf{Z}} c_\nu N_{2n,j}^{(j)}(2k-\nu) = 0, \quad k \in \mathbf{Z}.$$

Let

$$p(z) := \sum_{\nu \in \mathbf{Z}} N_{2n,j}^{(j)}(\nu) z^\nu, \quad q(z) := \sum_{\nu \in \mathbf{Z}} c_\nu z^\nu.$$

Equation (5.4) is equivalent to

$$p(z)q(z) + p(-z)q(-z) = 0, \quad \text{for all } z.$$

Since $N_{2n,j}$ has compact support, p is a polynomial. If p has zeros $\{z_j\}$, then it does not have zeros $\{-z_j\}$, and so $\{-z_j\}$ are zeros of q . We can therefore write

$$(5.5) \quad q(z) = p(-z)r(z).$$

Then $p(z)p(-z)r(z) + p(-z)p(z)r(-z) = 0$, which gives $r(z) = -r(-z)$. Therefore, we can write

$$r(z) = \sum_{\nu \in \mathbf{Z}} a_\nu z^{2\nu-1},$$

where $(a_\nu)_{\nu \in \mathbf{Z}}$ has exponential decay as $|\nu| \rightarrow \infty$, by (5.5). Also by (5.5),

$$(5.6) \quad c_\nu = \sum_{l \in \mathbf{Z}} a_l N_{2n,j}^{(j)}(\nu-2l+1)(-1)^{\nu+1}.$$

Substituting (5.6) into (5.3) and using (5.1) leads to (5.2). \square

Proposition 5.1. *Let $g \in D\widetilde{\mathcal{F}}_{2n}^r$ such that $g(x)$ decays exponentially as $|x| \rightarrow \infty$, and $g^{(j)}(\nu) = 0, \nu \in \mathbf{Z}, j = 0, \dots, r-1$. Then*

$$(5.7) \quad g(x) = \sum_{j=0}^{r-1} \sum_{\nu \in \mathbf{Z}} a_{j\nu} f_j(x-\nu),$$

where $(a_{j\nu})_{\nu \in \mathbf{Z}}$ decays exponentially as $|\nu| \rightarrow \infty$.

Proof. Since $g \in D\widetilde{\mathcal{S}}_{2n}^r$, $g = g_0 + g_1 + \cdots + g_{r-1}$, where $g_j \in D\widetilde{\mathcal{S}}_{2n,j}^r$, $j = 0, \dots, r-1$, with $g_j^{(j)}(\nu) = 0$, $\nu \in \mathbf{Z}$. The result follows from the previous lemma. \square

For $j = 0, \dots, r-1$, let $\psi_j := f_j^{(n)}$, where f_j is given by (5.1). Then $\psi_j \in W_0$ and is supported on $[0, 2n - 2r + 1]$.

Theorem 5.1. *The set $\{T_\nu \psi_j : \nu \in \mathbf{Z}, j = 0, \dots, r-1\}$ is complete in W_0 .*

Proof. It is sufficient to show that for $k = 0, \dots, r-1$,

$$(5.8) \quad N_{n,k}(2x) = f(x) + \sum_{j=0}^{r-1} \sum_{\nu \in \mathbf{Z}} a_{j\nu} \psi_j(x - \nu),$$

where $f \in V_0 := \widetilde{\mathcal{S}}_n^r$ has exponential decay as $|x| \rightarrow \infty$ and $(a_{j\nu})_{\nu \in \mathbf{Z}}$ decays exponentially as $|\nu| \rightarrow \infty$.

Choose $M \in D\widetilde{\mathcal{S}}_{2n}^r$ with

$$(5.9) \quad M^{(n)}(x) = N_{n,k}(2x).$$

Since $N_{n,k}(2x)$ is supported on $[0, \frac{n}{2} - r + 1]$, M may be taken as a polynomial of degree $n-1$ outside $[0, \frac{n}{2} - r + 1]$. It can therefore be written as $M = F + G$, where $G \in D\widetilde{\mathcal{S}}_{2n}^r$ has compact support, and $F \in \widetilde{\mathcal{S}}_{2n}^r$ equals a polynomial of degree $n-1$ outside a compact set. Now let H be the unique function in $\widetilde{\mathcal{S}}_{2n}^r$ such that

$$H^{(j)}(\nu) = G^{(j)}(\nu), \quad j = 0, \dots, r-1, \quad \nu \in \mathbf{Z}.$$

Since G has compact support, the theory of cardinal Hermite interpolation shows that $H(x)$ decays exponentially as $|x| \rightarrow \infty$. Let $R := G - H$. Then $R \in D\widetilde{\mathcal{S}}_{2n}^r$, $R^{(j)}(\nu) = 0$, $j = 0, \dots, r-1$, $\nu \in \mathbf{Z}$, and $R(x)$ decays exponentially as $|x| \rightarrow \infty$, and we can write $M = F + H + R$. By Proposition 5.1, R can be expressed as

$$R(x) = \sum_{j=0}^{r-1} \sum_{\nu \in \mathbf{Z}} a_{j\nu} f_j(x - \nu),$$

where $(a_{j\nu})_{\nu \in \mathbf{Z}}$ decays exponentially as $|\nu| \rightarrow \infty$. Also by (4.3)

$$H(x) = \sum_{j=0}^{r-1} \sum_{\nu \in \mathbf{Z}} b_{j\nu} N_{2n,j}(x - \nu),$$

where $(b_{j\nu})_{\nu \in \mathbf{Z}}$ decays exponentially as $|\nu| \rightarrow \infty$. By (5.9)

$$\begin{aligned} N_{n,k}(2x) &= F^{(n)}(x) + H^{(n)}(x) + R^{(n)}(x) \\ &= F^{(n)}(x) + \sum_{j=0}^{r-1} \sum_{\nu \in \mathbf{Z}} b_{j\nu} N_{2n,j}^{(n)}(x - \nu) + \sum_{j=0}^{r-1} \sum_{\nu \in \mathbf{Z}} a_{j\nu} \psi_j(x - \nu). \end{aligned}$$

Since $F^{(n)} \in V_0$ has compact support, and $H^{(n)} \in V_0$ has exponential decay, it follows that $F^{(n)} + H^{(n)} \in V_0$ has exponential decay. \square

REFERENCES

1. G. Battle, *A block spin construction of ondelettes, Part I: Lemarié functions*, Comm. Math. Phys. **110** (1987), 601–615.
2. G. Battle and P. Federbush, *Ondelettes and phase cell cluster expansion; a vindication*, Comm. Math. Phys. **109** (1987), 417–419.
3. Charles K. Chui and Jian-zhong Wang, *A cardinal spline approach to wavelets*, Proc. Amer. Math. Soc. **113** (1991), 785–793.
4. —, *A general framework of compactly supported splines and wavelets*, J. Approx. Theory **71** (1992).
5. I. Daubechies, *Orthonormal bases of compactly supported wavelet*, Comm. Pure Appl. Math. **41** (1988), 909–996.
6. P. R. Halmos, *A Hilbert space problem book*, 2nd ed., Springer-Verlag, 1982.
7. S. Jaffard and Y. Meyer, *Bases d'ondelettes dans des ouverts de \mathbb{R}^n* , J. Math. Pures Appl. **68** (1989), 95–108.
8. S. L. Lee, *B-splines for cardinal Hermite interpolation*, Linear Algebra and Appl. **12** (1975), 269–280.
9. —, *Fourier transforms of B-splines and fundamental splines for cardinal Hermite interpolations*, Proc. Amer. Math. Soc. **57** (1976), 291–296.
10. S. L. Lee and A. Sharma, *Cardinal lacunary interpolation by g-splines. I. The characteristic polynomials*, J. Approximation Theory **16** (1976), 85–96.
11. P. G. Lemarie, *Ondelettes a localisation exponentielles*, J. Math. Pures Appl. **67** (1988), 227–236.
12. S. Mallat, *Multiresolution approximations and wavelet orthonormal bases of $L^2(\mathbb{R})$* , Trans. Amer. Math. Soc. **315** (1989), 69–87.
13. —, *A theory for multiresolution signal decomposition: the wavelet representation*, IEEE Trans. on Pattern Analysis and Machine Intelligence **2** (1989), 674–693.
14. —, *Multifrequency channel decomposition of images and wavelet models*, IEEE Trans. ASSP **37** (1989), 2091–2110.
15. Y. Meyer, *Ondelettes et fonctions splines*, Seminaire Equations aux Derivees Partielles, École Polytechnique, Paris, 1986.
16. —, *Principe d'incertitude, bases hilbertiennes et algèbres d'opérateurs*, Bourbaki seminar, #662, 1985–1986.
17. J. B. Robertson, *On wandering subspaces for unitary operators*, Proc. Amer. Math. Soc. **16** (1965), 233–236.
18. I. J. Schoenberg, *Cardinal spline interpolation*, CBMS-NSF Series in Appl. Math., no. 12, SIAM, Philadelphia, Pa., 1973.

DEPARTMENT OF MATHEMATICS, TEXAS A & M UNIVERSITY, COLLEGE STATION, TEXAS 77843

Current address: Department of Mathematics and Computer Science, University of Dundee, Dundee, DD1 4HN, Scotland

E-mail address: tgoodman@mcs.dundee.ac.uk

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, 10 KENT RIDGE CRES-CENT, SINGAPORE 0511

E-mail address, S. L. Lee: matleesl@nusvm.bitnet

E-mail address, W. S. Tang: mattws@nusvm.bitnet