THE PLANAR CLOSING LEMMA FOR CHAIN RECURRENCE

MARIA LUCIA ALVARENGA PEIXOTO AND CHARLES CHAPMAN PUGH

ABSTRACT. If z is a chain recurrent point of a C^r planar flow, all of whose fixed points are hyperbolic, then it is proved that the orbit through z becomes periodic under a perturbation that is C^r small in the Whitney topology.

1. INTRODUCTION

Let φ be a flow on M , a metric space. The forward orbit of $p \in M$ under φ is

$$O_+(p) = \{\varphi_t p : t \geq 0\}.$$

There are several natural ways to extend O_+ , φ -invariantly. The simplest is to form

$$\overline{O_+}(p) = O_+(p) \cup \omega(p)$$

where $\omega(p)$ is the ω -limit set of p ,

$$\omega(p) = \{x \in M : \varphi_{t_n}(p) \rightarrow x \text{ for some } t_n \rightarrow \infty\}.$$

Thus, $\overline{O_+}(p)$ is the closure of $O_+(p)$. Although $\omega(p)$ is closed and φ -invariant, there may be points q near it which tend to new limits. This leads one to define the *first prolongation* of $O_+(p)$ as

$$P_1(p) = O_+(p) \cup J_1(p),$$

$$J_1(p) = \{y \in M : \varphi_{t_n}(x_n) \rightarrow y \text{ for some } t_n \rightarrow \infty \text{ and } x_n \rightarrow x\}.$$

Repeating the process transfinitely, with neighborhood closure at limit ordinals, leads to the ultimate prolongation of $O_+(p)$:

$$P(p) = O_+(p) \cup \bigcup_{\alpha} J_{\alpha}(p) = \bigcup_{\alpha}^{\uparrow} P_{\alpha}(p)$$

where α ranges through all ordinals. See Bhatia and Szegő (1970) for details. Correspondingly we write $J(p) = \bigcup_{\alpha} J_{\alpha}(p)$. If M is separable then it turns out that some countable ordinal $\beta = \beta(\varphi)$ suffices, that is

$$P_{\alpha}(p) = P_{\beta}(p) \quad \text{for all } \alpha \geq \beta.$$

Surprisingly, a more general extension $O_+(p)$ exists based on the concept of (ε, T) -chains.

Received by the editors March 4, 1991 and, in revised form, May 29, 1991.

1991 *Mathematics Subject Classification*. Primary 58Fxx; Secondary 58F10, 58F30, 34C28, 34D10, 34D30.

Definitions. An *orbit segment* of φ is a set $\{\varphi_t x : 0 \leq t \leq \tau\}$ for some $\tau \geq 0$. We denote it by $[x, y]$ where $y = \varphi_\tau(x)$. Its *duration* is τ . An (ε, T) -*chain* from p to q is a finite collection Γ of orbit segments $[x_1, y_1], \dots, [x_n, y_n]$ each of duration $\geq T$ such that $d(y_i, x_{i+1}) \leq \varepsilon$, $0 \leq i \leq n$, where we understand $p = y_0$ and $q = x_{n+1}$. See Figure 1.1.

FIGURE 1.1. An (ε, T) -chain from p to q .

The *chain extension* of $O_+(p)$ is $O_+(p) \cup E_+(p)$ where $E_+(p) = \{q \in M : \text{for all } \varepsilon, T > 0, \text{ there exists an } (\varepsilon, T)\text{-chain from } p \text{ to } q\}$.

If M is not compact then the ε and T above should be thought of as positive continuous functions defined on M . They are “local constants”. An (ε, T) -chain is a finite collection of orbit segments, as before, satisfying

- (a) $[x_i, y_i]$ has duration $\geq T(x_i)$, $1 \leq i \leq n$;
- (b) $d(y_i, x_{i+1}) < \varepsilon(y_i)$, $0 \leq i \leq n$.

Note. Nitecki (1978) requires T to be constant, even when M is noncompact. In Appendix 1, we show that this does not affect the definition of chain extension.

Corresponding to concepts of orbit extension are concepts of recurrence. The point $p \in M$ is

fixed	if $\{p\} = O_+(p)$,
periodic	if $p \in O_+(p) \neq \{p\}$,
ω -recurrent	if $p \in \omega(p)$,
nonwandering	if $p \in J_1(p)$,
prolongationally recurrent	if $p \in \bigcup_\alpha J_\alpha(p)$,
chain recurrent	if $p \in E_+(p)$.

How much do these recurrences coincide? The Closing Lemma may be viewed as an attempt to force coincidence by approximation: ω -recurrent

points and J_1 recurrent points become periodic after a perturbation of the dynamics. See Pugh (1967a, and b) respectively. The Closing Lemma for more general recurrence has not met with success, although in Pugh (1984), an example is given to show that the J_2 Closing Lemma is false on noncompact surfaces, and in Pugh (1975) there is an example against C^2 -double closing of J_2 -recurrence on the 2-torus. Here we verify the Closing Lemma for some planar chain recurrence.

1.1. Theorem. *Suppose X is a C^r vectorfield on an open subset M of S^2 , $1 \leq r \leq \infty$, X generates a flow ϕ , and $z \in M$ is chain recurrent. Suppose all the fixed points of ϕ are hyperbolic. If $\mathcal{U}$ is a neighborhood of X in the C^r Whitney topology then $\mathcal{U}$ contains a field Y for which z is periodic.*

See §5 for the proof of this theorem.

Remarks. S^2 minus a point is R^2 so the theorem applies to flows on R^2 . The assumption about hyperbolic fixed points reflects our inability to solve the following.

Local singular connecting problem. Let p be an isolated fixed point of the smooth flow ϕ on R^2 . Suppose $\omega(x) = p = \alpha(y)$ for some points $x, y \neq p$.

Can ϕ be C^r perturbed to put x, y on the same trajectory?

FIGURE 1.2. Connecting x to y via p in the hyperbolic case; Y approximates $X = \phi$.

FIGURE 1.3. Connecting x to y via p in a nonhyperbolic case; Y approximates $X = \phi$.

If p is hyperbolic then the answer is “yes”; see Figure 1.2. In some other cases (Figure 1.3) the answer is also “yes”, but in general the problem may be hard.

2. THE WHITNEY TOPOLOGY

If M is a compact manifold, it is natural to topologize the space of C^r tangent vector fields on M , $\mathcal{X}^r(M)$, with the C^r -uniform topology. When M is no longer compact, such as $M = \mathbf{R}^2$, one uses the Whitney topology. It can be defined in several equivalent ways, one of which is the following. Note that the definition we give is simpler than the usual one which involves coordinate coverings.

First take the case that M is Euclidean space, $M = \mathbf{R}^n$. If ε is a positive continuous function on $\mathbf{R}^n$ and $X \in \mathcal{X}^r(\mathbf{R}^n)$ then we set

$$\mathcal{N}_\varepsilon(X) = \{Y \in \mathcal{X}^r(\mathbf{R}^n) : \text{for all } x \in \mathbf{R}^n, \\ |Y(x) - X(x)| + \cdots + \|D^r(Y - X)_x\| < \varepsilon(x)\}.$$

These $\mathcal{N}_\varepsilon(X)$ are a neighborhood base at X for the C^r Whitney topology on $\mathcal{X}^r(\mathbf{R}^n)$.

Next take the case that M is a general smooth m -manifold. The meaning of $\|D^r(Y - X)\|$ becomes a problem which we bypass by choosing

- (a) a proper smooth embedding $M^m \hookrightarrow \mathbf{R}^n$,
- (b) the proper smooth tubular neighborhood retraction $\rho: N \rightarrow M$ along short normal segments to M in $\mathbf{R}^n$,
- (c) a smooth bump function $\beta: N \rightarrow [0, 1]$ which is identically zero near ∂N and identically one near M .

Then we define the extension operator $e: \mathcal{X}^r(M) \rightarrow \mathcal{X}^r(\mathbf{R}^n)$ by

$$e(X) = \begin{cases} \beta(x)\rho^*X(x), & x \in N, \\ 0, & x \notin N, \end{cases}$$

where ρ^*X is the parallel extension of X to N along ρ -fibers. That is, $\rho^*X(x) \in T_x(\mathbf{R}^n)$ is the $\mathbf{R}^n$ -parallel-translate of $X(\rho(x))$.

The Whitney topology on $\mathcal{X}^r(M)$ is the e -pullback of that on $\mathcal{X}^r(\mathbf{R}^n)$. That is, a neighborhood base at $X \in \mathcal{X}^r(M)$ is given by $\{Y \in \mathcal{X}^r(M) : e(Y) \in \mathcal{N}_\varepsilon(eX)\}$ where ε runs through all positive continuous functions on $\mathbf{R}^n$. It is independent of the choices made in (a), (b), (c).

2.1. Convexity Proposition. *$\mathcal{X}^r(M)$ has a neighborhood base $\mathcal{V}$ with the following property. If V is a $\mathcal{V}$ neighborhood of X , if $\Delta_1, \Delta_2, \dots, \Delta_n$ in $\mathcal{X}^r(M)$ have disjoint supports, and if $Y_i = X + \Delta_i$ lies in V for each i then $Y = X + \Sigma \Delta_i$ also lies in V .*

Proof. Suppose $M = \mathbf{R}^n$. Since $\mathcal{N}_\varepsilon(X)$ is defined pointwise, it is clear that it has the above property. If $M \hookrightarrow \mathbf{R}^n$ then $\mathcal{X}^r(M)$ inherits this property from $\mathcal{X}^r(\mathbf{R}^n)$ under e -pullback. $\square$

3. FLOWBOXES AND LIFT

To quantify the perturbations we use for the Closing Lemma, we introduce flowboxes in a uniform way.

Let φ be a C^r flow on a manifold M^m . We assume M equipped with some fixed smooth Riemann structure. As standing notation we call M_0 the set of regular points of φ ,

$$M_0 = \{x \in M : x \text{ is not a fixed point of } \varphi\}.$$

Choose and fix a smooth $(m-1)$ plane bundle over M_0 complementary to X ,

$$\Pi \oplus \text{span}(X) = TM_0.$$

For any $x \in M_0$, there exists $\nu > 0$ such that the map

$$f_x^\nu : [-\nu, \nu] \times \Pi_x(\nu) \rightarrow M, \quad (t, v) \rightarrow \varphi_t(\exp_x(v))$$

is a C^r embedding, where $\Pi_x(\nu)$ is the disc of radius ν in Π_x . Since M is locally compact we may determine some positive continuous function ν on M_0 such that $f_x^{\nu(x)}$ is an embedding for each $x \in M_0$. Fix such a flowbox-radius function ν .

Definition. The *canonical flowbox* of radius $\nu(x)$ at $x \in M_0$ is the range F_x^ν of the chart map $f_x^{\nu(x)}$. If $0 < c \leq 1$ then the *c-central portion* of F_x^ν is the set

$$\{\varphi_t(\exp_x(v)) : |t| \leq \nu(x) \text{ and } v \in \Pi_x(c\nu(x))\}.$$

3.1. Lift Lemma. Let $\mathcal{U}$ be a neighborhood of $X = \dot{\varphi}$ in the C^r Whitney topology on $\mathcal{X}^r(M)$, $r \geq 1$, and let $\mu : M_0 \rightarrow \mathbb{R}$ be a continuous function such that $0 < \mu(x) < \nu(x)$ for all $x \in M_0$. Then there exists a positive continuous function c on M_0 such that if y, y' lie at opposite ends of the $c(x)$ -central portion of the canonical F_x^ν then there exists a field $Y \in \mathcal{U}$ with $Y \equiv X$ off F_x^μ and, under the Y -flow, y flows to x which flows to y' across the $c(x)$ -central portion of F_x^ν .

Proof. Consider Figure 3.1. $\square$

FIGURE 3.1. Y lifts y to x and x to y' .

4. QUASIORBITS

A C^1 curve $\lambda(t)$ in M is an ε -quasiorbit of X if $|\lambda'(t) - X(\lambda(t))| < \varepsilon$ for all t where λ is defined. In other words, it is an ε -approximate solution

of the differential equation $x' = X(x)$. When M is noncompact it is natural to replace the constant ε with a positive continuous function $\varepsilon(x)$. Then we require $|\lambda'(t) - X(\lambda(t))| < \varepsilon(\lambda(t))$.

In Nitecki (1978) it is shown that a point z is chain recurrent if and only if for every positive continuous function $\varepsilon > 0$ on M there exists a periodic ε -quasiorbit through z .

4.1. Global Replacement Lemma. *Suppose that the fixed points of the X -flow on M^2 are hyperbolic, therefore countable in number, say $p_1, p_2, \dots$ and z is a chain recurrent point, $z \in M_0$. Then there is a neighborhood V of $\{p_i\}$ such that if $\varepsilon: M \rightarrow [0, \infty)$ is a given continuous function, positive on $M \setminus \bar{V}$, then there is a periodic ε -quasiorbit λ through z . Moreover λ is a simple closed curve.*

Remark. We permit $\varepsilon \equiv 0$ on $\bar{V}$. Then $\lambda \cap \bar{V}$ consists of true X -trajectory arcs.

4.2. Local Replacement Lemma. *Suppose p is a hyperbolic fixed point for a flow ϕ in $\mathbb{R}^2$ and U is a neighborhood of p on which ϕ is linearizable. Let $N \subset U$ be a small neighborhood of p and let $T_\pm, T_\pm^*$ be transversals to $W^s(p), W^u(p)$ as shown in Figure 4.1. Then, for every $\varepsilon > 0$ there exists $\delta > 0$ such that any δ -quasiorbit σ from T to T^* in U can be replaced by an ε -quasiorbit λ from T to T^* in U such that*

- (a) $\lambda = \sigma$ near T and T^* ,
- (b) $\lambda \cap N$ is a true orbit arc,
- (c) λ is an arc.

FIGURE 4.1

Remark. This result is valid in dimension ≤ 3 but not in dimension ≥ 4 . It is easy to see that (c) is implied by (a) and (b).

Proof. We draw flowboxes $F_{\pm}, F_{\pm}^*$ across $W^u(p), W^s(p)$ to get the situation shown in Figure 4.2.

FIGURE 4.2

FIGURE 4.3

Let σ be a δ -quasiorbit from $z \in T$ to $z^* \in T^*$ in U . When δ is small, the parts of σ outside N are nearly trajectory arcs in the C^1 sense. Near p , σ may be quite twisted, because the angle between X and σ' may be large when X and $|X - \sigma'|$ are small.

Suppose $z \in T_+$, $z^* \in T_+^*$. According to Appendix 2, we may replace σ with a φ -arc λ_0 from F_+ to F_+^* (necessarily in the first quadrant) together with δ -quasiorbits across F_+ , F_+^* as shown. See Figure 4.3(a). Note that z , z^* , σ need not lie in the first quadrant. See Figure 4.3(b).

The heavy line is the φ -arc λ_0 from F_+ to F_+^* . The dotted line in F_+ is an ε -quasiorbit λ_+ joining λ_0 to the incoming part of σ , while λ_+^* is a similar ε -quasiorbit in F_+^* .

Consider $\sigma \cap F_+$. It is either in the first quadrant or nearly so, because otherwise σ could not reach T_+^* .

Thus, for small δ , an ε -quasiorbit λ_+ exists as asserted; similarly λ_+^* .

If $z \in T_-$ and/or $z^* \in T_-^*$ then the corresponding construction places λ_0 in one of the other quadrants, verifying (a), (b), (c). $\square$

Proof of Proposition 4.1. Let $\{p_i\}$ be the set of fixed points of the X -flow and $z_0 \in M_0$ the given chain recurrent point. Choose disjoint neighborhoods U_i of p_i on which the X -flow can be linearized, $z \notin U_i$. In each U_i draw the transversals $T_\pm(i)$, $T_\pm^*(i)$ and the smaller neighborhood N_i of p_i as in Lemma 4.2. Choose a yet smaller neighborhood V_i of p_i with $\overline{V_i} \subset N_i$. Call $V = \bigcup_i V_i$. Let ε be a given nonnegative continuous function on M , positive on $M \setminus \overline{V}$. Since $\overline{V_i} \subset N_i$, $\varepsilon(x)$ is bounded away from 0 for $x \in U_i \setminus N_i$; say $\varepsilon_i = \inf\{\varepsilon(x) : x \in U_i \setminus N_i\} > 0$. Let δ_i be the number supplied by Lemma 4.2 and choose a continuous function $\delta(x)$ such that

$$0 < \delta(x) \leq \varepsilon(x) \quad \text{for all } x \in M \setminus U,$$

$$\delta(x) \leq \delta_i \quad \text{if } x \in U_i, \quad U = \bigcup_i U_i.$$

According to Nitecki's Lemma there is a periodic δ -quasiorbit σ through the chain recurrent point z . By Lemma 4.2 it can be replaced by an ε -quasiorbit λ whose intersection with N consists of true X -trajectory arcs. By a slight change we assume these X -trajectory arcs are distinct and λ is an immersed curve with generic self intersection. The self intersection points are finite in number and occur outside N . Cancelling extraneous loops and rounding off corners in the standard way, λ becomes a smooth simple ε -quasiorbit through z as asserted. $\square$

5. TWO-DIMENSIONAL CLOSING

Before starting the proof of the Closing Lemma, we need two more preliminary results. The first refers to the flowbox radius function $\nu: M_0 \rightarrow (0, \infty)$, M_0 being the set of nonfixed points of the X -flow. It was constructed in §3.

5.1. Lemma. *There exist positive continuous functions $\varepsilon_0, \omega, \mu: M_0 \rightarrow (0, 1)$ such that $0 < \mu < \omega < \nu$ and, writing F_x^ω for $F_x^{\omega(x)}$, etc.,*

- (a) *if $F_{x'}^\omega \cap F_x^\omega \neq \emptyset$ then $F_{x'}^\omega \subset F_x^\nu$;*
- (b) *if $x' \notin F_x^\omega$ then $F_{x'}^\mu \cap F_x^\mu = \emptyset$;*
- (c) *suppose $F_{x_1}^\omega, F_{x_2}^\omega$ are contained in F_x^ν and $F_{x_3}^\mu$ meets the convex hull of $F_{x_1}^\omega \cup F_{x_2}^\omega$ in F_x^ν . Then any ε_0 -quasiorbit σ through a point of $F_{x_3}^\mu$ stretches all the way across F_x^ν .*

Proof. If $\omega \ll \nu$ then intersections of ω -sized boxes occur only near the centers of ν -sized boxes, giving (a). If $\mu \ll \omega$ and $x' \notin F_x^\omega$ then $F_{x'}^\mu$ is so small that it cannot meet F_x^μ , giving (b). Since F_x^ν is compact, ω is bounded away from zero on it, and the centers x_1, x_2 of ω -sized boxes are bounded away from ∂F_x^ν . If $\mu \ll \omega$ then their convex hull H and any $F_{x_3}^\mu$ meeting H are also bounded away from ∂F_x^ν . For small enough ε_0 this gives (c). See Figure 5.1. Note that (c) applies to all $F_{x_3}^\mu$ which meet F_x^ω . $\square$

FIGURE 5.1

5.2. Lemma. *A simple loop in the plane or sphere which crosses an arc in one direction only, crosses the arc at most once.*

Proof. Consider the Poincaré-Bendixson picture in Figure 5.2. $\square$

FIGURE 5.2

Remark. In the proof of the Closing Lemma given below, we perturb X in a sequence of flowboxes along a periodic ε -quasiorbit through z . This resembles the box-chain construction in Peixoto (1988). The point to keep in mind is “does the perturbation act *only one time* on the periodic orbit being produced?”

Proof of the Closing Lemma (Theorem 1.1). Recall that we are given a C^r tangent vector field X on M , M is an open subset of S^2 , $\mathcal{U}$ is a given neighborhood of X in the C^r Whitney topology, $r \geq 1$, and z is a nonfixed chain recurrent point of the X -flow. The fixed points of X are assumed to be hyperbolic. We want to find $Y \in \mathcal{U}$ with a periodic orbit through z .

Step 1. By the construction in §3 we determine the flowbox radius function ν . It is positive and continuous on M_0 , the set of nonfixed points.

Step 2. By Lemma 5.1 we determine positive continuous functions $\omega, \mu, \varepsilon_0$ on M_0 .

Step 3. From hyperbolicity of the fixed points we use Proposition 4.1 to determine a suitably small neighborhood V of the fixed point set.

Step 4. From $\mathcal{U}, \mu, \omega$ we use the Lift Lemma 3.1 to determine a small enough $c: M_0 \rightarrow (0, 1]$. Besides, we require c to be so small that on the c -central portion of any F_x^ν, cF_x^ν , with $x \in M_0$, the following property holds:

(d) If $x' \in cF_x^\nu$ and $F_{x'}^\mu \subset F_x^\nu$ then cF_x^ν cuts across the middle of $F_{x'}^\mu$. See Figure 5.3.

FIGURE 5.3. The c -central portion of F_x^ν cuts across the middle of $F_{x'}^\mu$.

It follows from (d) that

(e) If $F_{x_1}^\mu, \dots, F_{x_k}^\mu$ are disjoint, contained in F_x^ν and their centers $x_1, \dots, x_k$ lie in cF_x^ν , then $cF_x^\nu \setminus \bigcup F_{x_i}^\mu$ consists of $k + 1$ topological rectangles as shown shaded in Figure 5.4.

FIGURE 5.4

Note that the actual shape of the $F_{x_i}^\mu$ can be somewhat contorted, as in Figure 5.5, but the monotonicity of the shaded rectangles persists.

FIGURE 5.5. Possible shapes of $F_{x_i}^\mu$.

Step 5. Choose $\varepsilon: M \rightarrow [0, 1]$ as follows. We require that ε is continuous, identically zero on $\overline{V}$, positive off $\overline{V}$, and $\varepsilon \leq \varepsilon_0$ everywhere. Also we require ε to be so small that

(f) If σ is any ε -quasiorbit through $x \in M_0$, then the connected component of $\sigma \cap F_x^\omega$ containing x is an arc in the c -central portion of F_x^ω .

It is possible to verify (f) because when x ranges over $M \setminus \overline{V}$, $|\sigma' - X|$ being small implies that the angle between σ' and X is small, while on $\overline{V}$, $\varepsilon \equiv 0$. See Figure 5.6.

FIGURE 5.6

Note that the c -central portion of F_x^ω is contained in the c -central portion of F_x^ν .

Step 6. Using this choice of ε and Proposition 4.1, we determine a smooth simple closed curve λ which is a periodic ε -quasiorbit through z . In V , λ

consists of true X -trajectory arcs. By Lemma 5.2,

(g) If $x \in \lambda$ then $F_x^\mu, F_x^\omega, F_x^\nu$ meet λ in nested arcs.

See Figure 5.7. (It is exactly here we use planarity.)

FIGURE 5.7

Step 7. Starting at $z = z_0$, we cover λ with ω -sized flowboxes as follows. By (f), we know that λ exits the right-hand edge of $F_{z_0}^\omega$ in the c -central portion, say at u_0 . Choose $z_1 \in \lambda$ slightly beyond u_0 so that $\lambda \cap F_{z_1}^\omega$ includes u_0 in its interior. Repeat the construction beyond the point u_1 where $\lambda \cap F_{z_1}^\omega$ exits $F_{z_1}^\omega$. Since λ is compact, contained in M_0 and ω is continuous, ω is bounded away from zero on λ and we eventually get a finite, indexed cover $F_{z_0}^\omega, \dots, F_{z_{n-1}}^\omega$ of λ . We set $z_n = z_0$ and write indices cyclically. We call z_i , *center points*. See Figure 5.8.

FIGURE 5.8

By (g), choice of $\varepsilon \leq \varepsilon_0$, and part (c) of Lemma 5.1, we know that $\lambda_0 = \lambda \cap F_{z_0}^\nu$ is an arc crossing all of $F_{z_0}^\nu$. By (f), λ_0 lies in the c -central portion of $F_{z_0}^\nu$. By part (a) of Lemma 5.1, $F_{z_{n-1}}^\omega, F_{z_0}^\omega, F_{z_1}^\omega$ are all contained in $F_{z_0}^\nu$. Because each z_i lies outside $F_{z_0}^\omega, \dots, F_{z_{i-1}}^\omega$, part (b) of Lemma 5.1 implies that $F_{z_i}^\mu$ is disjoint from $F_{z_0}^\mu, \dots, F_{z_{i-1}}^\mu$. Thus

(h) The boxes $F_{z_i}^\mu$ are disjoint, $i = 0, \dots, n-1$.

Step 8. We need to say a little more about the configuration of these boxes. Consider $F_{z_i}^\omega, F_{z_{i+1}}^\omega$. They lie in $F_{z_i}^\nu$ by construction. The part of $cF_{z_i}^\nu$ between $F_{z_i}^\mu$ and $F_{z_{i+1}}^\mu$ is a rectangle R_i , by (d). (If $i \neq n-1$, R_i is contained in $F_{z_i}^\omega$,

but for $i = n - 1$, all we can say is that R_i is part of the convex hull of $F_{z_i}^\mu \cup F_{z_{i+1}}^\mu$.) Suppose $F_{z_j}^\mu$ meets R_i . By (c) $\lambda \cap F_{z_j}^\mu$ is part of an arc stretching all the way across $F_{z_i}^\nu$. But λ meets $F_{z_i}^\nu$ in a unique arc, namely the one containing z_i . Therefore z_j lies on this arc. Since $F_{z_j}^\mu$ meets R_i , we see by (e) that z_j lies between z_i and z_{i+1} on λ , a contradiction. Therefore,

(i) the rectangle R_i which is the part of $cF_{z_i}^\nu$ between $F_{z_i}^\mu$ and $F_{z_{i+1}}^\mu$ meets no $F_{z_j}^\mu$. See Figure 5.9.

FIGURE 5.9

Step 9. Consider the point v_i where λ enters $F_{z_i}^\mu$. Let y_i be the point of the left edge of $F_{z_i}^\nu$ on the X trajectory through v_i . Thus, $[y_i, v_i]$ is a horizontal line in the c -central portion of $F_{z_i}^\nu$. Similarly, let y'_i be the point of the right edge of $F_{z_i}^\nu$, horizontally related to v_{i+1} . See Figure 5.10.

FIGURE 5.10

By the Lift Lemma 3.1, there is a perturbation of X , Y_i , such that $Y_i \equiv X$ off $F_{z_i}^\mu$ and under the Y_i -flow we have

$$y_i \rightarrow v_i \rightarrow z_i \rightarrow v_{i+1} \rightarrow y'_i.$$

For y_i, y'_i lie in the c -central portion of $F_{z_i}^\nu$. Let γ_i be the part of this Y_i -trajectory from v_i to v_{i+1} . It lies in the c -central portion of $F_{z_i}^\nu$ and therefore

(j) $\gamma_i \subset (F_{z_i}^\mu \cup R_i)$. See Figure 5.11.

FIGURE 5.11

Step 10. Let $Y = X + \sum_{i=0}^{n-1} (Y_i - X)$. By the convexity property of the Whitney topology mentioned in Proposition 2.1 and disjointness of the supports of the $Y_i - X$ shown in (h), $Y \in \mathcal{U}$. We claim that $\gamma_0 \cup \gamma_1 \cup \dots \cup \gamma_{n-1}$ is a periodic orbit of Y . By (i) none of the Y_j affects the part of γ_i in R_i and, by disjointness of the supports $F_{z_i}^\mu$ of $Y_i - X$, none of the $F_{z_j}^\mu$, $j \neq i$, affects the part of γ_i in $F_{z_i}^\mu$. Therefore γ is indeed Y -periodic. $\square$

APPENDIX 1

Above and in Nitecki (1978) one uses (ε, T) -chains on noncompact manifolds with ε a function but T a constant. Here we show that the natural definition in which T also is a function leads to the same chain extensions and chain limit sets.

Let φ be a flow on the metric space M . An *orbit segment* of φ is a set $\{\varphi_t(x); 0 \leq t \leq \tau\}$ which we denote by $[x, y]$, $y = \varphi_\tau(x)$. An (ε, T) -*chain from p to q* is a finite union of orbit segments

$$\Gamma = \bigcup_{i=1}^n \gamma_i, \quad \gamma_i = [x_i, y_i], \quad y_i = \varphi_{t_i}(x_i),$$

such that

- (a) $t_i \geq T(x_i)$, $1 \leq i \leq n$,
- (b) $d(y_i, x_{i+1}) < \varepsilon(y_i)$, $0 \leq i \leq n$.

In (b) we understand $p = y_0$ and $q = x_{n+1}$. See Figure 1. The *duration* of γ_i is t_i and the *duration* of Γ is $\sum t_i$.

FIGURE 1. An (ε, T) -chain from p to q .

Theorem. Suppose M is locally compact, $p, q \in M$, and for every constant $D > 0$ and every $\delta > 0$ on M , there is $(\delta, 1)$ -chain from p to q of duration $\geq D$. Then there exists an (ε, T) -chain from p to q .

Corollary. Nitecki's definition of chain extension and chain recurrence agree with ours.

Proof. Immediate. $\square$

Proof of the theorem. Without loss of generality we assume $\varepsilon \leq 1$ and $T \geq 2$ on M . Also, we rescale the metric so all unit balls on M are compact. Then we determine an appropriately small $\delta > 0$ on M as follows.

First, for each $y \in M$, we call

$$\overline{T}(y) = 3 \max\{T(z) : d(z, \varphi_t y) \leq 1 \text{ for some } t \in [0, 2T(y)]\}.$$

In particular $\overline{T} \geq 3T$. If $\delta > 0$ is small and Γ is a $(\delta, 1)$ -chain from y to y' of duration $\tau \leq \overline{T}(y)$ then

$$(*) \quad d(\varphi_\tau(y), y') \leq \varepsilon(\varphi_\tau y).$$

Such a δ exists. For the set of such Γ 's is compact and $y' \rightarrow \varphi_\tau(y)$ as $\delta \rightarrow 0$. The convergence is uniform over all Γ and all $\tau \in [0, \overline{T}(y)]$. That $\delta(y)$ can be chosen as a continuous positive function of $y \in M$ follows from continuity of φ and local compactness of M .

By hypothesis, there is a $(\delta, 1)$ -chain Γ from p to q of duration $\geq \overline{T}(p)$. Break its orbit segments $\gamma_1, \dots, \gamma_n$ into pieces, each of duration between 1 and 2. This introduces new jumps, but all of size zero. Thus, we assume, without loss of generality, that the orbit segments $\gamma_i = [x_i, y_i]$ of Γ already have duration $t_i \in [1, 2]$.

Successively choose $n_1 < n_2 < \dots$ such that

$$\begin{aligned} T(p) &\leq T_1 = t_1 + \dots + t_{n_1} \leq 2T(p), \\ T(y_{n_1}) &\leq T_2 = t_{n_1+1} + \dots + t_{n_2} \leq 2T(y_{n_1}), \end{aligned}$$

etc. Thus, the total duration of the first n_1 orbit segments is $T_1 \doteq T(p)$, the total duration of the next $n_2 - n_1$ is $T_2 \doteq T(y_{n_1})$, and so on. Since $T \geq 2$

and $t_i \in [1, 2]$ it is always possible to choose $n_1 < n_2 < \dots < n_K \leq n$ with these properties, $K \geq 1$ being the last choice. The final $n - n_K$ orbit segments have total duration $T_{K+1} < T(y_{n_K})$ and so we combine them with the preceding $n_K - n_{K-1}$ durations as

$$\tau_K = \sum_{n_{K-1} < i \leq n} t_i = T_K + T_{K+1}.$$

When $k = 1, \dots, K-1$ we set $\tau_k = T_k$ and

$$\begin{aligned} \lambda_1 &= \{\varphi_t(p) : 0 \leq t \leq \tau_1\} = [p_1, z_1], \\ &\vdots \\ \lambda_k &= \{\varphi_t(y_{n_{k-1}}) : 0 \leq t \leq \tau_k\} = [p_k, z_k], \\ &\vdots \\ \lambda_K &= \{\varphi_t(y_{n_{K-1}}) : 0 \leq t \leq \tau_K\} = [p_K, z_K]. \end{aligned}$$

The chain $\Lambda = \lambda_1 \cup \dots \cup \lambda_K$ starts at $p_1 = p$ and ends at z_K . The duration τ_k of λ_k satisfies

$$T(p_k) \leq \tau_k \leq 2T(p_k) < \bar{T}(p_k), \quad 1 \leq k \leq K-1,$$

by construction. For $k = K$ we have $p_K = y_{n_{K-1}}$ and

$$\begin{aligned} T(p_K) &= T(y_{n_{K-1}}) \leq T_K < T_K + T_{K+1} \\ &\leq 2T(y_{n_{K-1}}) + T(y_{n_K}) \\ &\leq \frac{2}{3}\bar{T}(y_{n_{K-1}}) + \frac{1}{3}\bar{T}(y_{n_{K-1}}) = \bar{T}(p_K). \end{aligned}$$

Thus, by (*)

$$d(z_k, p_{k+1}) \leq \varepsilon(p_k), \quad 0 \leq k \leq K,$$

where z_0 and p_{K+1} denote p and q . That is, Λ is an (ε, T) -chain from p to q and the theorem is proved. $\square$

APPENDIX 2

It is not hard to see that the property of hyperbolicity of fixed points we used was:

(H) p has a neighborhood N such that every ε -quasi-orbit in N can be δ -shadowed by a single φ -orbit-segment and $\delta \rightarrow 0$ uniformly as $\varepsilon \rightarrow 0$.

For conciseness we say p is *singly δ -shadowable*. It is easy to see that hyperbolic fixed points have property H. So does the fixed point of

$$(*) \quad \dot{x} = x^2 + y^2, \quad \dot{y} = 0.$$

See Figure 1.

FIGURE 1

Proposition. Let φ be a flow in the plane with an isolated fixed point p . Property H holds for p if and only if either

- (a) φ is topologically conjugate to $(*)$ near p .
- (b) φ is topologically conjugate to a hyperbolic flow near p .
- (c) p is a rotation point of φ with alternate inward-outward nonperiodic bands, no bands of closed orbits, and no semistable closed orbits.

Proof. In Hartman (1964) the local phase portrait of isolated planar fixed points is well described in terms of sectors.

First suppose p is not a rotation point. Assume there exists an elliptic solution E . Inside E is a second elliptic solution E' . See Figure 2(a).

FIGURE 2

The ε -quasiorbit along E followed by E' is not singly δ -shadowable. Thus, there are no elliptic sectors at p . All the sectors are hyperbolic or parabolic, and none of them has any elliptic parts. They are pure.

If p has a hyperbolic sector H and also a parabolic sector P , say positive, then an ε -quasiorbit along an interior base-solution in P joined to the negative base solution of H by a C^1 curve cannot be singly δ -shadowed. See Figure 2(b).

Therefore, if p is not a rotation point then all its sectors are pure parabolic or all are pure hyperbolic. In the former case, p is a source or sink. In the latter case the number n_h of hyperbolic sectors is even because of the Poincaré Index Formula (Hartman (1964)). If $n_h = 2$ then φ is topologically equivalent to $(*)$. If $n_h = 4$, it is a hyperbolic saddle. If $n_h \geq 6$ then an ε -quasiorbit along a positive base-solution (separatrix) joined to a nonadjacent negative base solution by a C^1 curve is not singly δ -shadowable. Consequently, p obeys (a), (b), or is a rotation point, the case we now analyse.

Rotation points p are encircled by arbitrarily small closed orbits. Let S be the set of all closed orbits near p . The set S contains no annulus A of closed orbits because there would exist ε -quasiorbits which are nearly dense in A . They would be singly unshadowable. Similarly an ε -quasiorbit along an ingoing spiral to a semistable closed orbit followed by an outgoing spiral (joined by a C^1 curve) is singly unshadowable.

Thus, the complement to $S \cup \{p\}$ near p consists of a countable number of bands of nonperiodic orbits. By Poincaré Bendixson theory, all the points of such a band B have the same ω -limit set, one of the two boundary curves of B , and all the points have the other boundary curve as their α -limit set. Say that B is *inward* if $\omega(x)$ is inside $\alpha(x)$ for all $x \in B$ and *outward* if the opposite.

FIGURE 3

If between two inward bands B_1 and B_2 there are only inward bands then there exist ε -quasiorbits marching forever inward from B_1 through B_2 which are singly unshadowable. This is the semistable contradiction again.

Thus property H implies (a), (b), or (c). The converse is easily verified and left to the reader. $\square$

REFERENCES

- N. P. Bhatia and G. P. Szegő, *Stability theory of dynamical systems*, Springer-Verlag, Berlin and New York, 1970.
- Z. Nitecki, *Explosions in completely unstable flows. I, Preventing explosions*, Trans. Amer. Math. Soc. **245** (1978), 43–61.
- M. L. A. Peixoto, *The closing lemma for generalized recurrence in the plane*, Trans. Amer. Math. Soc. **308** (1988).
- C. C. Pugh, *The closing lemma*, Amer. J. Math. (2) **89** (1967), 956–1009.
- , *An improved closing lemma and a general density theorem*, Amer. J. Math. **89** (1967), 1010–1021.

- , *Against the C^2 closing lemma*, J. Differential Equations **17** (1975), 435–443.
- , *The C^1 -connecting lemma, a counterexample* (preprint), 1984.

ESCOLA NACIONAL DE CIENCIAS ESTATISTICAS, RUA ANDRÉ CAVALCANTI 106, 20231 RIO DE JANEIRO, R.J., BRASIL

E-mail address: mpe@bridge.bitnet

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CALIFORNIA 94720