DISJOINT PATHS, PLANARIZING CYCLES, AND SPANNING WALKS

XINGXING YU

ABSTRACT. We study the existence of certain disjoint paths in planar graphs and generalize a theorem of Thomassen on planarizing cycles in surfaces. Results are used to prove that every 5-connected triangulation of a surface with sufficiently large representativity is hamiltonian, thus verifying a conjecture of Thomassen. We also obtain results about spanning walks in graphs embedded in a surface with large representativity.

1. INTRODUCTION

We begin with the definition of a k -walk. For a positive integer k , a k -walk in a graph is a walk that visits every vertex at least once and at most k times. Clearly a 1-walk is just a Hamilton cycle.

It is well known that not all 3-connected planar graphs are hamiltonian (for example, see [1]) and that every triangulation of the sphere with no separating triangles is hamiltonian [17]. Tutte [16] proved that every 4-connected planar graph has a Hamilton cycle. Extending Tutte's technique, Thomassen [12] proved that every 4-connected planar graph is in fact Hamilton connected. Thomassen's theorem implies that every 4-connected planar graph minus an arbitrary vertex is hamiltonian. Plummer [9] conjectured that every 4-connected planar graph minus two arbitrary vertices is hamiltonian. This conjecture was proved by Thomas and Yu [10] (by extending the techniques of Thomassen in various ways). With some additional techniques, it was also proved in [10] that every edge in a 4-connected projective-planar graph is contained in a Hamilton cycle, which implies the conjecture of Grünbaum [6] that every 4-connected projective-planar graph has a Hamilton cycle.

Grünbaum [6] and Nash-Williams [8] also conjectured that every 4-connected toroidal graph has a Hamilton cycle. While this is still unsolved, Thomassen [12] showed that there are 4-connected toroidal graphs in which certain edges are not contained in any Hamilton cycle. It is shown in [11] that every edge in a 5-connected graph in the torus is contained in a Hamilton cycle. The technique used in [11] can also be used to solve the conjecture of Molluzzo [7] that 6-connected toroidal graphs are Hamilton connected. We also mention that Duke [3] gave a function $c(g)$ for 2-connected graphs with the property that such a graph with orientable genus g

Received by the editors August 20, 1993.

1991 *Mathematics Subject Classification.* Primary 05C38, 05C10, 57M15.

Key words and phrases. bridge, disjoint paths, embedding, Hamilton cycle, representativity, walk.

Partially supported by NSF grants DMS-9105173 and DMS-9301909.

and connectivity at least $c(g)$ is hamiltonian. In general, this function is not best possible. For $g = 0$ the best value is 4 by Tutte [16]. For $g = 1$ the above-mentioned conjecture of Grünbaum and Nash-Williams would imply that the correct value is 4.

It is not difficult to see that the existence of a k -walk in a graph implies the existence of a spanning tree with maximum degree $k + 1$. Thomassen [14] showed that every triangulation of an orientable surface of genus g with no noncontractible cycle of length less than 2^{3g+4} contains a spanning tree of maximum degree at most 4. This upper bound 4 is best possible. He also raised two related conjectures in [14]. The first states that every 4-connected triangulation of an orientable surface with sufficiently large representativity has a spanning tree with maximum degree at most three. This was established by Ellingham and Gao [4]. The second conjecture in [14] states that every 5-connected triangulation of an orientable surface with sufficiently large representativity is hamiltonian. In this paper, we prove this conjecture for both orientable and non-orientable surfaces. We also prove that 4-connected graphs embedded in surfaces with large representativity admits 2-walks, which generalizes and improves a result of Ellingham and Gao [4]. We finally answer a question of Ellingham (personal communication) about the existence of a 3-walk in a 3-connected graph, which improves a result of Thomassen [14] on spanning trees. The large representativity condition in all the above results is essential as Archdeacon, Hartsfield and Little (reported in [4]) have shown the following: for each n there is a triangulation of some surface which is n -connected and has representativity at least n , and in which every spanning tree has maximum degree at least n .

In order to state our results precisely, we need some terminology. We follow that of Bondy and Murty [1]. Only simple graphs will be considered here. Let G and H be two graphs. The *intersection* and the *union* of G and H , denoted by $G \cap H$ and $G \cup H$, are the graphs $(V(G) \cap V(H), E(G) \cap E(H))$ and $(V(G) \cup V(H), E(G) \cup E(H))$, respectively. A *block* of a graph G is a maximal 2-connected subgraph of G . We view K_2 as being 2-connected. For any path P and $x, y \in V(P) \cup E(P)$, we use xPy to denote the subpath of P between x and y , where xPy does not include x if $x \in E(P)$ or y if $y \in E(P)$.

Let P be a subgraph of a graph G . Let S be a component of $G - V(P)$, and let B be the union of S and all edges with one end in $V(S)$ and the other in $V(P)$. Then B is called a P -*bridge* of G . Every vertex in $a(B) = V(B) \cap V(P)$ is said to be an *attachment* of B on P . This notation should not cause any confusion (as we will always say that “for a P -bridge B of G ”). In the sequel P will always be a path or a set of disjoint paths.

A *plane graph* is a graph embedded in the plane so that no two edges cross. A *plane subgraph* H of a plane graph G is a subgraph of G inheriting the embedding of G . Given a connected plane graph, the boundary of the infinite face is called the *outerwalk* of the graph, or *outercycle* if the graph is 2-connected. A walk is a *facial walk* in a plane graph if it bounds a face of the graph. Suppose that C is a facial walk of a plane graph G and that $x_1, \dots, x_m$ are distinct vertices or edges of G on C . Then we say that $x_1, \dots, x_m$ *appear on C in this order* if walking along C in the clockwise order from x_1 to x_m we encounter x_i before x_{i+1} and after the first visit to x_{i+1} we never encounter $x_1, \dots, x_i$ again. Given $x, y \in E(C) \cup V(C)$ on a facial walk C in a plane graph, we use xCy to denote the minimal subwalk of C from x to y in the clockwise order, where xCy does not include x if $x \in E(C)$ or y if $y \in E(C)$.

For convenience, we adopt the following convention. Let G be a plane graph with outerwalk C , and let x and y be two vertices on C . Then $G + xy$ (or $G + yx$) denotes the plane graph obtained from G by adding an edge between x and y so that $(C - (xCy - \{x, y\})) + xy$ (or $(C - (yCx - \{x, y\})) + yx$) is the outerwalk of the resulting graph.

Finally, we need some parameters for graphs embedded in surfaces. We refer to [13] for other related concepts. Suppose that G is embedded in a surface Σ . We usually do not distinguish between G and its embedding in Σ . A cycle of G is *noncontractible* if it is homotopically nontrivial in Σ , otherwise it is *contractible*. The *representativity* (or *face-width*) of G , denoted by $\rho(G, \Sigma)$, is $\min\{|\Gamma \cap V(G)| : \Gamma \text{ is a homotopically nontrivial simple closed curve in } \Sigma \text{ and } \Gamma \cap G \subset V(G)\}$. (The *edge-width* of G is the length of a shortest noncontractible cycle of G , which is the same as representativity when the graph is a triangulation.) We use $d_{G\Sigma}(x, y)$ to denote $\min\{|\Gamma \cap V(G)| : \Gamma \text{ is a simple curve in } \Sigma \text{ from } x \text{ to } y \text{ and } \Gamma \cap G \subset V(G)\}$. For two disjoint vertex sets C and D in G , let $d_{G\Sigma}(C, D) = \min\{d_{G\Sigma}(x, y) : x \in C \text{ and } y \in D\}$.

We now outline the rest of the paper. In Section 2 we prove a modification lemma which will be used in the proofs in Section 3 on disjoint paths. In Section 4 we generalize Thomassen's results on planarizing cycles of triangulations to general graphs. In Section 5 we prove Thomassen's conjecture that every 5-connected triangulation with large representativity has a Hamilton cycle. In Section 6 we find 2-walks in 4-connected graphs with large representativity and 3-walks in 3-connected graphs with large representativity.

2. PRELIMINARY RESULTS

Throughout the rest of the paper, when we say "by Theorem n (or by Lemma n), we find a path P " we mean that the path P must satisfy the relevant conclusions. Before we are able to describe a type of modification which is needed to simplify our statements, we need several lemmas. The first result (see [11], also independently obtained by D. P. Sanders) ensures a path between two specified vertices and using a specified edge.

Lemma 2.1. *Let G be a connected plane graph with outerwalk C and let $u, v \in V(G)$ and $e \in E(C)$. Suppose that G has a path from u to v using the edge e . Then there is a path P in G from u to v using e such that for every P -bridge B of G , (i) $|a(B)| \leq 3$, and (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap C \neq \emptyset$. $\square$*

The following result [10] guarantees a path between two given vertices which does not use two other specified vertices.

Lemma 2.2. *Let G be a connected plane graph with outerwalk C , and let x, y, u, v be four distinct vertices on C in this order such that $u, v \notin xCy$. Then G contains a path P from x to y missing u and v such that for every $(P \cup \{u, v\})$ -bridge B of G , (i) $|a(B)| \leq 3$, and (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap xCy \neq \emptyset$. $\square$*

The next lemma is similar to Lemma 2.2, and ensures a path between two given vertices and not using three specified vertices. The proof of Lemma 2.3 is similar to that of Lemma 2.7.

Lemma 2.3. *Let G be a connected plane graph with outerwalk C , and let x, y, u, v, w be five distinct vertices on C in this order such that $u, v, w \notin xCy$. Then G has a*

path P from x to y missing u, v, w such that for every $(P \cup \{u, v, w\})$ -bridge B of G , (i) $|a(B)| \leq 4$, (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap xCy \neq \emptyset$. $\square$

The following result [10] finds a path between two vertices using two specified edges.

Lemma 2.4. *Let G be a connected plane graph with outerwalk C , and let $u, v \in V(C)$ and $e, f \in E(C)$ so that u, e, f, v appear on C in this order and G has a path from u to v using e and f . Then G has a path P from u to v using e and f such that for every P -bridge B of G , (i) $|a(B)| \leq 3$, and (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap uCv \neq \emptyset$. $\square$*

For later convenience, we need another simple lemma.

Lemma 2.5. *Let G be a connected plane graph with outerwalk C , and let $x, y, z \in V(C)$ in this order such that $G - z$ has a path from x to y . Then G has a path P from x to y not using z such that for every $(P \cup \{z\})$ -bridge B of G , (i) $|a(B)| \leq 3$, and (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap xCz \neq \emptyset$.*

Proof. We consider $G + zx$. By Lemma 2.1 we find a path Q in the new graph from y to z using zx . Then $P = Q - zx$ is the desired path. $\square$

We are now ready to state our modification lemma, which will be used heavily in proofs in next section. The conditions in this lemma may seem complicated, but they are easy to verify when applied.

Lemma 2.6 (The Modification Lemma). *Let G be a connected plane graph with outerwalk C , and let α and β be two vertices on C with $L = \alpha C \beta$ simple. Let H be a plane subgraph of $G - V(L)$ with outerwalk D such that each $(H \cup L)$ -bridge of G has at least one attachment on L and at most one attachment on H . Suppose there are distinct vertices u and w in $D \cap C$ such that uDw is simple, α, β, w, u are on C in this order, and all attachments of $(H \cup L)$ -bridges on H are contained in uDw . Suppose that Q is a subgraph of H containing u and w such that for every Q -bridge B of H , (1) $|a(B)| \leq 4$, (2) $|a(B)| \leq 3$ if $(B - a(B)) \cap uDw \neq \emptyset$, (3) $B \cap uDw$ is a simple path.*

Then there is a path P in $G - V(Q)$ from α to β such that for every $(P \cup Q)$ -bridge B of G , (i) $|a(B)| \leq 4$ and (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap V(L) \neq \emptyset$.

Note that in Lemma 2.6 we may allow $u = w$ and let $uDw = D$. This note is used in the proof of Lemma 3.3.

Proof of Lemma 2.6. Since G is a plane graph, we can divide L into non-trivial edge-disjoint subpaths xLy of three types according to the Q -bridges of H : (a) for some Q -bridge B of H , xLy is a maximal subpath of L such that x and y belong to some $(H \cup L)$ -bridges of G each with an attachment in $B - a(B)$; (b) for some vertex v of Q , xLy is the maximal subpath of L such that both x and y belong to some $(H \cup L)$ -bridges of G with v as an attachment; (c) the remaining non-trivial edge maximal subpaths of L .

Obviously, each edge of L belongs to exactly one of these subpaths. We label these subpaths (from α to β) as L_i , $i = 1, \dots, m$. Let the endvertices of L_i be x_i and x_{i+1} such that $x_1, \dots, x_m$ are on C in this order, where $x_1 = \alpha$ and $x_{m+1} = \beta$. By (3), distinct L_i 's of type (a) are associated with distinct Q -bridges of H .

If L_i is of type (a), then in the union of L_i , B , and all $(H \cup L)$ -bridges of G with all their attachments in $L_i \cup (B - a(B))$, we find a path P_i from x_i to x_{i+1} not using vertices of $a(B)$ (Lemma 2.2 or 2.3).

If L_i is of type (b), then in the union of L_i and all $(H \cup L)$ -bridges of G with attachments in $L_i \cup \{v\}$, we find a path P_i from x_i to x_{i+1} not using v (Lemma 2.5).

If L_i is of type (c), then in the union of L_i and all $(H \cup L)$ -bridges of G whose attachments are in L_i , we find a path P_i from x_i to x_{i+1} (Lemma 2.1).

It is now easy to see that $P = \bigcup_{i=1}^m P_i$ is the desired path. $\square$

Remark. If in Lemma 2.6 we ask that for every Q -bridge B of H we have that $|a(B)| \leq 3$ and that $|a(B)| \leq 2$ if $(B - a(B)) \cap uDw \neq \emptyset$, then we can require that $|a(B)| \leq 3$ for every $(P \cup Q)$ -bridge B of G . The proof is exactly the same as that for Lemma 2.6 except in modifying L_i 's of type (a), we only apply Lemma 2.2. This remark is used in the proof of Lemma 3.3.

We note that the modification of L (to get P) takes place in the union of $L = \alpha C \beta$, H and all $(H \cup L)$ -bridges of G with attachments in $H \cup L$. In the rest of the paper, when we apply Lemma 2.6 we may simply say **by modifying $\alpha C \beta$ with respect to (H, Q) we get a path P** . We finish this section with a result slightly stronger than Lemma 2.6. But first we need the following easy lemma.

Lemma 2.7. *Let G be a connected plane graph with outerwalk C , and let x, y, u, v, w be five distinct vertices on C in this order with $u, v, w \notin xCy$. Then G contains a path P from x to y missing u, v and w such that for every $(P \cup \{u, v, w\})$ -bridge B of G , (i) $|a(B)| \leq 4$, (ii) $|a(B)| \leq 3$ if $(B - a(B)) \cap wCu \neq \emptyset$, and (iii) if $(B - a(B)) \cap wCu \neq \emptyset$ then $(B - a(B)) \cap uCw = \emptyset$.*

Proof. Let H be the union of xCy and all blocks of $G - \{u, v, w\}$ containing an edge of xCy . Let D be the outerwalk of H . Let $s, t \in yDx$ (in this order on D) with sDt shortest such that s and t are the attachments of some $(H \cup \{u, v, w\})$ -bridges of G containing u and w respectively. In H we use Lemma 2.4 to find a path P from x to y using s and t ($s = t$ is possible). It is easy to verify that P is the desired path. In fact there is at most one P -bridge of G with 4 attachments on P (which occurs only if $s = t$). $\square$

Lemma 2.8. *Let G, L, C, H, D, u, w, Q be as in Lemma 2.6, and let $w_1, \dots, w_m \in D$ (in this order on D) be the attachments of $(H \cup L)$ -bridges of G . Let $s_i L t_i$ be the maximal subpath of L such that s_i and t_i belong to some $(H \cup L)$ -bridges of G with w_i as an attachment and $s_i C t_i = s_i L t_i$, and let $\alpha, s_1, \dots, s_m, \beta$ appear on C in this order. Suppose z_1 and z_2 are vertices on L ($z_1 = z_2$ is possible) such that none of the following occurs: (I) $z_1, z_2 \in s_i L t_i - \{s_i, t_i\}$ for some i , or (II) $z_1, z_2 \in t_i L s_{i+1} - \{t_i, s_{i+1}\}$ for some i .*

Then we can find a path P as in Lemma 2.6 from α to β with an extra condition: either $\{z_1, z_2\} \cap P \neq \emptyset$ or z_1 and z_2 belong to distinct $(P \cup Q)$ -bridges of G .

Proof. As in the proof of Lemma 2.6, we divide L into edge-disjoint paths $L_i = x_i L x_{i+1}$ of three types. It is clear that if z_1 (or z_2) does not belong to any $L_i - \{x_i, x_{i+1}\}$ which is of type (a) or (b), then this path P can easily be found to use z_1 (or z_2). It is also clear that we may assume that $z_1, z_2 \in L_i - \{x_i, x_{i+1}\}$ for some i . Since (I) does not occur, we may assume that L_i is of type (a). Let B be the Q -bridge of H associated with L_i , and let $w_a, \dots, w_b \in B - a(B)$ in this order on

D (where $s_a = x_i$ and $t_b = x_{i+1}$). We now show how to modify L_i to get a path P_i to form the desired path P . We use J_j to denote the union of s_jLt_j and all $(H \cup L)$ -bridges of G with all attachments contained in $s_jLt_j \cup \{w_j\}$.

Suppose first that $z_1 \in s_kLt_k - \{s_k, t_k\}$. Again since (I) does not occur, $z_2 \notin s_kLt_k - \{s_k, t_k\}$. Since $z_1, z_2 \in L_i - \{x_i, x_{i+1}\}$, by symmetry we may assume that $w_{k+1} \in B - a(B)$ (the other case that $w_{k-1} \in B - a(B)$ can be treated similarly). Then we find a path R in B from w_k to w_{k+1} not using vertices of $a(B)$ (Lemma 2.2 or Lemma 2.7). In J_k we find a path R_k from w_k to s_k using z_1 and t_k (Lemma 2.4) if $t_k \neq s_{k+1}$; otherwise we simply find R_k from w_k to s_k not using t_k (Lemma 2.5). In J_{k+1} we find a path R_{k+1} from w_{k+1} to t_{k+1} using s_{k+1} (Lemma 2.1). We then use Lemma 2.6 to modify $x_iL_is_k$ and $t_{k+1}L_ix_{i+1}$ (both with respect to $(B, R \cup a(B))$) to complete P_i (together with R, R_k, R_{k+1}).

Hence we may assume that $z_1 \in t_kLs_{k+1}$ with k maximum and $z_2 \in t_lLs_{l+1}$ with l maximum, where $k \leq l$. Thus $t_{k+1} \neq z_1$ and $t_{l+1} \neq z_2$. Note that if $k = l$, then since (II) does not occur, $\{z_1, z_2\} \cap \{t_k, s_{k+1}\} \neq \emptyset$, and in this case we may assume by symmetry that $z_2 = s_{k+1}$. Thus we may assume that $z_2 \in s_{k+1}Lt_{l+1}$. Since $\{z_1, z_2\} \cap \{x_i, x_{i+1}\} = \emptyset$, we can take $c < k$ to be maximum such that $s_c \neq z_1$ (note that $c = k$ is possible). In B we find a path R from w_c to w_{k+1} not using the vertices in $a(B)$ (Lemma 2.2 or Lemma 2.7). In J_c we find a path R_c from w_c to s_c using t_c if $t_c \neq s_{k+1}$ (Lemma 2.1); otherwise $t_c = s_{k+1} = z_1$ and we just find a path R_c in J_c from w_c to s_c not using t_c (Lemma 2.5). In J_{k+1} we find a path R_{k+1} from w_{k+1} to t_{k+1} using s_{k+1} (Lemma 2.1). We then modify $x_iL_is_c$ and $t_{k+1}L_ix_{i+1}$ (both with respect to $(B, R \cup a(B))$) to complete P_i (together with R, R_c and R_{k+1}). It is easy to check that P_i is as desired. $\square$

We remark here that Lemma 2.8 will be used in the proof of Claim 2 in Theorem 3.4.

3. DISJOINT PATHS IN PLANAR GRAPHS

In this section we study the existence of certain disjoint paths in a planar graph. First we deal with disjoint paths among vertices lying on a single facial cycle.

Lemma 3.1. *Let G be a 2-connected plane graph with outercycle C , let x, u, v, y be four distinct vertices on C in this order, and let e be an edge on yCx . Suppose that G does not have a 2-cut $\{p, q\}$ with $p \in yCe$ and $q \in xCu - \{x\}$ such that the $\{p, q\}$ -bridge of G containing e is 2-connected. Then G has two disjoint paths P and Q with P from x to y using e and Q from u to v such that for every $(P \cup Q)$ -bridge B of G , (i) $|a(B)| \leq 4$, and (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap yCv \neq \emptyset$.*

Proof. Let H be the union of uCv and all blocks of $G - yCx$ containing an edge of uCv . Let $w \in V(H) \cap xCu$ with xCw shortest. By Lemma 2.1 we find a path Q in H from u to v using w . Let $w_i \in H$, $i = 1, \dots, m$, be the attachments of $(H \cup yCx)$ -bridges of G (in this order on the outercycle of H , where $w_m = w$). Let s_iCt_i be the maximal subpath of yCx such that s_i and t_i are in some $(H \cup yCx)$ -bridges of G with w_i as an attachment. Define J_i to be the union of s_iCt_i and all $(H \cup yCx)$ -bridges of G with all attachments contained in $s_iCt_i \cup \{w_i\}$. We divide yCx into subpaths $L_j = x_jCx_{j+1}$, $j = 1, \dots, n$, of types (a), (b) and (c) as in the proof of Lemma 2.6.

If $e \in L_j$ of type (c), then in the union of x_jCx_{j+1} and all $(H \cup yCx)$ -bridges of G with all attachments in x_jCx_{j+1} we use Lemma 2.1 to find a path P_j from x_j to

x_{j+1} using e . We now modify yCx_j and $x_{j+1}Cx$ (both with respect to (H, Q)) to get two paths which, together with P_j , give the desired path P from x to y using e . Hence we may assume that $e \in L_j$ of type (a) or (b).

Case 1. Suppose that $e \in t_kCs_{k+1} \subset L_j$ of type (a). Let B be the Q -bridge of H associated with L_j . Then $w_k, w_{k+1} \in B - a(B)$. In the union of t_kCs_{k+1} and all $(H \cup yCx)$ -bridges of G with all attachments contained in t_kCs_{k+1} , we find a path S from t_k to s_{k+1} using e (Lemma 2.1).

(1.1) If there is a $w_l \in B - a(B)$ with $l < k$ such that $s_l \neq t_k$, then take such w_l with l maximum. In B we find a path R from w_l to w_k (Lemma 2.2) not using vertices in $a(B)$. In J_k we find a path R_k from w_k to t_k using s_k if $s_k \neq s_l$ (Lemma 2.1), otherwise not using s_k (Lemma 2.5). In J_l we find a path R_l from w_l to s_l using t_l if $t_l \notin R_k$ (Lemma 2.1), otherwise not using t_l (Lemma 2.5). Finally modify yCs_l and $s_{k+1}Cx$ (both with respect to $(H, Q \cup R)$) to get two paths which, together with R, R_k, R_l and S , give the desired path P .

(1.2) Therefore we may assume that if $w_l \in B - a(B)$ with $l < k$ then $s_l = t_k$. By symmetry we may also assume that if $w_l \in B - a(B)$ with $l > k + 1$ then $t_l = s_{k+1}$. Let $R_k = \{s_k\}$ if $s_k = t_k$; otherwise we find a path R_k in J_k from s_k to t_k such that $w_k \in R_k$ or the R_k -bridge of J_k containing w_k has one attachment. We define R_{k+1} similarly. We modify yCs_k and $t_{k+1}Cx$ (both with respect to (H, Q)) to get two paths which, together with S, R_k, R_{k+1} , give the desired path P .

Case 2. Hence let $e \in J_k$ for some k . By our 2-cut assumption, $w_k \neq w_m$. If $w_k \in Q$, then we find a path P_k in $J_k - w_k$ from s_k to t_k using e (Lemma 2.1). If $w_k \in B - a(B)$ for some Q -bridge of H and $w_i \notin B - a(B)$ for every $i \neq k$, then we find a path P_k in J_k from s_k to t_k using e (Lemma 2.1). In either case, we modify yCs_k and t_kCx (both with respect to (H, Q)) to obtain two paths which, together with P_k , give the desired path P . Thus let $w_k \in B - a(B)$ for some Q -bridge B of H , and by symmetry we may assume that $w_{k-1} \in B - a(B)$.

(2.1) If there is some $l < k$ such that $w_l \in B - a(B)$ and $s_l \neq s_k$, we select l to be maximum. First we find a path R in B from w_l to w_k not using vertices in $a(B)$ (Lemma 2.2). Then in J_k we find a path R_k from w_k to t_k using s_k and e (Lemma 2.4). Finally we find a path R_l in J_l from w_l to s_l using t_l if $t_l \neq s_k$ (Lemma 2.1), otherwise not using t_l (Lemma 2.5). Now we just modify yCs_l and t_kCx (both with respect to $(H, Q \cup R)$) to get two paths P_1 and P_2 . It is easy to check that $P = R \cup R_k \cup R_l \cup P_1 \cup P_2$ is the desired path.

Hence we may assume that $s_i = s_k$ for every $w_i \in B - a(B)$ with $i < k$ and by symmetry that $t_i = t_k$ for every $w_i \in B - a(B)$ with $i > k$.

(2.2) Suppose $w_{k+1} \notin B - a(B)$. Then $w_{k-1} \in B - a(B)$. First we consider the case that s_k is not incident to e . If there is a path in $J_k - s_k$ from w_k to t_k using e , then we let N be the union of such a path and all blocks of $J_k - s_k$ containing an edge of that path. Let $z \in V(N) \cap C$ with s_kCz shortest. By Lemma 2.4 we find a path R_k in $J_k - s_k$ from w_k to t_k using z and e . Now in B we find a path R by Lemma 2.2 from w_k to w_{k-1} not using vertices in $a(B)$. In J_{k-1} we find a path R_{k-1} from w_{k-1} to $s_{k-1} = s_k$. We modify yCs_k and t_kCx (both with respect to $(H, Q \cup R)$) to get the desired path P . Hence we may assume that such a path in $J_k - s_k$ from w_k to t_k using e does not exist, and therefore G has a 2-cut $\{s_k, p\}$ with $p \in eCt_k$. We may select this 2-cut so that pCt_k is shortest.

If $p = t_k$, we then just find a path in J_k from s_k to t_k using e , and then modify yCs_k and t_kCx (both with respect to (H, Q)) to get the desired path P . Hence let $p \neq t_k$. Let T be the $\{s_k, p\}$ -bridge of J_k containing w_k . Then in $T - s_k$ we find a path R_1 from p to t_k such that $w_k \in R_1$ or the R_1 -bridge of $T - s_k$ containing w_k has exactly one attachment (Lemma 2.1), and in the other $\{s_k, p\}$ -bridge of J_k containing e , we find a path R_2 from s_k to p using e (Lemma 2.1). Now we can modify yCs_k and t_kCx (both with respect to (H, K)) to get the desired path P .

This reduces (2.2) to the case that s_k is incident to e . If t_k is also incident to e , we then just modify yCs_k and t_kCx (both with respect to (H, Q)), and then add the edge $e = s_k t_k$ to get the desired path P . So t_k is not incident to e . Let s be the end of e other than s_k and let S be the union of sCt_k and all blocks of $J_k - s_k$ containing an edge of sCt_k . Let $p \in S$ such that $p = w_k$ if $w_k \in S$; otherwise $p \in S$ is the cutvertex of $J_k - s_k$ separating w_k from sCt_k . Now we find a path in S from s to t_k using p (Lemma 2.1), and then we modify yCs_k and t_kCx (both with respect to (H, Q)) to get the desired path P (using e).

(2.3) Thus by symmetry we may assume that $w_{k-1}, w_{k+1} \in B - a(B)$. If $s_k Ct_k = s_k t_k$, then we just modify yCs_k and t_kCx (both with respect to (H, Q)) to get two paths which, together with e , give the desired path P . Hence by symmetry we may assume that s_k is not incident to e .

If in $J_k - s_k$ there is a path from w_k to t_k using e , we then let N and z be defined as in (2.2), and the same argument in (2.2) also works here. Hence we may assume that such a path does not exist. Again, as in (2.2) we have the same 2-cut $\{s_k, p\}$ with pCt_k shortest, and the case for $p = t_k$ is the same as in (2.2). So let $p \neq t_k$. We first find a path R in B from w_k to w_{k+1} not using the vertices in $a(B)$ (Lemma 2.2). Let S and T be the $\{s_k, p\}$ -bridges of J_k containing e and w_k , respectively. Since $p \neq t_k$, in $T - s_k$ we can find a path R_1 from p to w_k not using t_k (Lemma 2.5). In S we find a path R_2 from s_k to p using e (Lemma 2.1). In J_{k+1} we find a path R_3 from w_{k+1} to $t_{k+1} = t_k$. Finally, we modify yCs_k and t_kCx (with respect to $(H, Q \cup R)$) to get the desired path P (together with R, R_1, R_2 , and R_3). $\square$

Lemma 3.2. *Let G be a 2-connected plane graph with outercycle C , and let x, y, u, v, w be five distinct vertices on C in this order. Suppose that G does not have two disjoint paths X and Y , where X is from x to v using w and Y is from y to u or X is from x to w and Y is from y to v using u , such that for every $(X \cup Y)$ -bridge B of G , (i) $|a(B)| \leq 4$, and (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap wCu \neq \emptyset$. Then G has two disjoint paths V and W with V from x to w and W from y to u using v such that for every $(V \cup W)$ -bridge B of G , (i) $|a(B)| \leq 4$, and (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap wCu \neq \emptyset$.*

Proof. By Lemma 3.1 we know that G has 2-cuts $\{p, q\}$ and $\{r, s\}$, where v, p, w, x, q, y and x, r, y, u, s, v appear on C in this order, such that the $\{p, q\}$ -bridge L containing wCx and the $\{r, s\}$ -bridge R containing yCu are 2-connected. Therefore x, q, r, y must appear on C in this order, which implies that $r \neq x$ and $q \neq y$. (Note that $q = r$ is possible.) Also by Lemma 3.1 we may assume that $p \neq w$ and $s \neq u$ (as we can choose the edge e at w on vCw when applying Lemma 3.1). Let $M = G - ((L \cup R) - \{p, q, r, s\})$.

In $L + pw$ we find a path R_1 from x to q using pw (Lemma 2.1). In $M + pq$ we find a path R_2 from r to s using pq and v (Lemma 2.4). In $R + sr$ we find a path R_3 from y to u using sr (Lemma 2.1). Then $(R_1 - pw) \cup (R_2 - pq) \cup (R_3 - sr)$ gives

the desired paths V and W , where W uses v . (By a symmetric argument we can ask that V use v .) $\square$

We now turn our attention to disjoint paths among vertices on two facial cycles.

Lemma 3.3. *Let G be a 2-connected plane graph with outercycle C and another facial cycle D disjoint from C , and let $x, y \in C$ and $u, v \in D$ be four distinct vertices of G . Then G has two disjoint paths P and Q with P from x to y and Q from u to v such that for every $(P \cup Q)$ -bridge B of G we have $|a(B)| \leq 3$.*

Note that the sole purpose of $C \cap D = \emptyset$ is to ensure that no $(P \cup Q)$ -bridge B of G exists such that both $(B - a(B)) \cap C \neq \emptyset$ and $(B - a(B)) \cap D \neq \emptyset$.

Proof of Lemma 3.3. Let H be the block of $G - xCy$ containing D . If $H \cap (yCx - \{x, y\}) \neq \emptyset$, then we use Lemma 2.1 to find a path Q in H from u to v using a vertex in $H \cap (yCx - \{x, y\})$. Now we modify xCy with respect to (H, Q) to get the desired path P (the note and the remark following Lemma 2.6). Thus we may assume that $H \cap yCx = \emptyset$.

Suppose that H and $yCx - \{x, y\}$ belong to different components of $G - xCy$. Then G has a 2-cut $\{p, q\}$ separating H from $yCx - \{p, q\}$, where x, p, q appear on C in this order. Let T be the $\{p, q\}$ -bridge of G containing yCx . Let F be the outercycle of H . Let $J_1, \dots, J_m$ be the $(H \cup xCy)$ -bridges of G with an attachment in H such that $J_1 \cap C, \dots, J_m \cap C$ appear on C in this order. Let $J_i \cap H = \{r_i\}$. In H we find a path Q from u to v using a vertex in $r_m Fr_1$ (Lemma 2.1). In $T + pq$ we find a path P_1 from x to y using pq (Lemma 2.1). Now we modify pCq (with respect to (H, Q)) to get a path P_2 (by the note and the remark following Lemma 2.6). It is easy to see that $(P_1 - pq) \cup P_2$ gives the desired P .

Therefore we may assume that H and $yCx - \{x, y\}$ belong to a common component of $G - xCy$. Then G has a 3-cut $\{p, q, w\}$ with x, p, q appearing on C in this order and $w \in H$ is the cutvertex of $G - xCy$ separating $yCx - \{x, y\}$ from D . Let B be the $\{p, q, w\}$ -bridge of G containing yCx . In H we find a path Q from u to v using w (Lemma 2.1). In $B + pq - w$ we view the facial walk of $B + pq - w$ which bounds the face containing w as the outerwalk and find a path P_1 from x to y using pq (Lemma 2.1). Finally modifying pCq with respect to (H, Q) , we get a path P_2 (the note and the remark following Lemma 2.6). It is easy to verify that $(P_1 - pq) \cup P_2$ gives the desired path P . $\square$

We are now ready to prove our main results on disjoint paths.

Theorem 3.4. *Let G be a 2-connected plane graph with outercycle C and another facial cycle D disjoint from C . Let $x, y \in V(C)$ and $u, v \in V(D)$ be four distinct vertices. Suppose that (1) any simple closed curve in the plane separating C from D intersects G at least 7 times, and (2) any simple curve in the plane from C to D intersects G at least 8 times. Then G has two disjoint paths P and Q with P from x to y and Q from u to v such that for every $(P \cup Q)$ -bridge B of G , (i) $|a(B)| \leq 4$, and (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap (C \cup D) \neq \emptyset$.*

Proof. Let C' be the outercycle of the block of $G - C$ containing D , and let G' be the block of $G - (C \cup D)$ containing C' . Let D' be the facial cycle of G' (other than C') which is not a facial cycle of G . Our strategy is to find vertices $x', y' \in C'$ and $u', v' \in D'$ (and if necessary, we go to $G' - (C' \cup D')$) and to find paths P' and Q' in G' (as in Lemma 3.3) with P' from x' to y' and Q' from u' to v' , and then

to extend them to our desired paths in G by “modifying” C and D . Because of symmetry we only treat the “ C -side”, the “ D -side” can be treated in the same way.

Let $x'_1, \dots, x'_m \in C'$ (in this clockwise order on C') be the attachments of $(G' \cup C)$ -bridges of G . Let $s_i C t_i$ be the maximal subpath of C such that every $(G' \cup C)$ -bridge of G with x'_i as an attachment has all of its attachments in $s_i C t_i \cup \{x'_i\}$ and that $s_i C t_i - \{s_i, t_i\}$ contains no attachments of $(G' \cup C)$ -bridges of G with x'_j as an attachment for any $j \neq i$. Let J_i be the union of $s_i C t_i$ and all $(G' \cup C)$ -bridges of G with attachments in $s_i C t_i \cup \{x'_i\}$. Let T_i be the union of $s_i C s_{i+1}$ and all the $(G' \cup C)$ -bridges of G with attachments contained in $s_i C s_{i+1} \cup \{x'_i\}$.

We may assume $x \in s_k C s_{k+1} - \{s_k\}$, otherwise $s_i = x$ for each i and so there would be some x'_j such that a simple closed curve separating C from D intersects G only at x'_j and x . Similarly let $y \in s_l C s_{l+1} - \{s_l\}$. By a similar argument, we may also assume that $x \in t_{k'-1} C t_{k'} - \{t_{k'}\}$ and $y \in t_{l'-1} C t_{l'} - \{t_{l'}\}$.

In the case that $k = l$ and $k' = l'$ it is not difficult to see that $x, y \in s_k C t_k - \{s_k, t_k\}$ or $x, y \in t_k C s_{k+1}$ for some k . We may assume that s_k, x, y are on C in this clockwise order.

The rest of the proof is roughly divided into three stages. First we describe x', y', u' and v' . Then we find paths P' and Q' with P' from x' to y' and Q' from u' to v' . In the last stage we extend P' and Q' to P and Q , respectively.

Stage 1. This stage describes x', y', u', v' . We distinguish three cases: (a), (b) and (c).

(a) $k \neq l$, or $k' \neq l'$, or $x, y \in t_k C s_{k+1}$ for some k .

If $k \neq l$, then let $x' = x'_k$ and $y' = x'_l$. If $k' \neq l'$, then let $x' = x'_{k'}$ and $y' = x'_{l'}$. If $x, y \in t_k C s_{k+1}$, then we may select $x' = x'_k$ and $y' = x'_p$, where $x'_p C' x'_k$ is shortest with $s_p \neq t_k$. Note that such x'_p exists, otherwise there would be a simple closed curve in the plane separating C from D intersecting G only at s_k , contradicting (1).

Now the remaining cases will be for $x, y \in s_k C t_k - \{s_k, t_k\}$. It is clear that $s_k C t_k$ is contained in a block J of J_k . Let $z = x'_k$ if $J = J_k$; otherwise let $z \in J$ be the cutvertex of J_k separating x'_k from $s_k C t_k$.

(b) There are two disjoint paths X and Y in J with X from x to z using s_k and Y from y to t_k (or symmetrically, X from x to s_k and Y from y to z using t_k) such that for every $(X \cup Y)$ -bridge B of J , (i) $|a(B)| \leq 4$, and (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap s_k C t_k \neq \emptyset$.

In this case, we let $x' = x'_k$ and $y' = x'_p$, where $x'_p C' x'_k$ is shortest (or $x'_k C' x'_p$ is shortest) such that $s_p \neq s_k$ (or $t_p \neq t_k$). Note that such a vertex x'_p always exists, otherwise there would be a simple closed curve in the plane separating C from D which intersects G only at s_k, x'_k (or t_k, x'_k), contradicting (1).

(c) These two paths X and Y in (b) do not exist.

By Lemma 3.2, J contains two disjoint paths V and W with V from x to s_k using z and W from y to t_k such that for every $(V \cup W)$ -bridge B of J , (i) $|a(B)| \leq 4$, and (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap s_k C t_k \neq \emptyset$.

By condition (2) let H be the block of $G' - z$ containing D' and let E be the outercycle of H . By (1), $H \cap C' \neq \emptyset$. If $z \neq x'_k$, then $E = C'$ and in this case let $z_1 = z_2 = x'_k$. If $z = x'_k$, then let $z_1, z_2 \in C'$ be the attachments of $(H \cup \{z\})$ -bridges of G' such that z_1, z, z_2 appear on C' in this order. Note that $z_1 \neq z_2$, for otherwise there would be a simple closed curve in the plane separating

C from D which intersects G only at $z_1 = z_2$ and $x'_k = z$, contradicting (1). Also note that $z_2 C' z_1 = z_2 E z_1 = E \cap C'$. Let N be the block of $H - E$ containing D' , and let F be the outercycle of N . Let $w_1, \dots, w_n \in F$ (in this order on F) be the attachments of $(N \cup E)$ -bridges of H . We define $s'_i E t'_i$, J'_i and T'_i in H with respect to w_i, E and F much as we defined $s_i C t_i$, J_i and T_i in G with respect to x'_i, C and C' .

Claim 1. We can find $x'_a, x'_b \in z_2 C' z_1 - \{z_1, z_2\}$ with z_2, x'_a, x'_b in this order on C' such that (i) $s_a \neq t_b$ and $J_i \cap C = \{s_a = t_a\}$ for all $x'_i \in x'_a E x'_b - \{x'_a, x'_b\}$ or $J_i \cap C = \{s_b = t_b\}$ for all $x'_i \in x'_a E x'_b - \{x'_a, x'_b\}$, and (ii) no $w_c \in F$ exists with $x'_a, x'_b \in s'_c E t'_c - \{s'_c, t'_c\}$.

Proof of Claim 1. We first prove that we can select x'_a and x'_b satisfying (i). If $x'_i \notin z_2 C' z_1 - \{z_1, z_2\}$ for every i , then there is a simple closed curve in the plane separating C from D which only intersects G at z_1, z_2 and x'_k , a contradiction. If $z_2 C' z_1 - \{z_1, z_2\}$ contains exactly one x'_i , then there would be a simple closed curve in the plane separating C from D which only intersects G at x'_i, z_1, z_2 and x'_k . So we may assume that $z_2 C' z_1 - \{z_1, z_2\}$ contains at least two x'_i . Then this pair x'_a and x'_b can be selected unless $s_i = t_i = s$ for every $x'_i \in z_2 C' z_1 - \{z_1, z_2\}$. But then there is a simple closed curve in the plane separating C from D which intersects G only at z_1, z_2, s and x'_k , a contradiction.

Now suppose that (ii) is violated for all of those x'_a, x'_b satisfying (i), that is, there is a w_c such that $x'_a, x'_b \in s'_c E t'_c - \{s'_c, t'_c\}$.

Note that s'_c, z_2, z_1, t'_c cannot appear on E in this order, otherwise there would be a simple closed curve in the plane separating C from D which intersects G only at w_c, s'_c, t'_c and x'_k . We now show that z_1, z_2 cannot both be contained in $s'_c C' t'_c$. Suppose $z_1, z_2 \in s'_c C' t'_c$. Then there is $x'_r \in t'_c C' s'_c$, otherwise there would be a simple closed curve separating C from D which intersects G only at s'_c, t'_c and w_c . We may select x'_r so that $x'_b C' x'_r$ is shortest if $x'_a, x'_b \in z_2 C' t'_c$ or $x'_r C' x'_a$ is shortest if $x'_a, x'_b \in s'_c C' z_1$. It is easy to see that there is a $x'_d \in x'_a C' x'_r$ if $x'_a, x'_b \in z_2 C' t'_c$ or $x'_d \in x'_r C' x'_b$ if $x'_a, x'_b \in s'_c C' z_1$, such that x'_d and x'_r form the desired pair (using the fact that x'_a and x'_b satisfy (i)), a contradiction.

Therefore we can select $x'_r \in z_2 C' s'_c - \{z_2\}$ (or $x'_r \in t'_c C' z_1 - \{z_1\}$) such that $x'_r C' x'_a$ (or $x'_b C' x'_r$) is shortest; otherwise there would be a simple closed curve in the plane separating C from D which intersects G only (possibly) at $s'_c, t'_c, w_c, z_1, z_2$ and x'_k . It is now easy to check (by using the fact that x'_a and x'_b satisfy (i)) that we can find $x'_d \in x'_r C' x'_b$ (or $x'_d \in x'_a C' x'_r$) such that x'_r and x'_d give the desired pair, a contradiction. Hence Claim 1 follows.

By (ii) of Claim 1, we can find $w_p \in F$ such that $x'_a, x'_b \in t'_p C' s'_{p+1}$ or we can find $w_p, w_q \in F$ with $w_p \neq w_q$ such that $x'_a \in s'_p E s'_{p+1} - \{s'_p\}$ and $x'_b \in s'_q E s'_{q+1} - \{s'_q\}$ or (symmetrically) $x'_a \in t'_{p-1} E t'_p - \{t'_p\}$ and $x'_b \in t'_{q-1} E t'_q - \{t'_q\}$. In the former case we let $y' = w_p$ and $x' = w_q$, where $w_q F w_p$ is shortest with $s'_q \neq t'_p$. In the latter case we let $x' = w_p$ and $y' = w_q$.

By a similar treatment we have D', u', v' , and L at the “ D -side” precisely the same as C', x', y' , and F at the “ C -side”. By (2) we see that $F \cap L = \emptyset$.

Stage 2. We now find P' and Q' . Let H' be the plane subgraph of G such that: $H' = G'$ if $x', y' \in C'$ and $u', v' \in D'$; H' is the subgraph of G between C' and L (i.e., the subgraph of G contained in the closed annulus bounded by C' and L) if $x', y' \in C'$ and $u', v' \in L$; H' is the subgraph of G between F and D' if $x', y' \in F$

and $u', v' \in D'$; or H' is the subgraph of G between F and L if $x', y' \in F$ and $u', v' \in L$. For convenience we use F' and L' to denote the two facial cycles of H' which are not facial cycles of G , where F' is the outercycle. Hence $F' = F$ or $F' = C'$ and $L' = L$ or $L' = D'$.

By (2) we have $F' \cap L' = \emptyset$. Hence by Lemma 3.3 we can find two disjoint paths P' and Q' in H' with P' from x' to y' and Q' from u' to v' such that for every $(P' \cup Q')$ -bridge B of H' we have that $|a(B)| \leq 3$. Note that if $(B - a(B)) \cap F' \neq \emptyset$, then $(B - a(B)) \cap L' = \emptyset$.

Stage 3. We now extend P' to a path from x to y and Q' to a path from u to v . By symmetry we only show how to extend P' . Suppose first that $F' = C'$. Then we have (a) or (b). For (a) we may first assume that $x \in s_k C s_{k+1} - \{s_k\}$ and $y \in s_l C s_{l+1} - \{s_l\}$ with $x'_k \neq x'_l$. (Note that the case that $x \in t_{k'-1} C t_{k'} - \{t_{k'}\}$ and $y \in t_{l'-1} C t_{l'} - \{t_{l'}\}$ with $x'_{k'} \neq x'_{l'}$ can be treated in the same way.) In this case $x' = x'_k$ and $y' = x'_l$. In $T_k + s_{k+1}x'_k$ we find a path Q_k from x to s_k using $s_{k+1}x'_k$ (Lemma 2.1). In $T_l + s_{l+1}x'_l$ we find a path Q_l from y to s_l using $s_{l+1}x'_l$ (Lemma 2.1). We then modify $s_{k+1}C s_l$ and $s_{l+1}C s_k$ (both with respect to $(H', P' \cup Q')$) to get two paths R_1 and R_2 . It is easy to see that $P = P' \cup (Q_k - s_{k+1}x'_k) \cup (Q_l - s_{l+1}x'_l) \cup R_1 \cup R_2$ is as desired. So we may assume that $x, y \in t_k C s_{k+1}$ and s_k, x, y on C in this order. In this case $x' = x'_k$ and $y' = x'_p$. In J_p we find a path Q_p from x'_p to s_p using t_p if $t_p \neq t_k$ (Lemma 2.1), otherwise not using t_p (Lemma 2.5). In J_k we find a path Q_k from x'_k to t_k using s_k if $s_k \notin Q_p$ (Lemma 2.1), otherwise not using s_k (Lemma 2.5). In the union of the edge $s_{k+1}t_k$ and the $\{s_{k+1}, t_k\}$ -bridge of G containing x, y we find a path R_k from x to y using $s_{k+1}t_k$ (Lemma 2.1). Finally we modify $s_{k+1}C s_p$ with respect to $(H', P' \cup Q')$ to get a path R . Then $P = P' \cup Q_k \cup Q_p \cup (R_k - s_{k+1}t_k) \cup R$ is the desired path.

For (b) we use X and Y and $x' = x'_k$ and $y' = x'_p$. By symmetry we only consider the case when X is from x to z using s_k . In J_p we find a path Q_p from x'_p to s_p using t_p if $t_p \neq s_k$ (Lemma 2.1), otherwise not using t_p (Lemma 2.5). Let $Z = \{x'_k\}$ if $z = x'_k$, otherwise let Z be a path (Lemma 2.1) from z to x'_k in the z -bridge of J_k containing x'_k . Now we modify $t_k C s_p$ with respect to $(H', P' \cup Q')$ to get a path R . Hence $P = P' \cup X \cup Y \cup Z \cup Q_p \cup R$ is as desired.

Thus we have (c), i.e., $F' = F$. We first extend P' to a path P^* in H from x'_a to x'_b . Suppose first $x'_a \in s'_p E s'_{p+1} - \{s'_p\}$ and $x'_b \in s'_q E s'_{q+1} - \{s'_q\}$, where $w_p \neq w_q$. (The case that $x'_a \in t'_{p-1} E t'_p - \{t'_p\}$ and $x'_b \in t'_{q-1} E t'_q - \{t'_q\}$ with $w_p \neq w_q$ can be treated in the same way.) In this case $x' = w_p$ and $y' = w_q$. In $T'_p + s'_{p+1}w_p$ we find a path Q'_p from x'_a to s'_p using $s'_{p+1}w_p$. In $T'_q + s'_{q+1}w_q$ we find a path Q'_q from x'_b to s'_q using $s'_{q+1}w_q$. We modify $s'_{q+1}E s'_p$ and $s'_{p+1}E s'_q$ (both with respect to $(H', P' \cup Q')$) to get two paths R'_1 and R'_2 . Let $P^* = P' \cup (Q'_p - s'_{p+1}w_p) \cup (Q'_q - s'_{q+1}w_q) \cup R'_1 \cup R'_2$ (a path from x'_a to x'_b).

Hence by the above claim and by symmetry we may assume that $t'_p, x'_a, x'_b, s'_{p+1}$ are on C' in this order. In this case we have $x' = w_q$ and $y' = w_p$. In J'_q we find a path Q'_q from w_q to s'_q using t'_q (Lemma 2.1) if $t'_q \neq t'_p$, otherwise not using t'_q (Lemma 2.5). In J'_p we find a path Q'_p from w_p to t'_p using s'_p if $s'_p \notin Q'_q$ (Lemma 2.1), otherwise not using s'_p (Lemma 2.5). In the union of the edge $s'_{p+1}t'_p$ and the $\{t'_p, s'_{p+1}\}$ -bridge of H containing $x'_a C' x'_b$ we find a path R'_p from x'_a to x'_b using $s'_{p+1}t'_p$ (Lemma 2.1). Finally we modify $s'_{p+1}E s'_q$ with respect to $(H', P' \cup Q')$ to get a path R' . Let $P^* = P' \cup Q'_p \cup Q'_q \cup (R'_p - s'_{p+1}t'_p) \cup R'$. Note that $|a(B)| \leq 2$ if the P^* -bridge B of H has $(B - a(B)) \cap E \neq \emptyset$.

Suppose that z_1, z_2 do not both belong to $B - a(B)$ for any P^* -bridge B of H . In this case either $z = x'_k$ and $z_1 \neq z_2$ or $z \neq x'_k$ and $x'_k \in P^*$. We find a path Q_a in J_a from x'_a to s_a using t_a if $t_a \neq t_b$ (Lemma 2.1), otherwise not using t_a (Lemma 2.5). In J_b we find a path Q_b from x'_b to t_b using s_b if $s_b \notin Q_a$ (Lemma 2.1), otherwise not using s_b (Lemma 2.5). Finally we modify $t_k C s_a$ and $t_b C s_k$ (both with respect to $(G', P^* \cup Q' \cup \{x'_k\})$) to get two paths R_1 and R_2 . Then $P = P^* \cup V \cup W \cup Q_a \cup Q_b \cup R_1 \cup R_2$ is as desired.

Hence we may assume that $z_1, z_2 \in B - a(B)$ for some P^* -bridge B of H . We may also assume that both $x'_{k-1}, x'_{k+1} \in B - a(B)$; otherwise we can directly modify C exactly the same way as in the previous paragraph. Let $a(B) = \{v_1, v_2\}$ such that v_1, z_1, z_2, v_2 appear on C' in this order. We select P^* so that B is minimal.

Claim 2. There is some $w_d \in F$ such that $z_1, z_2 \in s'_d E t'_d - \{s'_d, t'_d\}$ or $z_1, z_2 \in t'_d E s'_{d+1} - \{t'_d, s'_{d+1}\}$.

Proof of Claim 2. Suppose this claim is not true. By Lemma 2.8 we have $z_1, z_2 \in s'_q E s'_{p+1}$ when $x'_a, x'_b \in t'_p C' s'_{p+1}$ (since in this case we modify $s'_{p+1} E s'_q$), and $z_1, z_2 \in T'_p$ or $z_1, z_2 \in T'_q$ when $x'_a \in T'_p$ and $x'_q \in T'_q$ with $w_p \neq w_q$ (since in this case we modify $s'_{q+1} E s'_p$ and $s'_{p+1} E s'_q$); otherwise we can modify E by Lemma 2.8 so that there is no P^* -bridge B of H with $z_1, z_2 \in B - a(B)$. If $z_1, z_2 \in s'_q E s'_{p+1}$, then $z_1, z_2 \in J'_q$ or $z_1, z_2 \in J'_p$ (otherwise $z_1, z_2 \in t'_q E s'_{q+1} - \{t'_q, s'_{q+1}\}$). By symmetry let $z_1, z_2 \in J'_q$. Then $\{z_1, z_2\} \cap \{s'_q, t'_q\} \neq \emptyset$ (otherwise $(Q'_q \cup Q'_p) \cap \{z_1, z_2\} \neq \emptyset$), but then $z_1, z_2 \in s'_q E t'_q - \{s'_q, t'_q\}$, a contradiction.

So by symmetry, let $z_1, z_2 \in T'_p$. Moreover, we may assume that $z_1 \in s'_p E t'_p$ and $z_2 \in t'_p E s'_{p+1}$ (otherwise the claim would follow). In this case we can change the part $Q'_p - s'_{p+1} w_p$ of P^* so that no P^* -bridge B of H exists with $z_1, z_2 \in B - a(B)$. First note that $x'_a \notin s'_p E t'_p$, otherwise z_1 and z_2 do not belong to the same P^* -bridge of H , a contradiction. So $x'_a \in t'_p E s'_{p+1} - \{t'_p\}$. Then we find a path in J'_p from w_p to s'_p using t'_p (Lemma 2.1). In the union of $t'_p E s'_{p+1}$ and all $(E \cup N)$ -bridges of H with attachments contained in $t'_p E s'_{p+1}$, we find a path from x'_a to s'_{p+1} not using t'_p (Lemma 2.5). Now the union of these two paths is used to replace the $Q'_p - s'_{p+1} w_p$ part of P^* . Clearly no P^* -bridge B of H exists such that $B - a(B)$ contains both z_1 and z_2 , a contradiction.

By Claim 2, we have two cases. Note that we may change x' and y' in the following arguments. Also note that since $|a(B)| = 2$, $v_1, v_2 \in s'_d E t'_d$ or $v_1, v_2 \in t'_d E s'_{d+1}$ according as $z_1, z_2 \in s'_d E t'_d - \{s'_d, t'_d\}$ or $z_1, z_2 \in t'_d E s'_{d+1} - \{t'_d, s'_{d+1}\}$.

(I) $z_1, z_2 \in s'_d E t'_d - \{s'_d, t'_d\}$.

By (1) we may assume that s'_d, z_1, z_2 are on E in this order. Note that $x_{k-1}, x_{k+1} \in B - a(B)$. Hence either there are $x'_g \in x'_k C' t'_d - \{x'_k, t'_d\}$ and $x'_h \in t'_d C' s'_d - \{s'_d\}$ such that $s_g \neq t_h$ and $s_i = t_i = s_g$ for all $x'_i \in x'_g C' x'_h - \{x'_g, x'_h\}$ (or $s_i = t_i = t_h$ for all $x'_i \in x'_g C' x'_h - \{x'_g, x'_h\}$) or there are $x'_g \in s'_d C' x'_k - \{x'_k, s'_d\}$ and $x'_h \in t'_d C' s'_d - \{t'_d\}$ such that $s_h \neq t_g$ and that $s_i = t_i = t_g$ for all $x'_i \in x'_h C' x'_g - \{x'_h, x'_g\}$ (or $s_i = t_i = s_h$ for all $x'_i \in x'_h C' x'_g - \{x'_h, x'_g\}$). For otherwise we deduce that $s_i = t_i = w$ for every $x'_i \in t'_d C' s'_d$. Thus we have a simple closed curve in the plane separating C from D which intersects G only at s'_d, t'_d, w, w_d , contradicting (1).

Without loss of generality we may assume that $x'_g \in x'_k C' t'_d - \{x'_k, t'_d\}$ and $x'_h \in t'_d C' s'_d - \{s'_d\}$. Note that we may select $w_r \in F - w_d$ so that $x'_h \in t'_{r-1} C' t'_r - \{t'_r\}$, for otherwise $s'_i E t'_i \subset t'_d E x'_h$ for each $i \neq d$ and so there is a simple closed curve

in the plane separating C from D which intersects G only at w_d, t'_d, x'_h (and s_g or t_h if $s_g = t_g$ or $s_h = t_h$), violating (1). We now change x' and y' in H' to w_d and w_r , respectively, and use Lemma 3.3 to find the two paths P' and Q' with P' from $x' = w_d$ to $y' = w_r$. In $J'_d + w_d s'_d$ we use Lemma 2.4 to find a path Q'_d from x'_g to t'_d using $w_d s'_d$ and z_1 . (Note that $z_1 = x'_k$ if $z \neq x'_k$.) In the union of $t'_{r-1} E t'_r$, $w_r t'_{r-1}$ and all the $(N \cup E)$ -bridges of H with all attachments in $t'_{r-1} E t'_r \cup \{w_r\}$, we find a path Q'_r from t'_r to x'_h using $w_r t'_{r-1}$. Now we modify $t'_d E t'_{r-1}$ and $t'_r E s'_d$ (both with respect to $(H', P' \cup Q')$) to get two paths R'_1 and R'_2 . Let $P^* = P' \cup (Q'_d - w_d s'_d) \cup (Q'_r - w_r t'_{r-1}) \cup R'_1 \cup R'_2$.

We now extend the path P^* to x and y . In J_g we find a path Q_g from x'_g to s_g using t_g if $t_g \neq t_h$ (Lemma 2.1), otherwise not using t_g (Lemma 2.5). In J_h we find a path Q_h from x'_h to t_h using s_h if $s_h \notin Q_g$ (Lemma 2.1), otherwise not using s_h (Lemma 2.5). Finally we modify $t_k C s_g$ and $t_h C s_k$ (both with respect to $(G', P^* \cup Q')$) to get two paths R_1 and R_2 . Then $P = P^* \cup V \cup W \cup Q_g \cup Q_h \cup R_1 \cup R_2$ is as desired.

(II) $z_1, z_2 \in t'_d E s'_{d+1} - \{t'_d, s'_{d+1}\}$.

In this case it is easy to see that $v_1 = t'_d$ and $v_2 = s'_{d+1}$. We claim that there is $x'_p \in t'_{d+1} E t'_d - \{t'_{d+1}\}$ and $x'_q \in v_1 C' x'_k - \{v_1, x'_k\}$ (or $x'_p \in s'_{d+1} E s'_d - \{s'_d\}$ and $x'_q \in x'_k C' v_2 - \{x'_k, v_2\}$) such that $s_p \neq t_q$ (or $t_p \neq s_q$) and $s_i = t_i = s_p$ for all $x'_i \in x'_p C' x'_q - \{x'_p, x'_q\}$ or $s_i = t_i = t_q$ for all $x'_i \in x'_p C' x'_q - \{x'_p, x'_q\}$ (or $s_i = t_i = s_q$ for all $x'_i \in x'_q C' x'_p - \{x'_p, x'_q\}$ or $s_i = t_i = t_p$ for all $x'_i \in x'_q C' x'_p - \{x'_p, x'_q\}$). Otherwise $s_i = t_i = s$ for each $x'_i \in v_1 C' x'_k - \{v_1, x'_k\}$ and $s_i = t_i = t$ for each $x'_i \in x'_k C' v_2 - \{x'_k, v_2\}$, and if $x'_i \in t'_{d+1} C' s'_d$ then $s_i = t_i = s$ or $s_i = t_i = t$. Hence there would be a simple closed curve in the plane separating C from D which intersects G only at $s, t, v_1 = t'_d, v_2 = s'_{d+1}$, contradicting (1). Hence by symmetry we may assume that $x'_p \in t'_{d+1} E t'_d - \{t'_{d+1}\}$ and $x'_q \in v_1 C' x'_k - \{v_1, x'_k\}$.

Suppose first $x'_p = v_1$. Note that there is a $w_r \in w_{d+1} F w_d - \{w_d, w_{d+1}\}$, otherwise there would be a simple closed curve in the plane separating C from D which intersects G only at w_d and w_{d+1} . We can select such w_r that $w_{d+1} F w_r$ is shortest with $v_2 \neq t'_r$, for otherwise there would be a simple closed curve in the plane separating C from D which only intersects G at w_d, v_2 . Now in H' we replace x' and y' by w_{d+1} and w_r and we find the paths P' and Q' by Lemma 3.3 with P' from w_{d+1} to w_r .

We extend P' to a path P^* in H from $x'_p = v_1$ to x'_q . In J'_{d+1} we find a path R'_{d+1} from w_{d+1} to v_2 using t'_{d+1} if $t'_{d+1} \neq t'_r$ (Lemma 2.1), otherwise not using t'_{d+1} (Lemma 2.5). In J'_r we find a path R'_r from w_r to t'_r using s'_r if $s'_r \notin R'_{d+1}$ (Lemma 2.1), otherwise not using s'_r (Lemma 2.5). In $B - \{v_1, z\}$ we find a path R' from x'_q to v_2 using z_2 (Lemma 2.1). (Note that $x'_k = z_2$ if $z \neq x'_k$.) Now modify $t'_r E v_1$ with respect to $(H', P' \cup Q')$ to get a path to complete P^* . We now extend P^* to P . In J_p we find a path R_p from $v_1 = x'_p$ to s_p using t_p if $t_p \neq t_q$ (Lemma 2.1), otherwise not using t_p (Lemma 2.5). In J_q we find a path R_q from x'_q to t_q using s_q if $s_q \notin R_p$ (Lemma 2.1), otherwise not using s_q (Lemma 2.5). Finally we modify $t_k C s_p$ and $t_q C s_k$ (both with respect to $(G', P^* \cup Q')$) to get two paths R and S . Then $P = P^* \cup V \cup W \cup R_p \cup R_q \cup R \cup S$ is the desired path.

Hence we may assume that $x'_p \neq v_1$. We select w_r with $t'_r \in x'_p E v_1 - \{x'_p\}$ so that $x'_p E t'_r$ is shortest (where $w_r = w_d$ is possible). Now in H' we use Lemma 3.3 to find two paths P' and Q' with P' from $x' = w_r$ to $y' = w_{d+1}$.

We extend P' to a path P^* in H from x'_p to x'_q . In J'_{d+1} we find a path R'_{d+1} from w_{d+1} to t'_{d+1} using v_2 (Lemma 2.1). In the union of $w_r t'_{r-1}$, $t'_{r-1} E t'_r$ and all $(E \cup N)$ -bridges of H with all attachments in $t'_{r-1} E t'_r \cup \{w_r\}$, we find a path R'_r from x'_p to t'_r using $w_r t'_{r-1}$ (Lemma 2.1). In $(B + v_2 v_1) - \{z\}$ we find a path R' from v_2 to x'_q using $v_2 v_1$ (Lemma 2.1). Finally modify $t'_{d+1} E t'_{r-1}$ and $t'_r E v_1$ (both with respect to $(H', P' \cup Q')$) to get two paths which, together with R'_{d+1} , $R'_r - w_r t'_{r-1}$, $R' - v_2 v_1$ and P' , give P^* . By the minimality of B (since $x'_q \neq v_1$), z_1 and z_2 belong to different P^* -bridges of H or $\{z_1, z_2\} \cap P^* \neq \emptyset$. We now extend P^* to the desired path P from x to y . In J_p we find a path R_p from x'_p to s_p using t_p if $t_p \neq t_q$ (Lemma 2.1), otherwise not using t_p (Lemma 2.5). In J_q we find a path R_q from x'_q to t_q using s_q if $s_q \notin R_p$ (Lemma 2.1), otherwise not using s_q (Lemma 2.5). Now we modify $t_k C s_p$ and $t_q C s_k$ (both with respect to $(G', P^* \cup Q')$) to get two paths which, together with P^* , R_p , R_q , V and W , give the desired path P . $\square$

The following lemma is a direct consequence of Lemma 2.1.

Lemma 3.5. *Let G be a 2-connected plane graph with outercycle C and another disjoint facial cycle D , and let $x, y \in V(C)$ and $u, v \in V(D)$ be four distinct vertices. Then G has two disjoint paths from $\{x, y\}$ to $\{u, v\}$ such that for every $(P \cup Q)$ -bridge B of G we have $|a(B)| \leq 3$.* $\square$

The next result plays a similar role as Theorem 3.4 in Sections 5 and 6. Its proof is exactly the same as that of Theorem 3.4 except that (in Stages 2 and 3) we use Lemma 3.5 instead of Lemma 3.3.

Theorem 3.6. *Let G be a 2-connected plane graph with outercycle C and another facial cycle D , and let $x, y \in V(C)$ and $u, v \in V(D)$ be four distinct vertices. Suppose that (1) any simple closed curve in the plane separating C from D intersects G at least 7 times, and (2) any simple curve in the plane from C to D intersects G at least 8 times. Then G has two disjoint paths P and Q from $\{x, y\}$ to $\{u, v\}$ such that for every $(P \cup Q)$ -bridge B of G , (i) $|a(B)| \leq 4$, and (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap (C \cup D) \neq \emptyset$.* $\square$

In order to include the non-orientable surfaces in the rest of this paper, we need a result (similar to Theorem 3.4) for graphs embedded in the projective plane. But first we need the following result from [10].

Lemma 3.7. *Let G be a 2-connected graph embedded in the projective plane, and let e be an edge on a homotopically trivial facial cycle C of G . Then G has a cycle T using e such that for every T -bridge B of G , (i) B is planar and $|a(B)| \leq 3$, and (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap C \neq \emptyset$.*

Theorem 3.8. *Let G be 2-connected graph embedded in the projective plane Σ , and let C be a homotopically trivial facial cycle of G with two vertices $x, y \in C$. Suppose that (1) every simple closed curve in Σ homotopic to C and disjoint from the open disc bounded by C must intersect $V(G)$ at least 7 times, and (2) every nontrivial simple closed curve intersecting the open disc bounded by C intersects G at least 8 times. Then G has a path P from x to y such that for every P -bridge B of G , (i) B is planar and $|a(B)| \leq 4$, and (ii) $|a(B)| \leq 2$ if $(B - a(B)) \cap C \neq \emptyset$.*

Proof. By condition (2), G contains three homotopically trivial disjoint cycles around C . Let C' be the cycle in G disjoint from C and homotopic to C such that the subgraph of G contained in the closed annulus bounded by C and C' is

minimal. We define x'_i, s_i, t_i, J_i exactly the same as in the proof of Theorem 3.4. We also proceed in three stages as in the proof of Theorem 3.4, and we define E and F in the same way (using condition (2)). Now the only difference is that we will use Lemma 3.7 instead of Lemma 3.3. We omit the details. $\square$

4. PLANARIZING CYCLES

The results of this section are similar to those of Thomassen's in [15]. The objective here is to reduce graphs embedded in a surface to graphs embedded in the sphere. The proofs are essentially the same as those of [15], though we work on general graphs instead of triangulations. But for the sake of completeness we sketch the proofs here.

First we need some terminology about embeddings of graphs in surfaces. We refer to [13] for terms not defined here. By a result of [13], every nonplanar embedding of a graph in a surface (i.e., an embedding in which the graph is not contained in a disc) has a noncontractible cycle. If C is a contractible cycle of G which is embedded in a surface Σ other than the sphere, then C bounds a unique open disc which is called the *interior* of C in Σ .

Let G be a graph embedded in a surface Σ . We use $eg(G, \Sigma)$ to denote the number satisfying $|V(G)| - |E(G)| + |F(G)| = 2 - eg(G, \Sigma)$, where $F(G)$ is the set of faces. Suppose that $G = (G_1, \dots, G_m)$ is a sequence of graphs and $\Sigma = (\Sigma_1, \dots, \Sigma_m)$ is a sequence of disjoint surfaces such that G_i is embedded in Σ_i . Then we also say that G is embedded in Σ . For technical reason we let $\rho(G, \Sigma) = \infty$ if Σ is the sphere. We also write $eg(G, \Sigma) = \max\{eg(G_i, \Sigma_i) : i = 1, \dots, m\}$ and $\rho(G, \Sigma) = \min\{\rho(G_i, \Sigma_i) : i = 1, \dots, m\}$.

If $C = x_1 \cdots x_k x_1$ is a cycle in a graph G embedded in a surface Σ with $eg(G, \Sigma) = g$, then by *cutting G and Σ along C* we can form a new graph G' embedded in a new surface Σ' , or a pair of new graphs $G' = (G_1, G_2)$ embedded in a pair of new surfaces $\Sigma' = (\Sigma_1, \Sigma_2)$ (with G_i in Σ_i , respectively): if C is orientation preserving, then we first add a cycle $C' = x'_1 \cdots x'_k x'_1$. If an edge yx_i is incident with C on its left side, we delete yx_i and add instead yx'_i . We finally attach a disc to C and another disc to C' to get the surface Σ' . (Note that we used the same symbol C for cycles in G and G' .) If C is orientation reversing, we delete C and add a new cycle $C' = x_1 \cdots x_k x'_1 \cdots x'_k x_1$: we have a local orientation at each vertex of C when walking along C in Σ , and we add the edge yx_i if yx_i is to the right of C at x_i and yx'_i otherwise. Finally we attach a disc to C' to get the surface Σ' . When C is non-contractible, $eg(G', \Sigma') \leq g - 2$ if C is orientation preserving, and $eg(G', \Sigma') = g - 1$ if C is orientation reversing.

Let $C_1, \dots, C_m$ be a collection of pairwise disjoint cycles in a connected graph G embedded in a surface Σ . If cutting along these cycles we obtain a graph H embedded in a disjoint union of spheres, then we say that $\{C_1, \dots, C_m\}$ is a *set of planarizing cycles* of G in Σ . We use the following convention throughout this section. Given a graph G in a surface Σ , we deal with simple curves which intersect G only at its vertices, and the *length of a simple curve* is the number of vertices in G on this curve.

The following lemma was proved in [15] for triangulations; the proof we present here is essentially the same.

Lemma 4.1. *Suppose that G is a connected graph embedded in a surface Σ (other than the sphere) with $\rho(G, \Sigma) \geq 2k + 2$ (k is a positive integer). Then for every*

vertex $x \in V(G)$ and every integer i with $1 \leq i \leq k$ there is an induced contractible cycle C in G containing x in its interior such that for every $z \in V(C)$ there is a simple curve γ of length $d_{G\Sigma}(z, x) = i + 1$ in the closed disc bounded by C from z to x with $\gamma \cap C = \{z\}$.

Proof. We use induction on k . For $k = 1$, $\rho(G, \Sigma) \geq 4$. Let C be a cycle in G which bounds a disc in Σ containing x such that for every vertex z on C there is a simple curve in Σ from z to x of length 2 in the closed disc bounded by C . If C is induced, then C is the desired cycle. Suppose that C has a chord uv . Let P and Q be the two paths of C between u and v , and let R be a simple curve in the interior of C from u to v meeting G only at u, v and x . Then $R \cup P$ and $R \cup Q$ are contractible. Since $\rho(G, \Sigma) \geq 4$, $R + uv$ is contractible. Hence both $P + uv$ and $Q + uv$ are contractible. We can choose the notation so that $P + uv$ is a contractible cycle containing x and Q in its interior. We replace C by $P + uv$. Continuing this procedure, we eventually obtain the desired cycle.

Hence we may assume that $k \geq 2$. Suppose that we can find desired contractible cycles for $i \leq k - 1$. Let D be the contractible cycle with respect to $k - 1$. We now contract D and its interior to a single vertex x' to obtain a new graph G' in Σ (loops and multiple edges are deleted). It is easy to check that $\rho(G', \Sigma) \geq 4$. Thus by the argument in the above paragraph we find an induced contractible cycle C in G' around x' . It is not difficult to check that C is the desired cycle with respect to $i = k$. $\square$

Let C be a noncontractible cycle of G which is embedded in a surface Σ . We cut G and Σ along C to get G' and Σ' . If C is orientation preserving, then we have two cycles C and C' in G' (embedded in Σ'): in the interior of C and C' , respectively, we add vertices x and x' and join x to each x_i and join x' to each x'_i . If C is orientation reversing, then we have a single cycle C' obtained from C : in this case we simply add a vertex x' in the interior of C' and join x' to each x_i and each x'_i . In either case we have a new graph G'' embedded in Σ' (or possibly in the orientation preserving case, a pair of new graphs $G'' = (G'_1, G'_2)$ embedded in $\Sigma' = (\Sigma_1, \Sigma_2)$).

Lemma 4.2. *Let G be a connected graph embedded in a surface Σ with $\rho(G, \Sigma) \geq 5$. Then for every shortest nontrivial simple closed curve Γ in Σ , G has an induced noncontractible cycle C homotopic to Γ such that for every vertex $z \in C$, $d_{G\Sigma}(z, \Gamma \cap V(G)) \leq 2$ and after cutting along C we have $\rho(G'', \Sigma') \geq \rho(G, \Sigma)/2$.*

Proof. Let Γ be a simple closed curve in Σ intersecting G the least number of times. If Γ is orientation preserving, then (since $\rho(G, \Sigma) \geq 5$) there is a cycle C of G homotopic to Γ (we may specify C to be on the left side of Γ) so that for each vertex $z \in C$, $d_{G\Sigma}(z, \Gamma \cap V(G)) \leq 2$. Suppose that uv is a chord of C . Let P and Q be the two subpaths of C between u and v . Then one of $P + uv$ and $Q + uv$ must be trivial; otherwise let P' (and Q') be a simple curve along P (and Q) from u to v which only meet G in $V(G) \cap \Gamma$ such that $P' \cup Q'$ is a simple closed curve homotopic to Γ . Let T be a simple curve from u to v along uv intersecting G only at u and v . Then $P' \cup T$ is homotopic to $P + uv$ and $Q' \cup T$ is homotopic to $Q + uv$. Since $\rho(G, \Sigma) \geq 5$, $|P' \cap V(G)| \geq 5$ and $|Q' \cap V(G)| \geq 5$, and so $P' + uv$ is a nontrivial simple closed curve shorter than Γ , a contradiction. Hence let $P + uv$ be trivial. We replace C by $Q + uv$ (which is homotopic to Γ). Continuing this procedure, we obtain the desired induced cycle C . The case that Γ is orientation reversing can

be treated in the same way (where we use the local orientation at each vertex of C when we find P' and Q').

We now cut G and Σ along C and obtain G'' in Σ' . We prove that $\rho(G'', \Sigma') \geq \rho(G, \Sigma)/2$. If Σ' is the sphere, then $\rho(G'', \Sigma') = \infty > \rho(G, \Sigma)/2$. So we may assume that Σ' is not the sphere. Let γ be a nontrivial simple closed curve in Σ' .

Suppose first that C is orientation preserving. If γ contains a simple curve γ' from C to C' meeting $C \cup C'$ only at its ends, then in Σ , γ' starts from the left side of C and ends on the right side of C . We can extend γ' slightly if necessary that γ' is from the left side of Γ to the right side of Γ , and let P be the shorter subcurve of Γ between the two ends of γ' . Then $P \cup \gamma'$ is a nontrivial simple closed curve in Σ , and so must intersect G at least $\rho(G, \Sigma)$ times. Hence γ' (and also γ) intersects G'' at least $\rho(G, \Sigma)/2$ times. Therefore we may assume that no subcurve of γ goes from C to C' in Σ' . If γ does not go through x or x' , then γ is a nontrivial simple closed curve in Σ , and so intersects G'' at least $\rho(G, \Sigma)$ times. Thus we may assume that γ goes through x' but not x (or equivalently through x but not x'). Then γ has a subcurve R joining two vertices y and z on C' with $R \cap C' = \{y, z\}$ such that R together with a simple curve from y to z intersecting G'' only at y, z, x' is nontrivial. We may extend R to meet Γ , and let P be the shorter subcurve of Γ between the two ends of R . Then $P \cup R$ is also nontrivial in Σ . Hence R (and also γ) intersects G at least $\rho(G, \Sigma)/2$ times.

Now let C be orientation reversing and let $C' = x_1 \cdots x_k x'_1 \cdots x'_k x_1$. If $x' \notin \gamma$ and γ meets C' at most once, then γ is also a nontrivial simple closed curve in Σ , and so γ intersects G'' at least $\rho(G, \Sigma)$ times. So we may assume that γ intersects C' at least twice. Hence γ has a subcurve P intersecting C' only at its ends, say y and z , such that $P + yx'z$ is nontrivial in Σ' . If $\{y, z\} = \{x_i, x'_i\}$, then P is a nontrivial simple closed curve in Σ , and so γ intersects G'' at least $\rho(G, \Sigma)/2$ times. Hence we may assume that $\{y, z\} \neq \{x_i, x'_i\}$. We may extend P to meet Γ , and let Q and R be the two subcurves of Γ between the two ends of P . Both $P \cup Q$ and $P \cup R$ are nontrivial in Σ ; for suppose that $P \cup Q$ is trivial in Σ , then it bounds a disc in Σ which implies that $P \cup Q$ bounds a disc in Σ' , contradicting the fact that $P \cup Q$ is homotopic to $P + yx'z$ in Σ' . Hence γ must have length at least $\rho(G, \Sigma)/2$. $\square$

Note that in the above lemma $\rho(G, \Sigma) \geq 3$ is sufficient if we do not ask C to be induced. We now reduce graphs embedded in a surface other than the sphere to a graph embedded in a disjoint union of spheres.

Theorem 4.3. *Let G be a connected graph embedded in a surface Σ (other than the sphere) with $eg(G, \Sigma) = g$ and $\rho(G, \Sigma) \geq 8(d+1)(2^g - 1)$. Then G can be reduced to a graph H embedded in a disjoint union S of spheres by cutting along a set of planarizing cycles $\{C_1, \dots, C_m\}$ (in this order) such that*

- (i) *each C_i is induced,*
- (ii) *for every integer k with $0 \leq k \leq d/2$ there is an induced cycle D_i^k (and D_i^k if C_i is orientation preserving) which bounds a closed disc in S containing C_i' (and C_i) such that for every vertex $z \in D_i^k$ (and $z \in D_i^k$) there is a simple curve P in S from z to C_i' (and C_i) with length equal to $d_{HS}(z, C_i') = k+1$ (and $d_{HS}(z, C_i) = k+1$) and $P \cap D_i^k = \{z\}$ (and $P \cap D_i^k = \{z\}$), and*
- (iii) *all D_i^k and D_i^k are disjoint, and for each integer k with $0 \leq k \leq d/2$, the closed disc bounded by D_i^k containing C_i' is disjoint from the closed disc bounded*

by D_i^k containing C_i , and both do not contain the disc bounded by $D_j^{k'}$ or D_j^k containing C_j' or C_j for any $j > i$.

Proof. We use induction on $eg(G, \Sigma) = g$. By Lemma 4.2, given a shortest non-trivial simple closed curve Γ in Σ , we find an induced noncontractible cycle C in G homotopic to Γ such that after cutting along C we have $\rho(G'', \Sigma') \geq \rho(G, \Sigma)/2$ (also for every vertex z on C , $d_{G\Sigma}(z, \Gamma \cap V(G)) \leq 2$).

First, we deal with the case that Σ' is the sphere. If C is orientation preserving, then let $C_1 = C$. By a similar argument as in the proof of Lemma 4.2, $d_{G'\Sigma'}(C, C') \geq \rho(G, \Sigma)/2 \geq 4(d+1)$. Let $D_1^0 = C$ and $D_1'^0 = C_1'$. Suppose we have constructed D_1^k for $0 \leq k \leq d/2 - 1$. Let F^k be the facial cycle of the block of $G' - D_1^k$ containing C' which is not a facial cycle of G' . If F^k is induced, then let $D_1^{k+1} = F^k$. Otherwise let uv be a chord of F^k . Now one of the two cycles in $F^k + uv$ containing uv separates C from C' , and we use such a cycle to replace F^k . Continuing this procedure we eventually get an induced cycle D_1^{k+1} . Similarly we can construct $D_1'^k$ for $0 \leq k \leq d/2$. Note that conditions (i) and (ii) are satisfied because of the construction, and condition (iii) is satisfied since $d_{G'\Sigma'}(C, C') \geq 4(d+1)$.

Now assume C is orientation reversing. In this case Σ is the projective plane. Let $x \in V(G)$. We first find a noncontractible cycle C_1 such that $d_{G\Sigma}(x, C_1) \geq 4d$. By Lemma 4.1, there are disjoint contractible cycles F_i for $1 \leq i \leq 4d+3$ such that F_i contains $x, F_1, \dots, F_{i-1}$ in its interior for $i \geq 2$. If C does not intersect the open disc bounded by F_{4d+3} , then $C_1 = C$. Otherwise, C contains a subpath P from x to y , where $x, y \in F_{4d+3}$, such that P and a subpath Q of F_{4d+3} between x and y form a noncontractible cycle C_1 . Clearly C_1 does not intersect the open disc bounded by F_{4d+3} . All possible chords of C_1 are from P to Q , and so we may assume that C_1 is induced (or else we use the same argument as in proof of Lemma 4.1 to get rid of those chords). Clearly $d_{G\Sigma}(x, C_1) \geq 4d$. Now after cutting along C_1 we obtain a graph G' embedded in the sphere Σ' with a cycle C_1' and $d_{G'\Sigma'}(x, C_1') \geq 4d$. Let $D_1'^0 = C_1'$. Suppose we have constructed $D_1'^k$ for $0 \leq k \leq d/2 - 1$. Let F'^k be the facial cycle of the block of $G' - D_1'^k$ containing x which is not a facial cycle of G' . Let $D_1'^{k+1} = F'^k$ if F'^k is induced. Otherwise let uv be a chord of F'^k . Then one of the cycles in $F'^k + uv$ containing uv separates C_1' from x , and we use such a cycle to replace F'^k . Continuing this procedure we will eventually get the desired induced cycle $D_1'^{k+1}$. Clearly conditions (i)-(iii) are satisfied.

We now assume that after cutting along C , G' is embedded in Σ' other than the sphere. By Lemma 4.2, $\rho(G'', \Sigma') \geq 4(d+1)(2^g - 1)$. By Lemma 4.1 we have the contractible induced cycles D'^k (and D^k if C is orientation preserving) in Σ' such that for every vertex $z \in D'^k \cup D^k$ there is a simple curve from z to x' (and to x) in the closed disc in Σ' bounded by D'^k (and D^k) of length $d_{G'\Sigma'}(z, x') = k+1$ (and $d_{G'\Sigma'}(z, x) = k+1$) meeting $D'^k \cup D^k$ exactly at z , where $D'^0 = C'$ (and $D^0 = C$) and $k \leq \rho(G'', \Sigma')/2 - 2$.

We claim that when C is orientation preserving the closed disc in Σ' bounded by D'^{d+1} containing C' is disjoint from the closed disc in Σ' bounded by D'^{d+1} containing C . First we show that in Σ' the closed disc bounded by D'^{d+1} containing C' does not contain C . Otherwise let C be contained in the interior of D'^k with k minimum ($k \leq d+1$). Then there is a simple curve P of length at most $2k+2$ in the interior of D'^k (in Σ') meeting C' only at its ends such that the two new closed curves in $P \cup C'$ contain C in their interiors (in Σ'). We may extend P a little to

meet Γ . Let Q and R be the two subcurves of Γ between the two ends of P . Then both $P \cup Q$ and $P \cup R$ are nontrivial in Σ . Since Γ is shortest in Σ , Γ would have length at most $2(2k+4) - 2 \leq 4d+6$, a contradiction. Similarly we can prove that the closed disc in Σ' bounded by D^{d+1} containing C does not contain C' . Hence if the closed disc in Σ' bounded by D'^{d+1} containing C' intersects the closed disc bounded by D^{d+1} containing C , then in Σ there is a simple curve from the left side of C to the right side of C of length at most $2d+3$. But this would imply that Γ has length at most $4d+6$, a contradiction.

Now let $E' = D'^{d+1}$ (and let $E = D^{d+1}$ if C is orientation preserving). We form a new graph G_0 embedded in Σ' by contracting the interior of E' (and the interior of E in the orientation preserving case) to a vertex y' (and y) and deleting the resulting loops and multiple edges.

We now verify that $\rho(G_0, \Sigma') \geq \rho(G'', \Sigma') - 4(d+1)$. Let γ be a nontrivial simple closed curve in Σ' . If γ does not contain any of y or y' , then γ intersects $V(G_0)$ only at vertices of $V(G'')$, and so γ has length at least $\rho(G'', \Sigma')$. If γ uses y' , then γ has a subcurve from $z_1 \in E'$ to $z_2 \in E'$ which only intersects $V(G_0)$ at y', z_1 and z_2 . We can replace this subcurve of γ by a simple curve in Σ' from z_1 to z_2 in the interior of E' (in G'') which intersects $V(G'')$ at most $2(d+1) + 1$ times. A similar modification is performed if γ contains y . Thus γ intersects $V(G_0)$ at least $\rho(G'', \Sigma') - 4(d+1)$ times. Hence $\rho(G_0, \Sigma') \geq 8(d+1)(2^{g-1} - 1)$.

We apply induction to G_0 which is embedded in the surface Σ' . We find a set of induced disjoint cycles $\{C_2, \dots, C_m\}$ and by cutting along these cycles in the order $C_2, \dots, C_m$ we reduce G_0 to a graph H_0 in a disjoint union S of spheres. We now obtain a graph H from H_0 by replacing y' and y by the interiors of D'^{d+1} and D^{d+1} in G' (in Σ'). We also modify the cycles $\{C_2, \dots, C_m\}$ to get the desired set of planarizing cycles in G . If y is used by some C_j ($2 \leq j \leq m$), then let uyv be a subpath of C_j , where $u, v \in V(C_j)$. Since $uy, vy \in E(G_0)$, there are disjoint paths from u and v to the interior of E (in Σ') which are internally disjoint from E . We can rewrite C_j by replacing uyv by the union of a path from u to $y_1, y_1 \dots y_r$, and a path from v to y_r , where $y_1 \dots y_r \subset G'$ is in the interior of E in Σ' and $d_{G' \Sigma'}(y_i, E) = 1$ for each i . We do the same when y' is used by C_j . Clearly H and $\{C_1, \dots, C_m\}$ are as desired. $\square$

5. HAMILTON CYCLES

We now prove the following result, establishing a conjecture of Thomassen [14].

Theorem 5.1. *Let G be a 5-connected triangulation of a surface Σ with $eg(G, \Sigma) = g$ and $\rho(G, \Sigma) \geq 96(2^g - 1)$. Then G has a Hamilton cycle.*

Proof. By Theorem 4.3 we can cut the surface along a set of planarizing cycles to get a new graph H embedded in a disjoint union of spheres with disjoint induced cycles C_i, C'_i, D_i^5 , and $D_i'^5$ for $i = 1, \dots, m$. Note that $d = 11$. Since G is a triangulation, all faces of H are triangles except possibly those bounded by C_i or C'_i , $i = 1, \dots, m$.

When C_i is orientation preserving, let H_i be the graph obtained from H by taking the union of the subgraph of H contained in the closed disc bounded by $D_i'^5$ containing C'_i and the subgraph of G contained in the closed disc bounded by D_i^5 containing C_i (note that these two subgraphs of H are disjoint), and identifying C_i and C'_i back to C_i . We view H_i as a plane graph with outercycle $D_i'^5$ and another disjoint facial cycle D_i^5 . Note that every simple curve in the plane from D_i^5 to

$D_i'^5$ must intersect H_i at least 11 times. Let M_i be the block of $H_i - (D_i^5 \cup D_i'^5)$ containing the cycle $C_i = C_i'$, and F_i and F_i' be the new facial cycles of M_i with F_i' being the outercycle.

When C_i is orientation reversing, let H_i be the subgraph of H contained in the closed disc bounded by $D_i'^5$ containing C_i' , and then identify C_i' back to C_i . We view H_i as a graph embedded in the projective plane with a facial cycle $D_i'^5$. Let M_i be the block of $H_i - D_i'^5$ containing C_i , and let F_i' be the facial cycle of M_i such that every $(M_i \cup D_i'^5)$ -bridge of H_i has at most one attachment on F_i' .

Since for every $z \in D_i^5 \cup D_i'^5$ there is a simple curve in Σ from z to C_i or C_i' intersecting G exactly 6 times, every $(M_i \cup D_i'^5 \cup D_i^5)$ -bridge has either one attachment on $D_i'^5 \cup D_i^5$ or two attachments on $D_i'^5$ or D_i^5 which are adjacent vertices.

Claim 1. Every $(M_i \cup D_i^5 \cup D_i'^5)$ -bridge of H_i is an edge.

Proof of Claim 1. Suppose that B is a $(M_i \cup D_i^5 \cup D_i'^5)$ -bridge of H_i which is not an edge. Since G is 5-connected, B must contain exactly one of C_i' or C_i . But this implies that there is a nontrivial simple closed curve in Σ (along B) intersecting G only at $a(B)$ (which has at most three vertices), a contradiction.

For $i = 1, 2, \dots, m$ and in this order, we replace with a vertex u_i' (or a vertex u_i) the edges and vertices contained in the open disc bounded by $D_i'^5$ (or bounded by D_i^5) containing C_i' (or C_i), and then add edges from u_i' (or u_i) to every vertex on $D_i'^5$ (or D_i^5) which is adjacent to at least two vertices on F_i' (or F_i). We use H' to denote the resulting graph. Note that H' needs not to be connected. Similarly we can obtain a graph M_i' from M_i by replacing by u_j' or u_j those open discs bounded $D_j'^5$ or D_j^5 containing C_j' or C_j which are contained in M_i , where $M_i' = M_i$ if M_i contains no other C_j' or C_j .

Note that the degrees of u_i' and u_i in H' or M_i' are sufficiently large (by the large representativity and the fact that G is a triangulation).

Claim 2. Every component of H' is 4-connected.

Proof of Claim 2. Suppose that a component of H' has a cutset S of size less than or equal to 3. By the choice of D_i^5 and $D_i'^5$, we may assume that $|S \cap D_i'^5| \geq 2$ (or equivalently, $|S \cap D_i^5| \geq 2$). Note that G is a triangulation. If $u_i' \in S$, then $D_i'^5$ has a chord, a contradiction. So $u_i' \notin S$. Let A be the component of $H' - S$ not containing u_i' . Then all vertices in $(A \cap D_i'^5) - S$ are adjacent to exactly one vertex of F_i' , say v , and so $S \cup \{v\}$ is a 4-cut of G , a contradiction. Thus each component of H' is 4-connected.

By Claim 2, each component of H' has a Hamilton cycle. We use T to denote the union of all those Hamilton cycles. Let the two vertices on T adjacent to u_i' (or u_i) be p_i' and q_i' (or p_i and q_i). By the choice of u_i' and u_i , we may assume that p_i' and q_i' are adjacent to x_i' and y_i' in F_i' , respectively, and that p_i and q_i are adjacent to x_i and y_i in F_i , respectively. Note that we may choose $x_i' \neq y_i'$ and $x_i \neq y_i$.

Claim 3. Every 3-cut of M_i' must have two vertices on F_i' or on F_i and each M_i' is 3-connected.

Proof of Claim 3. The first half of the claim follows from the same argument as in the proof of the above claim.

Suppose that S is a 2-cut of M'_i . We may assume that $S \subset F'_i$ (or equivalently, $S \subset F_i$). Let A be the component of $M'_i - S$ not containing C_i . Since $D_i'^4$ is induced and G is a triangulation, S cannot be contained in $D_i'^4$. Since every vertex of $D_i'^5$ has a simple curve in Σ to C_i intersecting G exactly 5 times, A is incident to at most two vertices on $D_i'^5$ which must be adjacent. Hence S together with the vertices of $D_i'^5$ incident with A forms a 4-cut, a contradiction. Therefore M'_i is 3-connected.

We now extend these disjoint paths $T - \{u_i, u'_i : i = 1, \dots, m\}$ to a Hamilton cycle of G by including those vertices in M'_i .

If C_m is orientation reversing, then we just use Theorem 3.8 to find a path P'_m in M'_m from x'_m to y'_m . Let $T_1 = (T - u'_m) \cup P'_m \cup \{p'_m x'_m, q'_m y'_m\}$. If C_m is orientation preserving and u_m and u'_m are in the same component of T , then we use Theorem 3.4 to find two disjoint paths P_m and P'_m with P_m from x_m to y_m and P'_m from x'_m to y'_m ; otherwise we use Theorem 3.6 to find two disjoint paths P_m and P'_m from $\{x_m, y_m\}$ to $\{x'_m, y'_m\}$. Let $T_1 = (T - \{u_m, u'_m\}) \cup P_m \cup P'_m \cup \{p_m x_m, q_m y_m, p'_m x'_m, q'_m y'_m\}$. By Claim 3, T_1 uses all vertices in M'_m . Suppose that we have included vertices in M'_j and the current T_{m-j+1} uses all vertices in M'_j . For $u'_i, u_i \in M'_j$, we define $x_i, x'_i, y_i, y'_i, p_i, q_i, p'_i, q'_i$ much the same as we did for those from T . We then in M'_{j-1} find two disjoint paths P_{j-1} and P'_{j-1} (or a path P'_{j-1} when C_j is orientation reversing) by Theorem 3.4 or Theorem 3.6 (or Theorem 3.8). By the above claim T_{m-j+2} uses all vertices of M'_{j-1} .

Note that conclusion (iii) of Theorem 4.3 guarantees that we can safely continue this procedure to find paths P'_i and/or P_i in M'_i . It is now easy to see that T_m is a Hamilton cycle of G . $\square$

To end this section, we point out that our technique fails for the general 5-connected graphs because we cannot find a way to get M'_i 3-connected, where the main problem is that a 2-cut may be contained in $D_i'^4$. It would be interesting to know if 5-connected graphs with sufficiently large representativity are hamiltonian.

6. 2-WALKS AND 3-WALKS

We end this paper by presenting an application of the generalized version of Thomassen's planarizing cycles. A *chain of blocks* is either a 2-connected graph or a connected graph in which every block contains at most two cutvertices and every cutvertex is contained in exactly two blocks. In a chain of blocks an *endblock* is a block containing at most one cutvertex. A *circuit graph* (G, C) is a 2-connected plane graph G with a facial cycle C such that for any 2-cut T of G , each T -bridge of G must contain a vertex of C . An *annulus graph* (G, C_1, C_2) is a 2-connected plane graph with two facial cycles C_1 and C_2 such that for each 2-cut T of G , every T -bridge of G contains a vertex of $C_1 \cup C_2$.

The following result is from [5] which was used by Gao and Richter to establish a conjecture of Jackson and Wormald.

Lemma 6.1. *Let (G, C) be a circuit graph and $x, y \in V(C)$. Then G has a 2-walk that visits each of x and y only once and traverses every edge of C exactly once.*

We need the following easy generalization of Lemma 6.1.

Lemma 6.2. *Let G be a plane graph which is a chain of blocks $G_1, \dots, G_n$ such that (G_i, C_i) is a circuit graph, where C_i is the outercycle of some G_i . Let $x \in V(C_1)$*

and $y \in V(C_n)$ such that neither x nor y is a cutvertex of G . Then G has a 2-walk which visits x and y exactly once.

It is not difficult to see that if a graph has a 2-walk, then it also has a spanning tree with maximum degree at most 3. Thomassen [14] showed that every triangulation of orientable genus g with no noncontractible cycle of length less than 2^{3g+4} contains a spanning tree of maximum degree at most 4. He also conjectured in [14] that the maximum degree can be lowered to 3 provided that the triangulation is 4-connected. This was verified by Ellingham and Gao [4] and they need 2^{3g+5} for the length of a shortest noncontractible cycle. In this section we prove the following stronger result.

Theorem 6.3. *Let G be a 4-connected graph embedded in a surface Σ with $eg(G, \Sigma) = g$ and $\rho(G, \Sigma) \geq 48(2^g - 1)$. Then G has a 2-walk.*

Proof. We use Theorem 4.3 and an idea similar to that in the proof of Theorem 5.1. We know that G has a set of planarizing cycles $C_1, \dots, C_m$ with D_i^2 and $D_i'^2$ as in Theorem 4.3. Note that $d = 5$. We now cut G along these cycles to obtain a graph H (embedded in a disjoint union of spheres) with disjoint induced cycles C_i' and $D_i'^2$ (and C_i and D_i^2 if C_i is orientation preserving). Let H_i , M_i , M_i' , F_i' and F_i be defined the same as in the proof of Theorem 5.1 (note that we use $D_i'^2$ and D_i^2 instead of $D_i'^5$ and D_i^5). Unlike the case in Theorem 5.1, M_i' may have 2-cuts contained in F_i' or in F_i .

As in the proof of Theorem 5.1, every $(M_i \cup D_i^2 \cup D_i'^2)$ -bridge of H_i is an edge. By a result on matchings in bipartite graphs (see [1] for example) and by the fact that G has very large representativity, we conclude that there is a very large matching A_i' (or A_i) between $D_i'^2$ and F_i' (or between D_i^2 and F_i).

We now construct new graphs H' and M_i' the same as in the proof of Theorem 5.1, and we use the same notation for vertices u_i' and u_i , where we join u_i' (and/or u_i) to the vertices in $D_i'^2 \cap V(A_i')$ (and/or $D_i^2 \cap V(A_i)$).

As in the proof of Theorem 5.1, we find a cycle T_m in G by finding disjoint cycles $T = T_0$ in H' using Lemma 2.1, and then finding paths P_i' and P_i in each M_i' using Lemma 3.3 or Lemma 3.7 (or Lemma 3.5 if u_i and u_i' are in different components of T_{m-i}). This can be done as every $u_i, u_i' \in H'$ is used by T and every $u_i, u_i' \in M_j'$ is used by $P_j' \cup P_j$. Clearly every T_m -bridge of G is either a T -bridge of H' containing a vertex of $D_i^2 \cup D_i'^2$ not as an attachment or a $(P_i' \cup P_i)$ -bridge of M_i' (or a P_i' -bridge of M_i') containing a vertex of $F_i' \cup F_i$ (or F_i') not as an attachment. We now extend T_m to a 2-walk of H by including vertices in those T -bridges of H' and those P_i' -bridges or $(P_i \cup P_i')$ -bridges of M_i' .

First we include those vertices in T -bridges of H' containing a vertex of $D_i'^2 \cup D_i^2$ not as an attachment. Without loss of generality we may assume that $B_1, \dots, B_k$ are those T -bridges of H' appearing on $D_i'^2$ in this order. Then by symmetry we may assume that $a(B_j) = \{a_j, b_j, c_j\}$, where $a_j D_i'^2 b_j \subset B_j$. Since G is 4-connected, $B_j - \{b_j, c_j\}$ is a chain of blocks with a_j in an endblock not as a cutvertex. By Lemma 6.2 we can find a 2-walk W_j in $B_j - \{b_j, c_j\}$ such that a_j is only visited once by W_j and we attach W_j to T_m . Similarly, we can include all vertices of T -bridges of H' containing a vertex of D_i^2 .

Now we consider those vertices in $(P_i \cup P_i')$ -bridges or P_i' -bridges of M_i' . Let $B'_1, \dots, B'_n$ be the $(P_i \cup P_i')$ -bridges of M_i' on F_i' in this order. Let $a(B'_j) = \{a'_j, b'_j, c'_j\}$ with $a'_j F_i' b'_j \subset B'_j$. Then since G is 4-connected, $B'_j - \{b'_j, c'_j\}$ is a chain

of blocks with a'_j in an endblock not as a cutvertex. Hence by Lemma 6.2 we can find a 2-walk W'_j of $B'_j - \{b'_j, c'_j\}$ such that a'_j is only visited once. We attach W'_j to T_m and we can similarly include vertices of those bridges of M'_i containing vertices of F_i .

It is now straightforward to verify that T_m together with all those W_j and W'_j forms a 2-walk of G . $\square$

A corollary of this result is that every 4-connected graph in a surface of orientable or nonorientable genus g with representativity at least $48(2^g - 1)$ has a spanning tree of maximum degree 3, which generalizes and improves a result of [4] (and so implies a conjecture of [14]).

Finally we deal with 3-walks in 3-connected graphs. For convenience we view a k -walk as an eulerian graph by doubling or tripling an edge if it is used twice or three times. Ellingham (personal communication) asked if a 3-connected graph in a surface with large representativity has a 3-walk? Here we answer this question in the affirmative.

Theorem 6.4. *Let G be a 3-connected graph embedded in a surface Σ with $eg(G, \Sigma) = g$ and $\rho(G, \Sigma) \geq 48(2^g - 1)$. Then G has a 3-walk.*

Sketch of a proof. By Theorem 4.3, G contains a set of planarizing cycles $C_1, \dots, C_m$ and cycles $D_i'^2$ and D_i^2 . Note that $d = 5$. Define H, H_i, M_i in the same way as in the proof of Theorem 5.1. Let H'' be the graph obtained from H by deleting all M_i . Then each component of H'' is a 2-connected plane graph, and every 2-cut of H'' is contained in some $D_i'^2$ or D_i^2 . Let M_i'' be obtained from M_i by deleting all M_j which is contained in M_i . Then M_i'' is a 2-connected plane graph, and every 2-cut of M_i'' is contained in some $D_j'^2$ or D_j^2 . We now prove the following statement.

Let K be a component of H'' and x be a vertex in K . Then K contains a 3-walk using x at most twice such that for each $D_i'^2 \subset K$ (respectively, each $D_i^2 \subset K$) the following hold.

- (i) if D_i^2 (respectively, $D_i'^2$) contains no 2-cut of K , then every vertex of D_i^2 (respectively, $D_i'^2$) is used at most twice, and
- (ii) for every maximal 2-cut $\{u, v\} \subset D_i^2$ (respectively, $\{u, v\} \subset D_i'^2$) and the $\{u, v\}$ -bridge B of K not containing any other D_j^2 or $D_j'^2$, the vertices in $B - \{u, v\}$ are used at most twice.

First, for every maximal 2-cut $\{u, v\}$ of K contained in some D_i^2 or $D_i'^2$, we replace with an edge uv the $\{u, v\}$ -bridge of K not containing D_j^2 or $D_j'^2$ for any $j \neq i$. This reduces K to a 3-connected plane graph which, by Lemma 6.1, contains a 2-walk W .

Second, we extend W to include the vertices in W -bridges of K containing edges of D_i^2 . Let $B_1, B_2, \dots, B_n$ be the W -bridges of K appearing on D_i^2 in this order. Let $a(B_j) = \{a_j, b_j\}$ with $a_j D_i^2 b_j \subset B_j$. We may assume, by symmetry, that $x \notin \{a_1, \dots, a_n\}$. Since G is 3-connected, $B_j - b_j$ is a chain of blocks with a_j in an endblock not as a cutvertex.

If the edge $a_j b_j$ is not used by W , then by Lemma 6.2, $B_j - b_j$ contains a 2-walk W_j using a_j only once. If the edge $a_j b_j$ is used by W , then $B_j + a_j b_j$ is a circuit graph, where $a_j D_i^2 b_j + a_j b_j$ is the outercycle. Now by Lemma 6.1, $B_j + a_j b_j$ contains a 2-walk using a_j and b_j just once and using all edges of $a_j D_i^2 b_j + a_j b_j$, and let W_j be obtained from the 2-walk by deleting the edge $a_j b_j$.

Now the union of W and all W_j 's includes all vertices in W -bridges of K containing edges of D_i^2 . Similarly we can extend W to include all vertices in W -bridges of K containing an edge of $D_i'^2$. Hence we get the desired 3-walk of K as in the statement.

Since G is 3-connected, there is an edge from $x_i \in V(M_i)$ to $y_i \in D_i^2$ (or $y_i \in D_i'^2$) such that y_i is used at most twice by 3-walks in M_l for $l < i$ (where $M_0 = H''$). In a similar way, we can find 3-walks in M_i'' such that (i) and (ii) above hold with $x, D_j'^2, D_j^2$ being replaced by $x_i, D_j'^2 \subset M_i'', D_j^2 \subset M_i''$.

Now by combining the 3-walks in H'', M_i'' together with the edges $x_i y_i$ we produce a 3-walk in G . $\square$

We point out here that Theorem 6.4 also generalizes a result of Thomassen [14] on spanning trees in triangulations.

ACKNOWLEDGMENT

I am grateful to the referee for his valuable suggestions.

REFERENCES

1. J. A. Bondy and U. S. R. Murty, *Graph Theory with Applications*. American Elsevier, 1976. MR **54**:117
2. R. Brunet, M. N. Ellingham, Z. Gao, A. Metzlar, and R. B. Richter, Spanning planar subgraphs of graphs in the torus and Klein bottle. *J. Combin. Theory Ser. B* 65 (1965) 7–22. MR **96g**:05045
3. R. A. Duke, On the genus and connectivity of hamiltonian graphs. *Discrete Math.*, Vol. 2, No. 3 (1972) 199–206. MR **47**:3221
4. M. N. Ellingham and Z. Gao, Spanning trees in locally planar triangulations. *J. Combinat. Theory Ser. B*, 61 (1994) 178–198. MR **95m**:05080
5. Z. Gao and R. B. Richter, 2-Walks in circuit graphs. *J. Combin. Theory Ser. B* 62 (1994) 259–267. MR **96e**:05093
6. B. Grünbaum, Polytopes, graphs, and complexes. *Bull. Amer. Math. Soc.* 76 (1970) 1131–1201. MR **42**:959
7. J. C. Molluzzo, Problem. in “*Second International Conference on Combinatorial Mathematics*” (A. Gewirtz and L. V. Quintas, Eds.) pp. 556–567, The New York Academy of Sciences, New York, 1979. MR **80j**:05003
8. C. St. J. A. Nash-Williams, Unexplored and semi-explored territories in graph theory. *New Directions in Graph Theory*, Academic Press, New York (1973) 169–176. MR **52**:7944
9. M. D. Plummer, Problem in infinite and finite sets. *Coll. Math. Soc. J. Bolyai 10*, Vol. III (A. Hajnal, R. Rado and V. T. Sós, Eds.), North-Holland, Amsterdam (1975) 1549–1550. MR **50**:12526
10. R. Thomas and X. Yu, 4-Connected projective-planar graphs are hamiltonian. *J. Combinat. Theory Ser. B* 62 (1994) 114–132. MR **95j**:05133
11. R. Thomas and X. Yu, 5-connected toroidal graphs are hamiltonian. *J. Combinat. Theory Ser. B.*, in press.
12. C. Thomassen, A theorem on paths in planar graphs. *J. Graph Theory* 7 (1983) 169–176. MR **84i**:05075
13. C. Thomassen, Embeddings of graphs with no short noncontractible cycles. *J. Combinat. Theory Ser. B* 48 (1990) 155–177. MR **91b**:05069
14. C. Thomassen, Trees in triangulations. *J. Combinat. Theory Ser. B* 60 (1994) 56–62. MR **95e**:05039
15. C. Thomassen, 5-Coloring maps on surfaces. *J. Combinat. Theory Ser. B* 59 (1993) 89–105. MR **94h**:05031

- 16. W. T. Tutte, A theorem on planar graphs. *Trans. Amer. Math. Soc.* 82 (1956) 99–116. MR **18**:408e
- 17. H. Whitney, A theorem on graphs. *Ann. of Math.* 32 (1931) 378–390.

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GEORGIA 30332

Current address: Department of Mathematics, Vanderbilt University, Nashville, Tennessee 37240

E-mail address: `yu@math.vanderbilt.edu`