HECKE ALGEBRAS AND COHOMOTOPICAL MACKEY FUNCTORS

NORIIHIKO MINAMI

Dedicated to Professor Hiroshi Toda on his 70th birthday

ABSTRACT. In this paper, we define the concept of the cohomotopical Mackey functor, which is more general than the usual cohomological Mackey functor, and show that Hecke algebra techniques are applicable to cohomotopical Mackey functors. Our theory is valid for any (possibly infinite) discrete group. Some applications to topology are also given.

1. INTRODUCTION

Hecke algebras have played an important role in the study of the cohomology of groups, more precisely cohomology of subgroups of a fixed (possibly infinite) discrete group G [40, 7, 44]. Because of this, it was a natural hope to find similar applications to the study of more general Mackey functors. Unfortunately, Hecke algebras can act naturally only on the so-called cohomological G -functors [46] (since a G -functor and a Mackey functor are essentially the same, we call it a cohomological Mackey functor here). Therefore that hope seemed to be an impossible project, at first glance.

Now the purpose of this paper is to realize this hope in the spirit of homotopical algebra [38, 42]. We define the notion of the cohomotopical Mackey functor (Definition 3.8), which generalizes the notion of the cohomological Mackey functor. Its typical example is the hypercohomology of G with coefficients in a G -spectrum X in the sense of [42], which is a natural generalization of the cohomology of the group from the setting of homological algebra to homotopical algebra [38]. (This situation motivated us to use the terminology “cohomotopical Mackey functor.”) Then the purpose of this paper is to show that Hecke algebra techniques are applicable to the cohomotopical Mackey functors also.

More precisely, we fix a commutative ring k and focus our attention on the Hurewicz functor (Definition 2.3)

$$H_k : (\mathcal{M}_k)_{\mathcal{F}}^{\wedge} \rightarrow \mathcal{H}_k,$$

where $(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}$ is the cohomotopical Mackey category (Definition 2.35) and $\mathcal{H}_k$ is the Hecke category (Definition 2.3). These functors emerge as the cohomotopical Mackey functor is defined to be a k -additive functor from $(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}$ and the cohomological Mackey functor is nothing but a k -additive functor from $\mathcal{H}_k$ (which is Yoshida’s theorem [46]).

Received by the editors April 8, 1997.

1991 *Mathematics Subject Classification*. Primary 55R35, 55P42, 55N91, 19A22, 18G55; Secondary 20C11, 57S17, 55R10.

Key words and phrases. Mackey functor, stable homotopy theory, classifying spaces.

Our philosophy here is to apply the homological algebra technique (Hecke algebra or Hecke category) to the (stable) homotopical algebra situation (cohomotopical Mackey functors) through the Hurewicz map (Hurewicz functor). Typical questions we would like to consider are: *If two objects are equivalent in $\mathcal{H}_k$, are they equivalent in $(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}$? If there is an idempotent decomposition in $\mathcal{H}_k$, is there a corresponding idempotent decomposition in $(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}$?* Our main results in §4 answer these questions affirmatively.

This paper is organized as follows. In §2, we define and investigate basic properties of underlying categories like the Mackey category, the Hecke category, and the cohomotopical Mackey category. In §3, we define the Mackey functor and other related functors like the cohomotopical Mackey functor. In §4, we state and prove our main results. For §5 and §6, we restrict ourselves to the case G finite and $k = \mathbb{Z}_p^{\wedge}$. In §5, we recall a beautiful identity in the Hecke category by Webb [44] and discuss some applications. As a corollary we get a combinatorial formula involving Möbius functions for cohomotopical Mackey functors over $\mathbb{Z}_p^{\wedge}$. In §6, we give applications to the topology. Here we consider the p -completed stable homotopy type of quotient spaces through free G -action, especially classifying spaces. We also mention the case of general (not necessarily free) G -action. In this case, even though there is no spectra-level combinatorial formula, we do have one at the level of cohomology.

We should emphasize that the philosophy of this paper comes from the homotopy theory. First, our main theorem can be interpreted as a representation theoretical analogue of Nishida's nilpotency theorem on the stable homotopy group of the sphere [36]. Second, we can give a topological proof, which makes use of the Segal conjecture [6], for a special case (i.e. G finite and $k = \mathbb{Z}_p^{\wedge}$) of the main theorem. Third, the whole project was motivated by the stable splitting of classifying spaces business. Following a conjecture of Priddy, we first proved the corresponding result for classifying spaces (Theorem 6.6), which originally led the author to conjecture our main results in §4.

If the reader is only interested in such applications to stable splittings of classifying spaces of finite groups (e.g. the Minami-Webb formula), then the reader with a prior knowledge of Yoshida's theorem [46] can jump straight to Lemma 6.8 and avoid the cohomotopical development. Yoshida's theorem [46] itself is reproved in this paper as Theorem 3.7, using our theory.

The original version of this paper was first distributed to experts, including Priddy and Webb, in 1988. (This old version was once again distributed in 1990 as MSRI preprint series 00425-91 [32].) During this period, there has been some progress in the subject [27, 28, 41, 20, 26, 21]. Such updated progress has also occurred in the current version of this paper. First, we have formulated things so that G could be an arbitrary (possibly infinite) discrete group. Second, we do not need the Segal conjecture for our proof of the main theorem, unlike the original version [32]. However, we have also inserted a significantly simplified version of this topological proof based on the Segal conjecture (for the case G finite and $k = \mathbb{Z}_p^{\wedge}$) in the current paper, because we believe it will give the reader better insight about analogies between algebra and topology.

The author's debt to Professor Stewart Priddy can never be overestimated. Not only did Professor Priddy motivate this work, but also he carefully read a preliminary version of this paper and suggested some improvements. Had it not been for his help, this paper would not have appeared. Also, the author would like

to thank Professor Haynes Miller for suggesting the use of the Mackey category (Definition 2.1), which turned out to be already given by Linder [25].

2. UNDERLYING CATEGORIES

In this section, we define basic underlying categories and discuss their properties, which are needed for our applications. Most of the proofs are straightforward.

Throughout this paper, G is a (possibly infinite) discrete group, and k is a commutative ring, unless otherwise stated.

Definition 2.1. Let $\mathcal{M}$ be the *Mackey category* defined as follows:

$$\text{Ob}_{\mathcal{M}} = \{\text{finite } G\text{-sets}\},$$

where a finite G -set is a finite set with a G -action.

To define the morphisms, we first consider the set of equivalence classes of the diagrams of the form

$$\begin{array}{ccc} & X & \\ f \swarrow & & \searrow g \\ S & & T, \end{array}$$

where X is a finite G -set, and f_S and f_T are G -maps. Two such objects are set to be equivalent:

$$\begin{array}{ccc} & X & \\ f_1 \swarrow & & \searrow g_1 \\ S & & T \end{array} \simeq \begin{array}{ccc} & Y & \\ f_2 \swarrow & & \searrow g_2 \\ S & & T, \end{array}$$

if and only if there is an isomorphism $h : X \rightarrow Y$ of G -sets such that $f_2 \circ h = f_1$ and $g_2 \circ h = g_1$. Then the monoid structure is given on representative elements by

$$\left[\begin{array}{ccc} & X_1 & \\ f_1 \swarrow & & \searrow g_1 \\ S & & T \end{array} \right] + \left[\begin{array}{ccc} & X_2 & \\ f_2 \swarrow & & \searrow g_2 \\ S & & T \end{array} \right] = \left[\begin{array}{ccc} & X_1 \amalg X_2 & \\ f_1 \amalg f_2 \swarrow & & \searrow g_1 \amalg g_2 \\ S & & T \end{array} \right].$$

Now $\text{Mor}_{\mathcal{M}}(S, T)$ is defined to be the group completion of this monoid. For $T = S$, the identity $I_S \in \text{Mor}_{\mathcal{M}}(S, S)$ is defined by

$$\left[\begin{array}{ccc} & S & \\ 1_S \swarrow & & \searrow 1_S \\ S & & S. \end{array} \right]$$

Furthermore, the composition of

$$\left[\begin{array}{ccc} & X & \\ f \swarrow & & \searrow g \\ S & & T \end{array} \right] \in \text{Mor}_{\mathcal{M}}(S, T) \quad \text{and} \quad \left[\begin{array}{ccc} & Y & \\ k \swarrow & & \searrow l \\ T & & U \end{array} \right] \in \text{Mor}_{\mathcal{M}}(T, U)$$

is given on representative elements by the pullback diagram of G -sets:

$$\left[\begin{array}{ccccc} & & Z & & \\ & s \swarrow & & \searrow t & \\ & X & & Y & \\ f \swarrow & & g \searrow & k \swarrow & \searrow l \\ S & & T & & U \end{array} \right] \in \text{Mor}_{\mathcal{M}}(S, U).$$

Clearly, this is uniquely extended to a bilinear map

$$\text{Mor}_{\mathcal{M}}(S, T) \otimes \text{Mor}_{\mathcal{M}}(T, U) \rightarrow \text{Mor}_{\mathcal{M}}(S, U),$$

which defines the composition.

For any commutative ring k , $\mathcal{M}_k$, the *Mackey category over k* , is defined by

$$\text{Ob}_{\mathcal{M}_k} = \text{Ob}_{\mathcal{M}},$$

$$\text{Mor}_{\mathcal{M}_k}(S, T) = \text{Mor}_{\mathcal{M}}(S, T) \otimes_{\mathbb{Z}} k.$$

In particular, $\mathcal{M}_{\mathbb{Z}} = \mathcal{M}$.

Definition 2.2. Given a sequence of finite index subgroups $K \subseteq H \subseteq G$, we define

$$\text{Res}_K^H \in \text{Mor}_{\mathcal{M}_k}(G/K, G/H),$$

$$\text{Ind}_K^H \in \text{Mor}_{\mathcal{M}_k}(G/H, G/K),$$

$$c_g \in \text{Mor}_{\mathcal{M}_k}(G/K^g, G/K)$$

by

$$\begin{aligned} \text{Res}_K^H &= \left[\begin{array}{ccc} & G/K & \\ 1_{G/K} \swarrow & & \searrow p \\ G/K & & G/H \end{array} \right] \in \text{Mor}_{\mathcal{M}_k}(G/K, G/H), \\ \text{Ind}_K^H &= \left[\begin{array}{ccc} & G/K & \\ p \swarrow & & \searrow 1_{G/K} \\ G/H & & G/K \end{array} \right] \in \text{Mor}_{\mathcal{M}_k}(G/H, G/K), \\ c_g &= \left[\begin{array}{ccc} & G/K & \\ r_g \swarrow & & \searrow 1_{G/K} \\ G/K^g & & G/K \end{array} \right] \in \text{Mor}_{\mathcal{M}_k}(G/K^g, G/K), \end{aligned}$$

where $p : G/K \rightarrow G/H$ is the canonical projection and $r_g : G/K \rightarrow G/K^g$ is given by the right multiplication by g , $hK \mapsto hKg = hgK^g$.

Definition 2.3. For any commutative ring k , we define $\mathcal{H}_k$, the *Hecke category over k* , as the category of permutation G -modules which are finitely generated over k :

$$\text{Ob}_{\mathcal{H}_k} = \{\text{finite } G\text{-sets}\},$$

$$\text{Mor}_{\mathcal{H}_k}(S, T) = \text{Hom}_{kG}(kS, kT).$$

We define the *Hurewicz functor* $H_k : \mathcal{M}_k \rightarrow \mathcal{H}_k$ as follows: On objects, H_k is the identity, and on morphisms, H_k sends

$$\left[\begin{array}{ccc} & X & \\ f \swarrow & & \searrow g \\ S & & T \end{array} \right] \in \text{Mor}_{\mathcal{M}_k}(S, T)$$

to

$$\left(s \mapsto \sum_{x \in f^{-1}(s)} g(x) \right) \in \text{Hom}_{kG}(kS, kT).$$

Remark 2.4. Suppose G is finite. Then the Mackey category $\mathcal{M}$ is nothing but the full subcategory of the equivariant stable homotopy category $\bar{h}GSU$ whose objects are the suspension spectra $\Sigma^\infty S_+$, where S is a finite G -set (see [24, v.9] for more detail). This point of view makes the well-definedness of the composition of morphisms in Definition 2.1 trivial.

This point of view also offers another interpretation of the Hurewicz functor $H : \mathcal{M} \rightarrow \mathcal{H}$ for G finite. Actually, $H : \text{Mor}_{\mathcal{M}}(S, T) \rightarrow \text{Mor}_{\mathcal{H}}(S, T)$ is nothing but the evaluation by the 0-dimensional non-equivariant ordinary reduced homology:

$$\{\Sigma^\infty S_+, \Sigma^\infty T_+\}_G \rightarrow \text{Hom}_G(\tilde{H}_0(\Sigma^\infty S_+), \tilde{H}_0(\Sigma^\infty T_+)) = \text{Hom}_{\mathbb{Z}G}(\mathbb{Z}S, \mathbb{Z}T).$$

This is essentially the treatment in [43].

Lemma 2.5. *Suppose $S = \coprod_i S_i$ and $T = \coprod_j T_j$ as finite G -sets. Then we have the following commutative diagram with canonical isomorphisms as horizontal arrows:*

$$\begin{array}{ccc} \text{Mor}_{\mathcal{M}_k}(S, T) & \xrightarrow{\cong} & \bigoplus_{i,j} \text{Mor}_{\mathcal{M}_k}(S_i, T_j) \\ H_k \downarrow & & \bigoplus_{i,j} H_k \downarrow \\ \text{Mor}_{\mathcal{H}_k}(S, T) & \xrightarrow{\cong} & \bigoplus_{i,j} \text{Mor}_{\mathcal{H}_k}(S_i, T_j). \quad \square \end{array}$$

When G is finite, the upper isomorphism in the following lemma may be interpreted as the G -self-duality theorem of Wirthmüller [45] through the equivariant stable homotopy theoretical interpretation in Remark 2.4.

Lemma 2.6. *For any finite G -sets S , T , and U , we have the following commutative diagram with canonical isomorphisms as horizontal arrows:*

$$\begin{array}{ccc} \text{Mor}_{\mathcal{M}_k}(S, T \times U) & \xrightarrow{\cong} & \text{Mor}_{\mathcal{M}_k}(S \times T, U) \\ H_k \downarrow & & H_k \downarrow \\ \text{Mor}_{\mathcal{H}_k}(S, T \times U) & \xrightarrow{\cong} & \text{Mor}_{\mathcal{H}_k}(S \times T, U). \end{array}$$

Proof. The upper horizontal isomorphism is given by

$$\left[\begin{array}{ccc} & X & \\ f \swarrow & & \searrow g \times k \\ S & & T \times U \end{array} \right] \mapsto \left[\begin{array}{ccc} & X & \\ f \times g \swarrow & & \searrow k \\ S \times T & & U \end{array} \right],$$

and the lower horizontal isomorphism is given by

$$h \in \text{Hom}_{kG}(kS, kT \times U) \\ \mapsto (\rho \otimes 1_{kU}) \circ (1_{kT} \otimes \epsilon) \circ (h \otimes 1_{kT}) \in \text{Hom}_{kG}(kS \times T, kU),$$

where $\rho : kT \times T \rightarrow k$ is characterized by $\rho(t, t) = 1$ and $\rho(t, t') = 0$ if $t \neq t'$, and $\epsilon : kU \times T \rightarrow kT \times U$ is induced by the interchange of U and T .

To check the commutativity of this diagram, notice that the middle square in

$$\left[\begin{array}{ccccc} & & X & & \\ & 1_X \times g \swarrow & & \searrow g \times h & \\ X \times T & & & & T \times U \\ \swarrow f \times 1_T & & & & \searrow p_U \\ S \times T & & (g \times h) \times 1_T \swarrow & \nwarrow (1_T \times \epsilon) \circ (\Delta_T \times 1_U) & U \end{array} \right] \in \text{Mor}_{\mathcal{M}}(S \times T, U)$$

is a pullback diagram, where $\delta_T : T \rightarrow T \times T$ is a diagonal map, $\epsilon : U \times T \rightarrow T \times U$ is the interchange of U and T , and p_U is the projection onto U . $\square$

Definition 2.7. Given finite G -sets S, T, U and V , the exterior pairing is defined by

$$\mu : \text{Mor}_{\mathcal{M}_k}(S, T) \otimes_k \text{Mor}_{\mathcal{M}_k}(U, V) \rightarrow \text{Mor}_{\mathcal{M}_k}(S \times U, T \times V), \\ \left[\begin{array}{ccc} & X & \\ f \swarrow & & \searrow g \\ S & & T \end{array} \right] \otimes \left[\begin{array}{ccc} & Y & \\ k \swarrow & & \searrow l \\ U & & V \end{array} \right] \mapsto \left[\begin{array}{ccc} & X \times Y & \\ f \times k \swarrow & & \searrow g \times l \\ S \times U & & T \times V \end{array} \right].$$

Definition 2.8. Let $f : G_1 \rightarrow G_2$ be a group homomorphism, and let $\mathcal{M}_k(G_1)$ and $\mathcal{M}_k(G_2)$ be Mackey categories of G_1 and G_2 , respectively. Then we may define a functor

$$f^* : \mathcal{M}_k(G_2) \rightarrow \mathcal{M}_k(G_1)$$

by regarding a finite G_2 -set and a diagram of finite G_2 -sets as a finite G_1 set and a diagram of finite G_1 -sets, respectively, through $f : G_1 \rightarrow G_2$.

Similarly, $f : G_1 \rightarrow G_2$ induces a functor

$$f^* : \mathcal{H}_k(G_2) \rightarrow \mathcal{H}_k(G_1)$$

between the corresponding Hecke categories by regarding a permutation kG_2 -module (which is finitely generated over k) as a finitely generated permutation kG_1 -module through $f : G_1 \rightarrow G_2$. Clearly, we have the following commutative diagram:

$$\begin{array}{ccc} \mathcal{M}_k(G_2) & \xrightarrow{f^*} & \mathcal{M}_k(G_1) \\ H_k \downarrow & & \downarrow H_k \\ \mathcal{H}_k(G_2) & \xrightarrow{f^*} & \mathcal{H}_k(G_1). \end{array}$$

Proposition 2.9. Consider a system of the canonical quotient homomorphisms $\{\pi_N : G \rightarrow G/N\}$, where N runs over finite index normal subgroups of G . Then we

have the following commutative diagram and isomorphisms of categories:

$$\begin{array}{ccc} \varinjlim_{N \triangleleft G; \text{finite index}} \mathcal{M}_k(G/N) & \xrightarrow[\cong]{\varinjlim_{N \triangleleft G} \pi_N} & \mathcal{M}_k(G) \\ \varinjlim_{N \triangleleft G} H_k \downarrow & & H_k \downarrow \\ \varinjlim_{N \triangleleft G; \text{finite index}} \mathcal{H}_k(G/N) & \xrightarrow[\cong]{\varinjlim_{N \triangleleft G} \pi_N} & \mathcal{H}_k(G). \end{array}$$

In particular, given finite G -sets S and T ,

$$\begin{array}{ccc} \varinjlim_{N \triangleleft G; \text{finite index}} \text{Mor}_{\mathcal{M}_k(G/N)}(S, T) & \xrightarrow[\cong]{\varinjlim_{N \triangleleft G} \pi_N} & \text{Mor}_{\mathcal{M}_k(G)}(S, T) \\ \varinjlim_{N \triangleleft G} H_k \downarrow & & H_k \downarrow \\ \text{Mor}_{\mathcal{H}_k(G/N_0)}(S, T) & \xrightarrow[\cong]{\pi_{N_0}} & \text{Mor}_{\mathcal{H}_k(G)}(S, T), \end{array}$$

where $\varinjlim_{N \triangleleft G; \text{finite index}}$ runs over those finite index normal subgroups N which are contained in the intersection of all the isotropy subgroups of S and T (this condition enables us to view S and T as finite G/N -sets), and N_0 is any one such normal subgroup.

Proof. Everything is easy to see, except the isomorphism

$$\text{Mor}_{\mathcal{H}_k(G/N_0)}(S, T) \xrightarrow[\cong]{\pi_{N_0}} \text{Mor}_{\mathcal{H}_k(G)}(S, T).$$

But this easily follows from Lemma 2.5 and the fact that

$$\text{Mor}_{\mathcal{H}_k(G)}(G/H, G/K) = \text{Hom}_{kG}(kG/H, kG/K) \cong kH \backslash G/K. \quad \square$$

Remark 2.10. Unlike the Hecke category, we usually cannot find a finite index normal subgroup N_0 such that $\text{Mor}_{\mathcal{M}_k(G/N_0)}(S, T) \xrightarrow[\cong]{\pi_{N_0}} \text{Mor}_{\mathcal{M}_k(G)}(S, T)$. A simplest example is $G = \mathbb{Z}$, $S = T = \text{pt}$.

Definition 2.11. Let H be a finite-index subgroup of G . Let $\mathcal{M}_k(G)$ and $\mathcal{M}_k(H)$ be the Mackey categories over k of G and H , respectively. Now define the functor

$$F_H^G : \mathcal{M}_k(H) \rightarrow \mathcal{M}_k(G)$$

by

$$\begin{array}{l} \text{Ob } \mathcal{M}_k(H) \rightarrow \mathcal{M}_k(G), \\ S \mapsto G \times_H S, \\ \text{Mor}_{\mathcal{M}_k(H)}(S, T) \rightarrow \text{Mor}_{\mathcal{M}_k(G)}(G \times_H S, G \times_H T), \\ \left[\begin{array}{ccc} & X & \\ f \swarrow & & \searrow g \\ S & & T \end{array} \right] \mapsto \left[\begin{array}{ccc} & G \times_H X & \\ G \times_H f \swarrow & & \searrow G \times_H g \\ G \times_H S & & G \times_H T \end{array} \right]. \end{array}$$

Under the same condition, let $\mathcal{H}_k(G)$ and $\mathcal{H}_k(H)$ be the Hecke categories over k of G and H , respectively. Then, using the induced representation $M \mapsto kG \otimes_{kH} M$,

we may define the functor $F_H^G : \mathcal{H}_k(H) \rightarrow \mathcal{H}_k(G)$ so that the following diagram commutes:

$$\begin{array}{ccc} \mathcal{M}_k(H) & \xrightarrow{F_H^G} & \mathcal{M}_k(G) \\ H_k \downarrow & & \downarrow H_k \\ \mathcal{H}_k(H) & \xrightarrow{F_H^G} & \mathcal{H}_k(G). \end{array}$$

To show that $F_H^G : \mathcal{M}_k(H) \rightarrow \mathcal{M}_k(G)$ is really a functor, it suffices to establish

Proposition 2.12. *Suppose H is a subgroup of G , and*

$$\begin{array}{ccc} P & \xrightarrow{h_1} & Y \\ v_1 \downarrow & & \downarrow v_2 \\ X & \xrightarrow{h_2} & A \end{array}$$

is a pull-back diagram of H -sets. Then

$$\begin{array}{ccc} G \times_H P & \xrightarrow{G \times_H h_1} & G \times_H Y \\ G \times_H v_1 \downarrow & & \downarrow G \times_H v_2 \\ G \times_H X & \xrightarrow{G \times_H h_2} & G \times_H A \end{array}$$

is a pull-back diagram of G -sets.

However, this is an immediate consequence of the following easy lemma:

Lemma 2.13. *Suppose H is a subgroup of G , T is a G -set, S is a H -set, and*

$$f : T \rightarrow G \times_H S$$

is a G -map. Let $p : G \times_H S \rightarrow G/H$ be the obvious canonical projection. Then

1. $(p \circ f)^{-1}(H) \subseteq T$ *is an H -subset.*
2. *There is an isomorphism of G -sets:*

$$T \cong G \times_H ((p \circ f)^{-1}(H)).$$

3. *Let $f|_{(p \circ f)^{-1}(H)} : (p \circ f)^{-1}(H) \rightarrow S (= H \times_H S \subseteq G \times_H S)$ be the resulting H -map. Then*

$$f = G \times_H (f|_{(p \circ f)^{-1}(H)}).$$

Furthermore, up to the identification in 2, the decomposition of f in 3 is unique, i.e. the decomposition in 3 uniquely characterizes $f|_{(p \circ f)^{-1}(H)}$. $\square$

From this lemma, we also immediately get

Proposition 2.14. *Let H be a finite index subgroup of G . Then, for any finite H -set S and a finite G -set T , there is a canonical isomorphism*

$$\text{Mor}_{\mathcal{M}_k(G)}(G \times_H S, T) \cong \text{Mor}_{\mathcal{M}_k(H)}(S, T). \quad \square$$

Remark 2.15. Clearly, the above canonical isomorphism may be completed to the following commutative diagram with canonical horizontal isomorphisms:

$$\begin{array}{ccc} \mathrm{Mor}_{\mathcal{M}_k(G)}(G \times_H S, T) & \xrightarrow{\cong} & \mathrm{Mor}_{\mathcal{M}_k(H)}(S, T) \\ H_k \downarrow & & H_k \downarrow \\ \mathrm{Mor}_{\mathcal{H}_k(G)}(G \times_H S, T) & \xrightarrow{\cong} & \mathrm{Mor}_{\mathcal{H}_k(H)}(S, T). \end{array}$$

Definition 2.16. For any group G , its Burnside ring $A(G)$ is defined to be the Grothendieck ring of finite G -sets; its additive structure and multiplicative structures are induced from the disjoint union and the Cartesian product of finite G -sets, respectively. For any commutative ring k , we set $A(G)_k := A(G) \otimes_{\mathbb{Z}} k$.

We denote the product map by

$$\begin{aligned} \mu : A(G)_k \otimes_k A(G)_k &\rightarrow A(G)_k, \\ [X] \otimes_k [Y] &\mapsto [X \times Y]. \end{aligned}$$

Furthermore, when H is a subgroup of G and g is an element of G , we set $\mathrm{Res}_H^G : A(G) \rightarrow A(H)$, $\mathrm{Ind}_H^G : A(H) \rightarrow A(G)$ and $c_g : A(H) \rightarrow A(H^g)$ by

$$\begin{aligned} \mathrm{Res}_H^G : A(G) &\rightarrow A(H), \\ [X] &\mapsto [X], \\ \mathrm{Ind}_H^G : A(H) &\rightarrow A(G), \\ [Y] &\mapsto [G \times_H Y], \\ c_g : A(H) &\rightarrow A(H^g), \\ [Y] &\mapsto [g^{-1}H \times_H Y], \end{aligned}$$

where we must require that H is of finite index in G to define Ind_H^G .

Remark 2.17. If G is a compact Lie group, this definition of the Burnside ring does not coincide with tom Dieck's in general.

Definition 2.18. Given a finite G -set S , we define $A_G(S)$ to be the Grothendieck construction of finite G -sets over S . More precisely, consider diagrams of the form

$$f : T \rightarrow S,$$

where T is a finite G -set and f is a G -map. Two such diagrams $f_i : T_i \rightarrow S$ ($i = 1, 2$) are said to be equivalent if there is an isomorphism of finite G -sets $h : T_1 \rightarrow T_2$ such that $f_1 = f_2 \circ h$. Furthermore, the monoid structure is given by the disjoint union on representative elements

$$[g : U \rightarrow S] + [h : V \rightarrow S] = [g \coprod h : U \coprod V \rightarrow S].$$

Now $A_G(S)$ is defined to be the group completion of this monoid. (A_G was defined and denoted by Ω as the (Burnside) Green functor in [11, p.303].)

As usual, we set $A_G(S)_k := A_G(S) \otimes_{\mathbb{Z}} k$. $A_G(S)_k$ is equipped with an $A(G)_k$ -module structure by

$$\begin{aligned} A(G)_k \otimes_k A_G(S)_k &\rightarrow A_G(S), \\ [X] \otimes_k [f : T \rightarrow S] &\mapsto [f \circ p_T : X \times T \rightarrow S], \end{aligned}$$

where $p_T : X \times T \rightarrow T$ is the canonical projection.

Given a G -map $g : S_1 \rightarrow S_2$ between finite sets, its induced map g_* is defined by

$$\begin{aligned} g_* : A_G(S_1)_k &\rightarrow A_G(S_2)_k, \\ [f : T \rightarrow S_1] &\mapsto [g \circ f : T \rightarrow S_2]. \end{aligned}$$

Clearly, this is an $A(G)_k$ -module map.

Definition 2.19. Given finite G -sets S and T , we define the exterior pairing by

$$\begin{aligned} \mu : A_G(S)_k \otimes_k A_G(T)_k &\rightarrow A_G(S \times T)_k, \\ [g : U \rightarrow S] \otimes [h : V \rightarrow T] &\mapsto [g \times h : U \times V \rightarrow S \times T]. \end{aligned}$$

Clearly, μ is an $A(G)_k$ -bilinear map and factors through

$$A_G(S)_k \otimes_{A(G)_k} A_G(T)_k \rightarrow A_G(S \times T)_k.$$

Definition 2.20. For each finite G -set S , define a k -algebra map

$$\begin{aligned} \Delta_S : A(G)_k &\rightarrow \text{Mor}_{\mathcal{M}_k}(S, S), \\ [X] &\mapsto \left[\begin{array}{ccc} & S \times X & \\ p_S \swarrow & & \searrow p_S \\ S & & S \end{array} \right], \end{aligned}$$

where p_S is the projection onto the S factor.

Lemma 2.21. Let H be a finite index subgroup of G and let S be a finite H -set. Then we have the following commutative diagram of k -algebras and k -algebra maps:

$$\begin{array}{ccc} A(G)_k & \xrightarrow{\Delta_{G \times_H S}} & \text{Mor}_{\mathcal{M}_k(G)}(G \times_H S, G \times_H S) \\ \text{Res}_H^G \downarrow & & \uparrow F_H^G \\ A(H)_k & \xrightarrow{\Delta_S} & \text{Mor}_{\mathcal{M}_k(H)}(S, S). \end{array}$$

Proof. This easily follows from the following commutative diagram of G -sets and G maps:

$$\begin{array}{ccc} G \times_H (S \times X) & \xrightarrow[\cong]{f} & (G \times_H S) \times X \\ p_1 \downarrow & & \downarrow p_2 \\ G \times_H S & \xlongequal{\quad} & G \times_H S, \end{array}$$

where X and S are finite G -set and finite H -set, respectively, and

$$\begin{aligned} f : (g, (s, x)) &\mapsto ((g, s), gx), \\ p_1 : (g, (s, x)) &\mapsto (g, s), \\ p_2 : ((g, s), x) &\mapsto (g, s). \quad \square \end{aligned}$$

Lemma 2.22. For any finite G -sets S and T , the following three $A(G)_k$ -actions on $\text{Mor}_{\mathcal{M}_k}(S, T)$ are all well-defined and coincide:

$$\begin{aligned} (2.1) \quad & A(G)_k \otimes_k \text{Mor}_{\mathcal{M}_k}(S, T) \\ & \xrightarrow{\Delta_S \otimes_k \text{Mor}_{\mathcal{M}_k}(S, T)} \text{Mor}_{\mathcal{M}_k}(S, S) \otimes_k \text{Mor}_{\mathcal{M}_k}(S, T) \\ & \rightarrow \text{Mor}_{\mathcal{M}_k}(S, T), \end{aligned}$$

where the last map is the composition;

$$(2.2) \quad \begin{aligned} & \text{Mor}_{\mathcal{M}_k}(S, T) \otimes_k A(G)_k \\ & \xrightarrow{\text{Mor}_{\mathcal{M}_k}(S, T) \otimes_k \Delta_T} \text{Mor}_{\mathcal{M}_k}(S, T) \otimes_k \text{Mor}_{\mathcal{M}_k}(T, T) \\ & \rightarrow \text{Mor}_{\mathcal{M}_k}(S, T) \end{aligned}$$

where the last map is the composition;

$$(2.3) \quad A(G)_k \otimes_k \text{Mor}_{\mathcal{M}_k}(S, T) \rightarrow \text{Mor}_{\mathcal{M}_k}(S, T),$$

$$[Q] \otimes \left[\begin{array}{ccc} & X & \\ f \swarrow & & \searrow g \\ S & & T \end{array} \right] \mapsto \left[\begin{array}{ccc} & Q \times X & \\ f \circ p_X \swarrow & & \searrow g \circ p_X \\ S & & T \end{array} \right]$$

where p_X is the projection onto the X factor. $\square$

Corollary 2.23. *With respect to the $A(G)_k$ -module structure in Lemma 2.22, the composition of the Mackey category is an $A(G)_k$ -bilinear map and induces*

$$\text{Mor}_{\mathcal{M}_k}(S, T) \otimes_{A(G)_k} \text{Mor}_{\mathcal{M}_k}(T, U) \rightarrow \text{Mor}_{\mathcal{M}_k}(S, U),$$

for any finite G -sets S , T and U . $\square$

Lemma 2.24. *We have the following canonical natural isomorphisms of $A(G)_k$ -modules:*

1. $\text{Mor}_{\mathcal{M}_k}(S, T) \cong A_G(S \times T)_k$.
2. $A_G(\text{pt})_k \cong A(G)_k$.
3. For any finite G -sets S_i $i = 1, 2, \dots, n$, we have

$$A_G \left(\prod_{i=1}^n S_i \right)_k \cong \bigoplus_{i=1}^n A_G(S_i)_k$$

4. For any finite index subgroup H of G and finite H -set S ,

$$\begin{aligned} A_H(S)_k &\cong A_G(G \times_H S)_k, \\ [f : T \rightarrow S] &\mapsto [G \times_H f : G \times_H T \rightarrow G \times_H S]. \end{aligned}$$

5. For any finite G -set $S = \coprod_{i=1}^n G/H_i$, we have

$$A_G(S)_k \cong \bigoplus_{i=1}^n A(H_i)_k. \quad \square$$

Proposition 2.25. *For any finite G -sets S and T , let $S \times T \cong \coprod_{i=1}^n G/H_i$ be the decomposition into G -orbits. Then we have the following commutative diagram with canonical $A(G)_k$ -module isomorphisms as horizontal arrows:*

$$\begin{array}{ccc} \text{Mor}_{\mathcal{M}_k}(S, T) & \xrightarrow{\cong} & \bigoplus_i A(H_i)_k \\ H_k \downarrow & & \bigoplus_i \epsilon_i \downarrow \\ \text{Mor}_{\mathcal{H}_k}(S, T) & \xrightarrow{\cong} & \bigoplus_i k, \end{array}$$

where $\epsilon_i : A(H_i)_k \rightarrow k$ is the augmentation map, the $A(G)_k$ -module structure on the left hand side is given by any one of Lemma 2.22, and that of the right hand side is given through $\bigoplus_{i=1}^n \text{Res}_{H_i}^G$.

In particular, $H_k : \text{Mor}_{\mathcal{M}_k}(S, T) \rightarrow \text{Mor}_{\mathcal{H}_k}(S, T)$ is always onto.

Proof. The commutative diagram is obtained by

$$\begin{array}{ccccccc} \mathrm{Mor}_{\mathcal{M}_k}(S, T) & \xrightarrow{\cong} & \mathrm{Mor}_{\mathcal{M}_k}(S \times T, \mathrm{pt}) & \xrightarrow{\cong} & \bigoplus_i \mathrm{Mor}_{\mathcal{M}_k}(G/H_i, \mathrm{pt}) & \xrightarrow{\cong} & \bigoplus_i A(H_i) \\ H_k \downarrow & & H_k \downarrow & & \bigoplus_i H_k \downarrow & & \bigoplus_i \epsilon_i \downarrow \\ \mathrm{Mor}_{\mathcal{H}_k}(S, T) & \xrightarrow{\cong} & \mathrm{Mor}_{\mathcal{H}_k}(S \times T, \mathrm{pt}) & \xrightarrow{\cong} & \bigoplus_i \mathrm{Mor}_{\mathcal{H}_k}(G/H_i, \mathrm{pt}) & \xrightarrow{\cong} & \bigoplus_i k, \end{array}$$

where the commutativities of the left, middle and the right squares follow from Lemma 2.6, Lemma 2.5, and Remark 2.15, respectively. $\square$

Remark 2.26. A different way of looking at the embedding of each factor $A(H_i) \rightarrow \mathrm{Mor}_{\mathcal{M}_k}(S, T)$ in Proposition 2.25 is given as follows:

Fix each G -orbit $G/(H \cap K^g) \cong G \cdot (H \times g^{-1}K) \subset G/H \times G/K$. The corresponding factor $A(H \cap K^g)$ is embedded by the following map:

$$\begin{aligned} A(H \cap K^g) &\xrightarrow{\Delta_{\mathrm{pt}}} \mathrm{Mor}_{\mathcal{M}_k}(H \cap K^g, \mathrm{pt}, \mathrm{pt}) \xrightarrow{F_{H \cap K^g}^G} \mathrm{Mor}_{\mathcal{M}_k(G)}(G/(H \cap K^g), G/(H \cap K^g)) \\ &\xrightarrow{c_g \circ \mathrm{Ind}_{H \cap K^g}^{K^g} \circ - \circ \mathrm{Res}_{H \cap K^g}^H} \mathrm{Mor}_{\mathcal{M}_k(G)}(G/H, G/K), \end{aligned}$$

where $c_g \circ \mathrm{Ind}_{H \cap K^g}^{K^g} \circ - \circ \mathrm{Res}_{H \cap K^g}^H$ sends $f \in \mathrm{Mor}_{\mathcal{M}_k(G)}(G/(H \cap K^g), G/(H \cap K^g))$ to $c_g \circ \mathrm{Ind}_{H \cap K^g}^{K^g} \circ f \circ \mathrm{Res}_{H \cap K^g}^H$.

Explicitly, the embedding is given by

$$A(H \cap K^g) \hookrightarrow \mathrm{Mor}_{\mathcal{M}_k(G)}(G/H, G/K),$$

$$[X] \mapsto \left[\begin{array}{ccc} & G \times_{H \cap K^g} X & \\ p_1 \swarrow & & \searrow p_1 \\ G/(H \cap K^g) & & G/(H \cap K^g) \\ p_2 \swarrow & & \searrow rp \\ G/H & & G/K \end{array} \right],$$

where p_1 and p_2 are obvious canonical projections and $rp : G/(H \cap K^g) \rightarrow G/K$ is the composite of the canonical projection onto G/K^g and $r_{g^{-1}} : G/K^g \rightarrow G/K, hK^g \mapsto hg^{-1}K$, the right multiplication by g^{-1} .

Corollary 2.27. *If H is a finite-index subgroup of G and S and T are finite G -sets, suppose either $S = G \times_H S_0$ or $T = G \times_H T_0$ for some finite H -sets S_0 or T_0 . Then the canonical $A(G)_k$ -action on $\mathrm{Mor}_{\mathcal{M}_k(G)}(S, T)$ factors $\mathrm{Res}_H^G : A(G)_k \rightarrow A(H)_k$. $\square$*

Definition 2.28. For any finite index subgroup H of G and a nonnegative integer $t \geq 0$, define a decreasing filtration $\{\mathcal{F}^n A(H)_k\}_{n=0}^\infty$ on $A(H)_k$ by

$$\mathcal{F}^0 A(H)_k = A(H)_k,$$

$$\mathcal{F}^n A(H)_k = I(G)^n \cdot A(H)_k,$$

where $I(G)$ is the augmentation ideal of the Burnside ring $A(G)$. For any finite G -set $S = \coprod_i G/H_i$, define a decreasing filtration $\{\mathcal{F}^n A_G(S)_k\}_{n=0}^\infty$ on $A_G(S)_k$ by setting

$$\mathcal{F}^n A_G(S)_k = \bigoplus_i \mathcal{F}^n A(H_i)_k,$$

where we have used the identification $A_G(S)_k = \bigoplus_i A(H_i)_k$ (Lemma 2.24).

Similarly, for any G -sets S and T , we define a decreasing filtration

$$\{\mathcal{F}^n \mathrm{Mor}_{\mathcal{M}_k}(S, T)\}_{n=0}^\infty$$

using the filtration $\{\mathcal{F}^n A_G(S \times T)_k\}_{n=0}^\infty$ (see Lemma 2.24).

Definition 2.29. For any finite index subgroup H of G and a nonnegative integer $t \geq 0$, define a decreasing filtration $\{\mathcal{F}_t^n A(H)_k\}_{n=0}^\infty$ on $A(H)_k$ by induction on t :

$$\begin{aligned}\mathcal{F}_0^n A(H)_k &:= I(H)_k^n \\ \mathcal{F}_t^n A(H)_k &:= \text{the ideal generated by elements of the form}\end{aligned}$$

$$(\text{Ind}_{H_1}^H a_1) \cdot (\text{Ind}_{H_2}^H a_2) \cdots (\text{Ind}_{H_l}^H a_l),$$

where H_i is a finite index subgroup of H and $a_i \in \mathcal{F}_{t-1}^{n_i} A(H_i)_k$ ($1 \leq i \leq l$) with $n_1 + n_2 + \cdots + n_l \geq n$.

Assembling these, we define a new filtration $\{\mathcal{F}_\infty^n A(H)_k\}_{n=0}^\infty$ on $A(H)_k$ by

$$\mathcal{F}_\infty^n A(H)_k := \bigcup_{t=0}^\infty \mathcal{F}_t^n A(H)_k.$$

As special cases, we have

$$\begin{aligned}\mathcal{F}_\infty^0 A(H)_k &= \mathcal{F}_t^0 A(H)_k = \mathcal{F}_0^0 A(H)_k = A(H)_k, \\ \mathcal{F}_\infty^1 A(H)_k &= \mathcal{F}_t^1 A(H)_k = \mathcal{F}_0^1 A(H)_k = I(H)_k.\end{aligned}$$

Notice that this filtration $\{\mathcal{F}_\infty^n A(H)_k\}_{n=0}^\infty$ on $A(H)_k$ does not depend upon any particular finite index embedding in G at all, unlike the filtration $\{\mathcal{F}^n A(H)_k\}_{n=0}^\infty$.

For each finite G -set $S = \coprod_i G/H_i$, define a new decreasing filtration

$$\{\mathcal{F}_\infty^n A_G(S)_k\}_{n=0}^\infty$$

on $A_G(S)_k$ by setting

$$\mathcal{F}_\infty^n A_G(S)_k = \bigoplus_i \mathcal{F}_\infty^n A(H_i)_k,$$

where we have used the identification $A_G(S)_k = \bigoplus_i A(H_i)_k$ (Lemma 2.24).

Similarly, for each finite G -sets S and T , we define a new decreasing filtration

$$\{\mathcal{F}_\infty^n \text{Mor}_{\mathcal{M}_k}(S, T)\}_{n=0}^\infty$$

using the filtration $\{\mathcal{F}_\infty^n A_G(S \times T)_k\}_{n=0}^\infty$ (see Lemma 2.24).

Proposition 2.30. Suppose G is a finite group. For any subgroup H of G , define the completed Burnside rings $(A(H)_k)_{\mathcal{F}}^\wedge$ and $(A(H)_k)_{\mathcal{F}_\infty}^\wedge$ by

$$\begin{aligned}(A(H)_k)_{\mathcal{F}}^\wedge &:= \varprojlim_n A(H)_k / \mathcal{F}^n A(H)_k, \\ (A(H)_k)_{\mathcal{F}_\infty}^\wedge &:= \varprojlim_n A(H)_k / \mathcal{F}_\infty^n A(H)_k.\end{aligned}$$

Then the natural ring homomorphism $(A(H)_k)_{\mathcal{F}}^\wedge \rightarrow (A(H)_k)_{\mathcal{F}_\infty}^\wedge$ is a topological isomorphism.

Proof. By definition, it is clear that

$$\mathcal{F}^n A(H)_k \subseteq I(H)_k^n \subseteq \mathcal{F}_\infty^n A(H)_k.$$

Therefore, for any n we must find an appropriate N_H such that

$$(2.4) \quad \mathcal{F}_\infty^{N_H} A(H)_k \subseteq I(G)_k^n \cdot A(H)_k = \mathcal{F}^n A(H)_k$$

to prove the claim. We prove this by induction on H with respect to the inclusion.

First, (2.4) certainly holds for $H = \{e\}$ because $\mathcal{F}_\infty^n A(\{e\})_k = \mathcal{F}^n A(\{e\})_k = 0$ for any $n \geq 1$.

Now assume (2.4) is proved for all maximal subgroups H_i ($i = 1, \dots, m$) of H with an appropriate integer N_{H_i} . Since the $I(G)$ -adic topology and $I(H)$ -adic topology determine the same topology on $A(H)$ for any subgroup H of a finite group G (see [18]), we may choose M so that $I(H)_k^M \subseteq I(G)_k^n \cdot A(H)_k$. We claim that

$$N_H := M + \sum_{i=1}^m (M-1)(N_{H_i} - 1)$$

will do the job.

To see this, just notice that a general element in $\mathcal{F}_\infty^{N_H} A(H)_k$ is a linear combination of elements of the form

$$(2.5) \quad \left(\prod_{i=1}^m \left(\prod_{j=1}^{e_i} (\text{Ind}_{H_i}^H a_{i,j}) \right) \right) \cdot a,$$

where $e_i \geq 0$, $a_{i,j} \in \mathcal{F}_\infty^{n_{i,j}} A(H_i)_k$ with $n_{i,j} \geq 1$, and $a \in I(H)_k^e$ for some $e \geq 0$ such that

$$e + \sum_{i=1}^m \sum_{j=1}^{e_i} n_{i,j} \geq N_H = M + \sum_{i=1}^m (M-1)(N_{H_i} - 1).$$

This is because any Ind_K^H from a finite index proper subgroup $K \subsetneq H$ factors through $\text{Ind}_{H_i}^H$ for some finite-index maximal proper subgroup H_i of H . Notice that a contradiction would occur if we falsely assume all of the following:

$$\begin{aligned} e &\leq M-1, \quad e_i \leq M-1 \quad (i = 1, \dots, m), \\ n_{i,j} &\leq N_{H_i} - 1 \quad (i = 1, \dots, m, j = 1, \dots, e_i). \end{aligned}$$

Thus, either 1) $e \geq M$, or 2) $e_i \geq M$ for some i ($i = 1, \dots, m$), or 3) $n_{i,j} \geq N_{H_i}$ for some $i = 1, \dots, m$ and $j = 1, \dots, e_i$ occurs.

If 1) or 2) occurs, the element (2.5) clearly belongs to $I(H)_k^M \subseteq I(G)_k^n \cdot A(H)_k$.

If 3) occurs, by the inductive assumption, $a_{i,j} \in I(G)_k^n \cdot A(H_i)_k$, which implies $\text{Ind}_{H_i}^H a_{i,j} \in I(G)_k^n \cdot A(H)_k$ because $\text{Ind}_{H_i}^H$ is an $A(G)_k$ -module homomorphism.

In this way, we have verified that the element (2.5) always belongs to $I(G)_k^n \cdot A(H)_k$, which completes the proof of (2.4). $\square$

Lemma 2.31. *For both filtrations in Definition 2.28 and Definition 2.29, the conjugation $c_g : A(H)_k \rightarrow A(H^g)_k$, the induction $\text{Ind}_K^H : A(K)_k \rightarrow A(H)_k$, and the restriction $\text{Res}_K^H : A(H)_k \rightarrow A(K)_k$ are filtration preserving.*

Proof. c_g : This is clear for the both filtrations.

Ind_K^H : The claim for $\{\mathcal{F}^n\}_{n=0}^\infty$ follows from the projection formula, which claims Ind_K^H is an $A(H)_k$ -module homomorphism (and so an $A(G)_k$ -module homomorphism through Res_H^G). The claim for $\{\mathcal{F}_\infty^n\}_{n=0}^\infty$ follows from the definition; just notice that $\mathcal{F}_\infty^n A(K)_k = \bigcup_{t=0}^\infty \mathcal{F}_t^n A(K)_k$ and

$$\text{Ind}_K^H (\mathcal{F}_t^n A(K)_k) \subseteq \mathcal{F}_{t+1}^n A(H)_k \subseteq \mathcal{F}_\infty^n A(H)_k.$$

Res_K^H : The claim for $\{\mathcal{F}^n\}_{n=0}^\infty$ follows from the fact that Res_K^H is a ring homomorphism (and so an $A(G)_k$ -module homomorphism through Res_H^G). To prove the claim for $\{\mathcal{F}_\infty^n\}_{n=0}^\infty$, we show that

$$\text{Res}_K^H (\mathcal{F}_t^n A(H)_k) \subseteq \mathcal{F}_\infty^n A(K)_k$$

by induction on t . In fact, this is trivial for $t = 0$, as $\mathcal{F}_0^n A(H)_k = I(H)_k^n$. Now assume this is proved for $t - 1$. To show the claim for t , since Res_K^H is a ring homomorphism, it suffices to prove that

$$\text{Res}_K^H \text{Ind}_L^H(a) \in \mathcal{F}_\infty^m A(K)_k,$$

where L is a finite index subgroup of H and $a \in \mathcal{F}_{t-1}^m A(L)_k$ (see Definition 2.29). However, the double coset formula claims it is a linear combination of elements of the form

$$c_g \text{Ind}_{K^{g^{-1}} \cap L}^{K^{g^{-1}}} \text{Res}_{K^{g^{-1}} \cap L}^L(a),$$

where $\text{Res}_{K^{g^{-1}} \cap L}^L(a) \in \mathcal{F}_\infty^m A(K^{g^{-1}} \cap L)_k$ by the inductive assumption, and $\text{Ind}_{K^{g^{-1}} \cap L}^{K^{g^{-1}}}$ and c_g both preserve the filtration as was shown above. Thus the claim follows. $\square$

Lemma 2.32. *For both filtrations in Definition 2.28 and Definition 2.29, the product $\mu : A(H)_k \otimes_k A(H)_k \rightarrow A(H)_k$ respects the filtration, i.e. it induces*

$$\begin{aligned} \mathcal{F}^a A(H)_k \otimes \mathcal{F}^b A(H)_k &\rightarrow \mathcal{F}^{a+b} A(H)_k \\ \mathcal{F}_\infty^a A(H)_k \otimes \mathcal{F}_\infty^b A(H)_k &\rightarrow \mathcal{F}_\infty^{a+b} A(H)_k \end{aligned}$$

for any $a, b \geq 0$. $\square$

Proposition 2.33. *For both filtrations in Definition 2.28 and Definition 2.29, we have the followings:*

- (i) *All the isomorphisms in Lemma 2.24 are filtration preserving.*
- (ii) *For any G -map $f : S \rightarrow T$ of finite G -sets, the induced map $f_* : A_G(S)_k \rightarrow A_G(T)_k$ is filtration preserving.*
- (iii) *The exterior pairing $\mu : A_G(S)_k \otimes A_G(T)_k \rightarrow A_G(S \times T)_k$ defined in Definition 2.19 respects the filtration, i.e. it induces*

$$\begin{aligned} \mathcal{F}^a A_G(S)_k \otimes \mathcal{F}^b A_G(T)_k &\rightarrow \mathcal{F}^{a+b} A_G(S \times T)_k, \\ \mathcal{F}_\infty^a A_G(S)_k \otimes \mathcal{F}_\infty^b A_G(T)_k &\rightarrow \mathcal{F}_\infty^{a+b} A_G(S \times T)_k \end{aligned}$$

for any $a, b \geq 0$.

Proof. (i) is easy to see. To show (ii), we simply notice that the map $f_* : A_G(S)_k \rightarrow A_G(T)_k$, induced by a G -map of finite G -sets $f : S \rightarrow T$, is a linear combination of maps of the form $\text{Ind}_K^H : A(K)_k \rightarrow A(H)_k$ with respect to the decomposition in Lemma 2.24. So the claim follows from Lemma 2.31.

To show (iii), we may assume both S and T are transitive G -sets, say $S = G/H$, $T = G/K$ for some finite index subgroups H and K . Choose a transitive orbit $G/(H \cap K^g) \cong G \cdot (H \times g^{-1}K) \subseteq G/H \times G/K$ and let $\pi : A_G(G/H \times G/K)_k \rightarrow A_G(G/H \cap K^g)_k$ be the corresponding projection. Then, in view of Lemma 2.31 and Lemma 2.32, the claim will follow if we can show the composite

$$(2.6) \quad \begin{aligned} A(H)_k \otimes_k A(K)_k &\cong A_G(G/H)_k \otimes_k A_G(G/K)_k \\ &\xrightarrow{\mu} A_G(G/H \times G/K)_k \xrightarrow{\pi} A_G(G/(H \cap K^g))_k \cong A(H \cap K^g)_k \end{aligned}$$

is the same as

$$(2.7) \quad \begin{aligned} A(H)_k \otimes_k A(K)_k &\xrightarrow{A(H)_k \otimes_k c_g} A(H)_k \otimes_k A(K^g)_k \\ &\xrightarrow{\text{Res}_{H \cap K^g}^H \otimes_k \text{Res}_{H \cap K^g}^{K^g}} A(H \cap K^g)_k \otimes_k A(H \cap K^g)_k \xrightarrow{\mu} A(H \cap K^g)_k. \end{aligned}$$

Now, pick a representative element $[X] \otimes [Y] \in A(H)_k \otimes_k A(K)_k$, where X (resp. Y) is a finite H set (resp. K set). Then its image under the exterior pairing μ is

$$\left[(G \times_H X) \times (G \times_K Y) \xrightarrow{p_1 \times p_2} G/H \times G/K \right] \in A_G(G/H \times G/K)_k,$$

where p_i ($i = 1, 2$) are the canonical projection maps. To find out the π image of this element, consider the following commutative diagram of G -maps between G -sets:

$$\begin{array}{ccccc} (G \times_H X) \times (G \times_K Y) & \xrightarrow[\cong]{t_1} & G \times_H [(G \times_K Y) \times X] & \xleftarrow[\cong]{G \times_H i_1} & G \times_H [(Hg^{-1}K \times_K Y) \times X] \\ p_1 \times p_2 \downarrow & & G \times_H p_3 \downarrow & & G \times_H p_4 \downarrow \\ G/H \times G/K & \xrightarrow[\cong]{b} & G \times_H (G/K) & \xleftarrow[\cong]{G \times_H i_2} & G \times_H (Hg^{-1}K) \\ & & & & \cong \downarrow \\ & & & & G/(H \cap K^g), \end{array}$$

where i_1 and i_2 are canonical inclusions, p_1 and p_2 are canonical projections, and

$$\begin{aligned} t_1 : ((g_1, x), (g_2, y)) &\mapsto (g_1, ((g_1^{-1}g_2, y), x)), \\ b_1 : (g_1H, g_2K) &\mapsto (g_1, g_1^{-1}g_2K), \\ p_3 : ((g, y), x) &\mapsto gK, \\ p_4 : ((hg^{-1}k, y), x) &\mapsto hg^{-1}K. \end{aligned}$$

Notice that the desired π image is exhibited in this commutative diagram as the right upper vertical map, which is the G -extension, i.e. the image under the isomorphism $A_H(Hg^{-1}K)_k \cong A_G(G \times_H (Hg^{-1}K))_k$, of the left vertical map in the following commutative diagram of H -spaces and H -maps:

$$\begin{array}{ccc} (Hg^{-1}K \times_K Y) \times X & \xrightarrow[\cong]{t_2} & H \times_{H \cap K^g} ((g^{-1}K \times_K Y) \times X) \\ p_4 \downarrow & & \downarrow p_5 \\ Hg^{-1}K/K & \xrightarrow[\cong]{b_2} & H/(H \cap K^g), \end{array}$$

where p_5 is the canonical projection and

$$\begin{aligned} t_2 : ((hg^{-1}k, y), x) &\mapsto (h, (g^{-1}k, y, h^{-1}x)), \\ b_2 : hg^{-1}K &\mapsto h(H \cap K^g). \end{aligned}$$

This latter commutative diagram exhibits the desired π image as the H extension, i.e. the image under the isomorphism $A(H \cap K^g)_k \cong A_H(H/(H \cap K^g))_k \cong A_H(Hg^{-1}K)_k$ ($\cong A_G(G \times_H (Hg^{-1}K))_k$) of

$$[(g^{-1}K \times_K Y) \times X] \in A(H \cap K^g)_k.$$

However, $g^{-1}K \times_K Y$, regarded as a finite K^g -set (and so a finite $H \cap K^g$ -set by restricting the action), represents the image of $[Y] \in A(K)_k$ under the conjugation $c_g : A(K)_k \rightarrow A(K^g)_k$ by Definition 2.16. This implies (2.6) is the same as (2.7), which completes the proof. $\square$

Proposition 2.34. *For both filtrations in Definition 2.28 and Definition 2.29, the composition*

$$\mathrm{Mor}_{\mathcal{M}_k}(S, T) \otimes \mathrm{Mor}_{\mathcal{M}_k}(T, U) \rightarrow \mathrm{Mor}_{\mathcal{M}_k}(S, U)$$

respects the filtration, i.e. it induces

$$\begin{aligned} \mathcal{F}^a \mathrm{Mor}_{\mathcal{M}_k}(S, T) \otimes \mathcal{F}^b \mathrm{Mor}_{\mathcal{M}_k}(T, U) &\rightarrow \mathcal{F}^{a+b} \mathrm{Mor}_{\mathcal{M}_k}(S, U), \\ \mathcal{F}_\infty^a \mathrm{Mor}_{\mathcal{M}_k}(S, T) \otimes \mathcal{F}_\infty^b \mathrm{Mor}_{\mathcal{M}_k}(T, U) &\rightarrow \mathcal{F}_\infty^{a+b} \mathrm{Mor}_{\mathcal{M}_k}(S, U) \end{aligned}$$

for any $a, b \geq 0$.

Proof. The claim for the filtration $\{\mathcal{F}^n\}_{n=0}^\infty$ is clear from Corollary 2.23.

For the filtration $\{\mathcal{F}_\infty^n\}_{n=0}^\infty$ it suffices to check the claim when T is a transitive G -set, say G/H . Then it is easy to check the commutativity of the following diagram:

$$\begin{array}{ccc} \mathrm{Mor}_{\mathcal{M}_k}(S, G/H) \otimes \mathrm{Mor}_{\mathcal{M}_k}(G/H, U) & \xrightarrow{\mu} & \mathrm{Mor}_{\mathcal{M}_k}(S, U) \xleftarrow{\cong} A_G(S \times U)_k \\ \cong \downarrow & & \uparrow (\pi_{S \times U})^* \\ A_G(S \times G/H)_k \otimes A_G(G/H \times U)_k & & A_G(G/H \times (S \times U))_k \\ \cong \downarrow & & \uparrow \cong \\ A_G(G \times_H S)_k \otimes A_G(G \times_H U)_k & & A_G(G \times_H (S \times U))_k \\ \cong \downarrow & & \uparrow \cong \\ A_H(S)_k \otimes A_H(U)_k & \xrightarrow{\mu} & A_H(S \times U)_k, \end{array}$$

where $p_{S \times U} : G/H \times (S \times U) \rightarrow S \times U$ is the projection. In this diagram, the bottom μ respects the filtration as was shown in Proposition 2.33 (iii), and all the other maps are filtration preserving by Proposition 2.33 (i) (ii). Thus the claim follows. $\square$

Definition 2.35. For each $l \geq 1$, we define the reduced Mackey categories $\mathcal{M}_k/\mathcal{F}^l$ and $\mathcal{M}_k/\mathcal{F}_\infty^l$ by

$$\begin{aligned} \mathrm{Ob}_{\mathcal{M}_k/\mathcal{F}^l} &= \mathrm{Ob}_{\mathcal{M}_k/\mathcal{F}_\infty^l} = \{\text{finite } G\text{-sets}\}, \\ \mathrm{Mor}_{\mathcal{M}_k/\mathcal{F}^l}(S, T) &= \mathrm{Mor}_{\mathcal{M}_k}(S, T)/\mathcal{F}^l \mathrm{Mor}_{\mathcal{M}_k}(S, T), \\ \mathrm{Mor}_{\mathcal{M}_k/\mathcal{F}_\infty^l}(S, T) &= \mathrm{Mor}_{\mathcal{M}_k}(S, T)/\mathcal{F}_\infty^l \mathrm{Mor}_{\mathcal{M}_k}(S, T), \end{aligned}$$

where S and T are finite G -sets. Here the composition is induced by that of $\mathcal{M}_k$. Its well-definedness is guaranteed by Proposition 2.34.

Similarly, we define the completed Mackey categories $(\mathcal{M}_k)_{\mathcal{F}}^\wedge$ and $(\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge$ by

$$\begin{aligned} \mathrm{Ob}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge} &= \mathrm{Ob}_{(\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge} = \{\text{finite } G\text{-sets}\}, \\ \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge}(S, T) &= \varprojlim_l \mathrm{Mor}_{\mathcal{M}_k/\mathcal{F}^l}(S, T), \\ \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge}(S, T) &= \varprojlim_l \mathrm{Mor}_{\mathcal{M}_k/\mathcal{F}_\infty^l}(S, T). \end{aligned}$$

Notice that we have the following commutative diagram of categories and functors:

$$\begin{array}{ccccccc}
 (\mathcal{M}_k)_{\mathcal{F}}^{\wedge} & \longrightarrow & \mathcal{M}_k/\mathcal{F}^n & \longrightarrow & \mathcal{M}_k/\mathcal{F}^{n-1} & \longrightarrow & \mathcal{M}_k/\mathcal{F}^1 = \mathcal{M}_k/\mathcal{F} \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 (\mathcal{M}_k)_{\mathcal{F}_{\infty}}^{\wedge} & \longrightarrow & \mathcal{M}_k/\mathcal{F}_{\infty}^n & \longrightarrow & \mathcal{M}_k/\mathcal{F}_{\infty}^{n-1} & \longrightarrow & \mathcal{M}_k/\mathcal{F}_{\infty}^1 = \mathcal{M}_k/\mathcal{F}_{\infty}.
 \end{array}$$

We call $(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}$ the *cohomotopical Mackey category over k* . Note that

$$\mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}}(S, T) \cong \mathrm{Mor}_{\mathcal{M}_k}(S, T) \otimes_{A(G)_k} (A(G)_k)_{I(G)_k}^{\wedge},$$

when G is a finite group.

Corollary 2.36. *The Hurewicz functor $H_k : \mathcal{M}_k \rightarrow \mathcal{H}_k$ induces an isomorphism $\overline{H}_k : \mathcal{M}_k/\mathcal{F}_{\infty} \rightarrow \mathcal{H}_k$.*

Proof. For any finite G -sets S and T , we have the following commutative diagram from Proposition 2.25:

$$\begin{array}{ccc}
 \mathrm{Mor}_{\mathcal{M}_k}(S, T) & \xrightarrow{\cong} & \bigoplus_i A(H_i)_k \\
 H_k \downarrow & & \bigoplus_i \epsilon_i \downarrow \\
 \mathrm{Mor}_{\mathcal{H}_k}(S, T) & \xrightarrow{\cong} & \bigoplus_i k.
 \end{array}$$

On the other hand, from Definition 2.35 and Definition 2.29, we have the following commutative diagram:

$$\begin{array}{ccc}
 \mathrm{Mor}_{\mathcal{M}_k}(S, T) & \xrightarrow{\cong} & \bigoplus_i A(H_i)_k \\
 \downarrow & & \bigoplus_i \epsilon_i \downarrow \\
 \mathrm{Mor}_{\mathcal{M}_k}(S, T)/\mathcal{F}_{\infty} & & \bigoplus_i A(H_i)_k/I(H_i)_k \\
 \parallel & & \parallel \\
 \mathrm{Mor}_{\mathcal{M}_k/\mathcal{F}_{\infty}}(S, T) & \xrightarrow{\cong} & \bigoplus_i k
 \end{array}$$

From these commutative diagrams, we immediately find that H_k induces an isomorphism

$$\mathrm{Mor}_{\mathcal{M}_k/\mathcal{F}_{\infty}}(S, T) \xrightarrow{\cong} \mathrm{Mor}_{\mathcal{H}_k}(S, T),$$

which proves the claim. $\square$

Corollary 2.37. *If G is a finite group, then the natural map*

$$(\mathcal{M}_k)_{\mathcal{F}}^{\wedge} \rightarrow (\mathcal{M}_k)_{\mathcal{F}_{\infty}}^{\wedge}$$

is a topological isomorphism.

Proof. This is an immediate consequence of Proposition 2.30, in view of Definition 2.28, Definition 2.29 and Definition 2.35. $\square$

Our attention is now focused upon the Hurewicz functor

$$H_k : (\mathcal{M}_k)_{\mathcal{F}}^{\wedge} \rightarrow \mathcal{H}_k,$$

whose study is the main subject in §4.

3. FUNCTORS

Having established basic properties of the underlying categories in the previous section, we now define various Mackey functors as k -additive functors from those categories.

As in the previous section, G is a (possibly infinite) discrete group and k is a commutative ring, unless otherwise stated.

Definition 3.1. We call a category $\mathcal{C}$ an *additive category over k* if

1. For any objects A and B in $\mathcal{C}$, $\text{Mor}_{\mathcal{C}}(A, B)$ is a k -module.
2. For any objects A , B , and C in $\mathcal{C}$, the composition

$$\text{Mor}_{\mathcal{C}}(A, B) \otimes_k \text{Mor}_{\mathcal{C}}(B, C) \rightarrow \text{Mor}_{\mathcal{C}}(A, C)$$

is a k -module homomorphism.

A functor between additive categories over k is called a *k -additive functor*, if the induced maps on the sets of morphisms are k -module homomorphisms.

Then we define a *Mackey functor over k* as a k -additive (contravariant) functor from $\mathcal{M}_k$ to $\mathcal{C}$, an additive category over k . Since $\mathcal{M}_k = \mathcal{M}_k^{op}$, there is no difference whether we choose covariant or contravariant in the definition of the Mackey functor. However, our namings of the maps in Definition 2.2 fit more naturally with the definition which requires contravariance.

Remark 3.2. The traditional definition of the Mackey functor [14, 10, 25, 22, 24] is essentially an additive functor from the Mackey category $\mathcal{M}_k$ to the category of k -modules, which transforms finite sum in $\mathcal{M}_k$ to finite sum in the category of k -modules, with the assumption that G is finite. This latter condition is omitted in this paper. The use of $\mathcal{M}_k$ in the definition of the Mackey functor was first done by Linder [25] under a more general categorical setting.

Example 3.3. (i) Let M be a kG -module. Then the correspondence $G/H \mapsto H_*(H, M)$ defines a Mackey functor over k . Also, through the self-duality of the Mackey category, the correspondence $G/H \mapsto H^*(H, M)$ also defines a Mackey functor over k .

(ii) Let X be a free G -space. Then the correspondence $S \mapsto \Sigma^\infty X_+ \wedge_G S_+ = \Sigma^\infty (X \times_G S)_+$ is a Mackey functor (over $\mathbb{Z}$) in the category of spectra.

More examples will be stated later.

Definition 3.4. Let M be a Mackey functor over k in the category $\mathcal{C}$. Then we call $\mathcal{M}_M$ the *associated category of M* when

$$\text{Ob}_{\mathcal{M}_M} = \{\text{finite } G\text{-sets}\},$$

$$\text{Mor}_{\mathcal{M}_M}(S, T) = \text{Im}(M : \text{Mor}_{\mathcal{M}_k}(S, T) \rightarrow \text{Mor}_{\mathcal{C}}(M(S), M(T))).$$

Definition 3.5. A *cohomological Mackey functor over k* is a k -additive functor from $\mathcal{H}_k$.

Remark 3.6. (i) Originally [14, 1.4], the cohomological Mackey functor (over k) was defined to be a Mackey functor M (over k) in the sense of Remark 3.2, which further satisfies the following: If $H \subseteq K \subseteq G$ and $\beta \in M(G/K)$, then

$$\text{Ind}_H^K \circ \text{Res}_H^K(\beta) = |K : H|\beta.$$

The equivalence of these two conditions for those Mackey functors was originally proved by Yoshida [46]; we recover Yoshida's theorem below in Theorem 3.7.

(ii) Of course, the homology and the cohomology of groups are the main examples of cohomological Mackey functors (see Example 3.3).

The following was originally proved by Yoshida [46] (for the case $\mathcal{C}$ is the category of k -modules) by a direct method:

Theorem 3.7. *For Mackey functors in the category of k -modules, which transform finite sum in $\mathcal{M}_k$ to finite sum in the category of k -modules, the two definitions of the cohomological Mackey functors in Definition 3.5 and Remark 3.6 coincide.*

Proof. Let $M : \mathcal{M}_k \rightarrow \mathcal{C}$ be a cohomological Mackey functor over k in the sense of Remark 3.6. Then, by Proposition 2.25 and Remark 2.26, we can easily see that M , when restricted to each summand $A(H_i)_k$ in $\text{Mor}_{\mathcal{M}_k}(S, T)$, factors through the augmentation $\epsilon_i : A(H_i) \rightarrow k$. This immediately implies (using the fact that M transforms finite sum to finite sum) that M factorizes $\mathcal{M}_k/\mathcal{F}_\infty$, which is isomorphic to the category of the finitely generated permutation kG -modules $\mathcal{H}_k$ by Corollary 2.36. The other implication is trivial. $\square$

We now come to the central concept of this paper.

Definition 3.8. We call M , a Mackey functor over k , a *cohomotopical Mackey functor over k* , if it is a k -additive functor from $(\mathcal{M}_k)^\wedge_{\mathcal{F}}$.

Example 3.9. (i) Any cohomological Mackey functor is cohomotopical.

(ii) Let X be a G -equivariant infinite loop space (cf. [24]). Then the functor

$$S \mapsto \pi_n(\text{Map}_G(EG \times S, X))^\wedge,$$

where the completion is given by the skeletal filtration of EG , is a cohomotopical Mackey functor. This functor may be viewed as the “cohomology of group with coefficient” in homotopical algebra [42], [38]. So we could call it as “cohomotopy of group with coefficient.”

(iii) The Tate construction of [3] may be also viewed as a cohomotopical Mackey functor, just like (ii).

Now let us assume that G is a finite group and $k = \mathbb{Z}_p^\wedge$. We would like to show a characterization of the cohomotopical Mackey functor for this particular case.

However, we must first define more notation: We write $A(G)_p^\wedge$ instead of $A(G) \otimes_{\mathbb{Z}} \mathbb{Z}_p^\wedge$. Also, by $A(G)_{p+I(G)}^\wedge$, we mean the completion of $A(G)$ with respect to the ideal generated by p and $I(G)$. Notice that this is also the same as $(A(G)_p^\wedge)_{I(G)_p^\wedge}^\wedge$.

Fix a p -Sylow subgroup G_p of G . Then we may write any finite G -set S as

$$S \cong \coprod_{i=1}^n G/H_i,$$

so that $H_i \cap G_p$ is a p -Sylow subgroup of H_i for any $i = 1, 2, \dots, n$. Set $\tilde{S} := \coprod_{i=1}^n G/(H_i \cap G_p)$ and define

$$p_S := \prod_{i=1}^n p_i : \tilde{S} \cong \prod_{i=1}^n G/(H_i \cap G_p) \rightarrow \prod_{i=1}^n G/H_i \cong S,$$

where $p_i : G/(H_i \cap G_p) \rightarrow G/H_i$ is the canonical projection. We now set

$$R_S := \left[\begin{array}{ccc} & \tilde{S} & \\ p_S \swarrow & & \searrow 1_{\tilde{S}} \\ S & & \tilde{S} \end{array} \right] \in \text{Mor}_{\mathcal{M}}(S, \tilde{S}).$$

Proposition 3.10. *Let G be a finite group and $k = \mathbb{Z}_p^\wedge$. The following four conditions about M , a Mackey functor over $\mathbb{Z}_p^\wedge$, are equivalent:*

1. M is a cohomotopical Mackey functor.
2. For any finite G -sets T and S , the right composition with R_S induces an injection

$$M(R_S \circ -) : \text{Mor}_{\mathcal{C}}(M(T), M(S)) \rightarrow \text{Mor}_{\mathcal{C}}(M(T), M(\tilde{S})).$$

3. The canonical action of $A(G)_p^\wedge$ on $\text{Mor}_{\mathcal{C}}(M(T), M(S))$ factors through $A(G)_{p+I(G)}^\wedge$ for any finite G -sets T and S .
4. The canonical action of $A(G)_p^\wedge$ on the set of the morphisms of the associated category of M factors through $A(G)_{p+I(G)}^\wedge$.

Proof of 1 $\Rightarrow$ 2. It suffices to show that $R_S \in \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge}(S, \tilde{S})$ is a split-injection, i.e. there exists $g \in \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge}(\tilde{S}, S)$ such that $g \circ R_S = 1_S \in \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge}(S, S)$. Now, from the definition of R_S , it suffices to prove the claim for a transitive finite G -set $S = G/H$. (Here H is chosen so that $H \cap G_p$ is a p -Sylow subgroup of H as before.) Set

$$g_S := \left[\begin{array}{ccc} & \tilde{S} & \\ 1_{\tilde{S}} \swarrow & & \searrow p_S \\ \tilde{S} & & S \end{array} \right] \in \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge}(\tilde{S}, S).$$

Then, when $S = G/H$, the composition $g_S \circ R_S \in \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge}(G/H, G/H)$ is given by the image of $[H/(H \cap G_p)]$ under the composite (see Lemma 2.21)):

$$A(H)_{p+I(H)}^\wedge \xrightarrow{\Delta_{\text{pt}}} \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge}(H)(\text{pt}, \text{pt}) \rightarrow \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge}(G)(G/H, G/H).$$

However, $[H/(H \cap G_p)] \in A(H)_{p+I(H)}^\wedge$ is a unit, which implies

$$g_S \circ R_S \in \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge}(G/H, G/H)$$

is also a unit. Now the claim follows immediately.

Proof of 2 $\Rightarrow$ 3. Consider the following commutative diagram of the (p -completed) Burnside ring actions:

$$\begin{array}{ccc} \text{Mor}_{\mathcal{C}}(M(T), M(S)) \otimes_{\mathbb{Z}_p^\wedge} A(G)_p^\wedge & \xrightarrow{(R_S \circ -) \otimes_{\mathbb{Z}_p^\wedge} \text{Res}_{G_p}^G} & \text{Mor}_{\mathcal{C}}(M(T), M(\tilde{S})) \otimes_{\mathbb{Z}_p^\wedge} A(G_p)_p^\wedge \\ \downarrow & & \downarrow \\ \text{Mor}_{\mathcal{C}}(M(T), M(S)) & \xrightarrow[\subseteq]{R_S \circ -} & \text{Mor}_{\mathcal{C}}(M(T), M(\tilde{S})), \end{array}$$

where the commutativity follows from Corollary 2.27 and the bottom map is injective by the assumption 1.

Therefore $A(G)_p^\wedge$ -action on $\text{Mor}_C(M(T), M(S))$ factors through $\text{Im}\left(A(G)_p^\wedge \rightarrow A(G_p)_p^\wedge\right)$, which can be canonically identified with $A(G)_{p+I(G)}^\wedge$ from our knowledge of the $I(G)$ -adic topology and p -adic topology of $A(G)$ (see [18, 31]).

Proof of 3 $\Rightarrow$ 4. This is immediate from the definition of the associated category (see Definition 3.4).

Proof of 4 $\Rightarrow$ 1. This follows from $A(G)_{p+I(G)}^\wedge = (A(G)_k)_{I(G)_k}^\wedge$ with $k = \mathbb{Z}_p^\wedge$, since $\text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge}(S, T) = \text{Mor } \mathcal{M}_k(S, T) \otimes_{A(G)_k} (A(G)_k)_{I(G)_k}^\wedge$ when G is a finite group. $\square$

Corollary 3.11. *Let X be a free G -space. Then the correspondence (cf. Example 3.3)*

$$S \mapsto (\Sigma^\infty X \times_G S_+)_p^\wedge$$

is a cohomotopical Mackey Mackey functor over $\mathbb{Z}_p^\wedge$. $\square$

Remark 3.12. (i) Proposition 3.10 and its proof give yet another characterization of cohomotopical Mackey functors over $\mathbb{Z}_p^\wedge$ when G is finite. Let $(A_G)_{p+I(G)}^\wedge$ be the Green functor $S \mapsto A_G(S)_{p+I(G)}^\wedge$, where A_G is as in Definition 2.18 (which is the (Burnside) Green functor defined and denoted by Ω in [11, p.303]), and the completion is with respect to the ideal $p + I(G) \subseteq A(G)$ through its $A(G)$ -module structure. Then the cohomotopical Mackey functors over $\mathbb{Z}_p^\wedge$ for G finite are nothing but Green modules over the G/G_p -injective Green functor $(A_G)_{p+I(G)}^\wedge$. See Propositions 1.2 and 1.1' of [11] for some consequences.

(ii) Proposition 3.10 2 implies that a p -Sylow subgroup is a detecting subgroup.

4. THE MAIN THEOREM AND ITS PROOFS

In this section, we state and prove our main results. First we prove a list of results concerning the category $(\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge$ and its relations with the Hecke category $\mathcal{H}_k$ through the Hurewicz functor.

Proposition 4.1. 1. *For any finite G -set S , the Hurewicz functor $H_k : (\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge \rightarrow \mathcal{H}_k$ induces a surjection of k -algebras*

$$H_k : \left(\text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge}(S, S)\right) \twoheadrightarrow \left(\text{Mor}_{\mathcal{H}_k}(S, S)\right)$$

with its kernel quasi-regular, i.e. for any $k \in \text{Ker}(H_k : (\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge \rightarrow \mathcal{H}_k)$, $1 + k \in \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge}(S, S)$ is a unit.

2. *For any finite G -set S , the Hurewicz functor $H_k : (\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge \rightarrow \mathcal{H}_k$ induces a surjection of the group of units*

$$H_k^\times : \left(\text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge}(S, S)\right)^\times \twoheadrightarrow \left(\text{Mor}_{\mathcal{H}_k}(S, S)\right)^\times.$$

3. *The Hurewicz functor induces an isomorphism of semi-simple rings:*

$$\begin{aligned} & \left(\text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge}(S, S)\right) / J\left(\text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge}(S, S)\right) \\ & \xrightarrow{\cong} \left(\text{Mor}_{\mathcal{H}_k}(S, S)\right) / J\left(\text{Mor}_{\mathcal{H}_k}(S, S)\right) \\ & = \left(\text{Hom}_{kG}(kS, kS)\right) / J\left(\text{Hom}_{kG}(kS, kS)\right). \end{aligned}$$

Here $J(?)$ is the Jacobson radical of $(?)$, i.e. the intersection of all maximal left ideals (cf. [4, 9]).

4. Assume further that G is finite and $k = \mathbb{Z}_p^\wedge$. Then all of the above 1, 2, and 3 are also valid when $(\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge$ is replaced with $(\mathcal{M}_k)_{\mathcal{F}}^\wedge$. Furthermore, there exists $n_0 > 0$ such that, for any $f_i \in \text{Ker}(H_p : \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge}(S_i, S_{i+1}) \rightarrow \text{Mor}_{\mathcal{H}_k}(S_i, S_{i+1}))$ ($i = 1, \dots, n$) with $n \geq n_0$, we have

$$f_n \circ f_{n-1} \circ \dots \circ f_2 \circ f_1 \in p \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}_\infty}^\wedge}(S_1, S_{n+1}).$$

Proof of 1, 2 and 3. For 1, it is certainly surjective by Proposition 2.25. The remaining claim of 1 follows immediately from Proposition 2.34, which allows us to construct its inverse $\sum_{n=0}^\infty (-1)^n k^n$ explicitly. Then 2 immediately follows from 1. Furthermore, 3 is also an immediate consequence of 1 [4, 9.15, 15.3].

Proof of 4. The first claim is an immediate consequence of 1, 2, 3 and Corollary 2.37.

For the second claim, from Proposition 2.34, Corollary 2.37, and Proposition 2.25, it suffices to show there exists a certain n_0 such that for any subgroup H of a finite group G

$$(4.1) \quad \left(I(G)_{\mathbb{Z}_p^\wedge} \right)^n \cdot A(H)_{p+I(H)}^\wedge \subseteq p A(H)_{p+I(H)}^\wedge$$

for any $n \geq n_0$, because $A(H)_{p+I(H)}^\wedge = \left(A(H)_{\mathbb{Z}_p^\wedge} \right)_{I(G)_{\mathbb{Z}_p^\wedge}}^\wedge$.

To show (4.1), we choose a p -subgroup H_p of H , and notice that

$$\text{Res}_{H_p}^H : A(H)_{p+I(H)}^\wedge \hookrightarrow A(H_p)_{p+I(H_p)}^\wedge$$

is a topological split injection (cf. [31]) and the topology on $A(H_p)_{p+I(H_p)}^\wedge$ is given by the p -adic topology since H_p is a p -group ([18]).

Therefore, the $I(G)_{\mathbb{Z}_p^\wedge}$ -adic topology on $A(H)_{p+I(H)}^\wedge$ also becomes a p -adic topology, and so (since there are just finitely many subgroups inside a finite group G), there certainly exists some n_0 which satisfies (4.1). $\square$

Notice that, for G finite and $k = \mathbb{Z}_p^\wedge$, 4 implies 1, 2 and 3. We now offer a topological proof of this particular case, which is a simplified version of our original proof in [31]. Although the proof requires the Segal conjecture, we hope this would give the reader better ideas about the underlying relationship between algebra and topology.

A topological proof of 4. By Carlsson's affirmative solution of the Segal conjecture [6], the composite

$$\begin{aligned} \alpha_{(S_1, S_{n+1})} : \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge}(S_1, S_{n+1}) &\cong \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge}(S_1 \times S_{n+1}, \text{pt}) \\ &\cong \left(\{(S_1 \times S_{n+1})_+, S^0\}_G \right)_{p+I(G)}^\wedge \cong \{EG_+ \wedge_G (S_1 \times S_{n+1})_+, S^0\}_p^\wedge \end{aligned}$$

is an isomorphism. Under this isomorphism, the composite $f_n \circ f_{n-1} \circ \dots \circ f_2 \circ f_1$ is sent to

$$\begin{aligned} EG_+ \wedge_G (S_1 \times S_{n+1})_+ &\xrightarrow{1 \wedge (f_1 \times 1)} EG_+ \wedge_G (S_2 \times S_{n+1})_+ \xrightarrow{1 \wedge (f_2 \times 1)} \dots \\ &\dots \xrightarrow{1 \wedge (f_n \times 1)} EG_+ \wedge_G (S_{n+1} \times S_{n+1})_+ \xrightarrow{\alpha_{(S_{n+1}, S_{n+1})}(1_{S_{n+1}})} S_+^0. \end{aligned}$$

Now the point is that the correspondence

$$S \rightsquigarrow H_*(EG_+ \wedge_G (S \times S_{n+1})_+; \mathbb{Z}/p\mathbb{Z})$$

defines a cohomological Mackey functor, which, together with the assumption, implies

$$(1 \wedge (f_i \wedge 1))_* = 0 \quad (i = 1, \dots, n).$$

Therefore, $\alpha_{(S_1, S_{n+1})}(f_n \circ f_{n-1} \circ \dots \circ f_2 \circ f_1)$ has Adams filtration at least n . However, Carlsson's solution [6] of the Segal conjecture implies that

$$\{EG_+ \wedge_G (S_1 \times S_{n+1})_+, S^0\}_p^\wedge$$

is a finitely generated free $\mathbb{Z}_p^\wedge$ -module. Therefore, there exists some n_0 such that any element with the Adams filtration $\geq n_0$ is a multiple of p . Now the claim follows immediately. $\square$

Remark 4.2. 4.1 can be regarded as a representation theoretical analogue of the Nishida nilpotency theorem [36]. This states that any element in the kernel of the stable Hurewicz homomorphism $H : \pi_*^s(S^0) \rightarrow H_*(S^0)$ is nilpotent.

We have now come to the main theorem of this paper.

Main Theorem. (i) *Let G be a (possibly infinite) discrete group, and let S and T be finite G -sets such that the corresponding permutation kG -modules are isomorphic:*

$$kS \cong kT, \text{ as } kG\text{-modules.}$$

Then for any cohomotopical Mackey functor M ,

$$M(S) \cong M(T).$$

(ii) *Suppose $k = \mathbb{Z}_p^\wedge$. Then for any finite G -set S and any idempotent $e \in \text{Hom}_{kG}(kS, kS) = \text{Mor}_{\mathcal{H}_k}(S, S)$, there exists an idempotent $\tilde{e} \in \text{Mor}_{\mathcal{M}_k}(S, S)$, which lifts e .*

Proof of (i). From the assumption, there are

$$f \in \text{Hom}_{kG}(kS, kT) = \text{Mor}_{\mathcal{H}_k(G)}(S, T)$$

and

$$g \in \text{Hom}_{kG}(kT, kS) = \text{Mor}_{\mathcal{H}_k(G)}(T, S)$$

such that

$$(4.2) \quad f \circ g = 1_{kT}, \quad g \circ f = 1_{kS}.$$

Now, choose a finite index normal subgroup N_0 which is contained in the intersection of all the isotropy subgroups of S and T , as in Proposition 2.9. Then from the following commutative diagrams (see Proposition 2.9) and Proposition 2.25:

$$\begin{array}{ccc} \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge(G/N_0)}(S, T) & \xrightarrow{\pi_{N_0}} & \text{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^\wedge(G)}(S, T) \\ H_k \downarrow & & H_k \downarrow \\ \text{Mor}_{\mathcal{H}_k(G/N_0)}(S, T) & \xrightarrow[\cong]{\pi_{N_0}} & \text{Mor}_{\mathcal{H}_k(G)}(S, T), \end{array}$$

$$\begin{array}{ccc}
\mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G/N_0)}(T, S) & \xrightarrow{\pi_{N_0}} & \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G)}(T, S) \\
H_k \downarrow & & H_k \downarrow \\
\mathrm{Mor}_{\mathcal{H}_k(G/N_0)}(T, S) & \xrightarrow[\cong]{\pi_{N_0}} & \mathrm{Mor}_{\mathcal{H}_k(G)}(T, S),
\end{array}$$

we may find some lifts $\tilde{f} \in \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G)}(S, T)$, $\bar{f} \in \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G/N_0)}(S, T)$, $\tilde{g} \in \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G)}(T, S)$, and $\bar{g} \in \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G/N_0)}(T, S)$, so that

$$\pi_{N_0}(\tilde{f}) = \bar{f}, \quad H_k(\tilde{f}) = f, \quad \pi_{N_0}(\tilde{g}) = \bar{g}, \quad H_k(\tilde{g}) = g.$$

Although

$$\begin{aligned}
\bar{f} \circ \bar{g} &\neq 1 \in \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G/N_0)}(T, T), \\
\bar{g} \circ \bar{f} &\neq 1 \in \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G/N_0)}(S, S)
\end{aligned}$$

in general, their errors respectively belong to

$$\begin{aligned}
&\mathrm{Ker}\left(H_k : \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G/N_0)}(T, T) \rightarrow \mathrm{Mor}_{\mathcal{H}_k}(T, T)\right), \\
&\mathrm{Ker}\left(H_k : \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G/N_0)}(S, S) \rightarrow \mathrm{Mor}_{\mathcal{H}_k}(S, S)\right).
\end{aligned}$$

Thus, by Proposition 4.1 (which we may apply as G/N is finite), both $\bar{f} \circ \bar{g}$ and $\bar{g} \circ \bar{f}$ are units in the k -algebras $\mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G/N_0)}(T, T)$ and $\mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G/N_0)}(S, S)$, respectively.

Since

$$\begin{aligned}
&\mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G/N_0)}(T, T) \xrightarrow{\pi_{N_0}} \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G)}(T, T), \\
&\mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G/N_0)}(S, S) \xrightarrow{\pi_{N_0}} \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G)}(S, S)
\end{aligned}$$

are k -algebra homomorphisms, we now know that both $\tilde{f} \circ \tilde{g}$ and $\tilde{g} \circ \tilde{f}$ are units in the k -algebras $\mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G)}(T, T)$ and $\mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}(G)}(S, S)$, respectively. This implies the objects S and T are equivalent in the category $(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}$. Now the claim follows.

Proof of (ii). Arguing as the proof of (i), we may assume G is finite. Now let $e' \in \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}}(S, S)$ be any lift of e . Since e is an idempotent, we may write

$$e'^2 = e' + l,$$

where $l \in \mathrm{Ker}\left(H_p : \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}}(S, S) \rightarrow \mathrm{Mor}_{\mathcal{H}_k}(S, S)\right)$. Then, from Proposition 4.1, there exists some $n_1 \geq 1$ such that $k^{p^{n_1}} = px$ for some $x \in \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}}(S, S)$. Set $e'' := e'^{p^{n_1}}$. Then e'' is another lift of e such that

$$e''^2 = e'' + py$$

for some $y \in \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}}(S, S)$. Since $\mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}}(S, S)$ is p -adically complete (this is because G is finite), we can construct an idempotent lift of e by the standard argument using e'' ; just check that $\{e''^{p^n}\}_{n=1}^{\infty}$ is a Cauchy sequence with respect to the p -adic topology and let $\tilde{e} \in \mathrm{Mor}_{(\mathcal{M}_k)_{\mathcal{F}}^{\wedge}}(S, S)$ be the limit. Then $\tilde{e}$ is easily seen to be an idempotent lift of e . $\square$

5. WEBB'S FORMULAE AND APPLICATIONS

In [44], Webb proved interesting combinatorial formulae for Tate and ordinary cohomology of finite groups. Since the essence of his results is the corresponding combinatorial formulae about permutation $\mathbb{Z}_p^\wedge G$ -modules, we may use it as an input of our Main Theorem. For the rest of this paper, we assume G is a finite group. We first recall Webb's results:

Theorem 5.1 (Theorem A' of [44]). *Let G act simplicially on the simplicial complex Δ , suppose for each simplex σ the isotropy group G_σ fixes σ pointwise, and let p be a fixed prime. Assume that one of the following conditions holds:*

1. *for each $H \in \mathcal{C}$ with $O_p(H) \neq 1$ the fixed point complex Δ^H has Euler characteristic $\chi(\Delta^H) = 1$,*
2. *for each cyclic subgroup H of order p , Δ^H is acyclic.*

Then we have an isomorphism

$$\mathbb{Z}_p^\wedge G/G \cong \sum_{\sigma \in \Delta/G} (-1)^{\dim(\sigma)} \mathbb{Z}_p^\wedge G/G_\sigma,$$

as $\mathbb{Z}_p^\wedge G$ -modules, modulo projectives of the form $\mathbb{Z}_p^\wedge G/H$ with $(p, |H|) = 1$.

As was mentioned in [44], Quillen [39] provided interesting examples for this theorem.

Theorem 5.2 ([39]). *Theorem A' is valid when $\Delta = \mathcal{A}$ is the poset of all non-identity elementary abelian p -subgroups of G , or when $\mathcal{P}$ is the poset of non-identity p -subgroups of G , or the Tits building of a finite Chevalley group in defining characteristic p .*

In a step of his proof in [44], Webb proved the following result, which doesn't involve any equivalence modulo projectives. It is based upon Conlon's deep result [8] on the detecting subgroups of the Green ring, and the idempotent formula of the Burnside ring by [13, 47]. We denote the class of mod- p cyclic subgroups of G by $\mathcal{C}$:

$$\mathcal{C} = \{H \subseteq G \mid O_p(H), \text{ } p\text{-Sylow of } H, \text{ is a normal subgroup of } G \\ \text{and } H/O_p(H) \text{ is cyclic}\}.$$

Theorem 5.3 (Theorem D' of [44]). *Let $\mathcal{X}$ be a class of subgroups of G which is closed under taking conjugates and forming subgroups, and with $\mathcal{X} \supseteq \mathcal{C}$. Then in the Green ring of G defined over $\mathbb{Q}$,*

$$\mathbb{Z}_p^\wedge G/G = \sum_{H \in \mathcal{X}} \frac{\mu(H)}{[G:H]} \mathbb{Z}_p^\wedge G/H,$$

where $\mu: \mathcal{X} \rightarrow \mathbb{Z}$ is the function defined by the equations

$$\sum_{J \subseteq K \in \mathcal{X}} \mu(K) = 1 \text{ for every } J \in \mathcal{X}.$$

To apply Theorem 5.1 and Theorem 5.2, we provide the following lemma.

Lemma 5.4. *Let M be a cohomotopical Mackey functor over $\mathbb{Z}_p^\wedge$ such that $M(G/\{e\}) = 0$. Then for any subgroup $H \subseteq G$ such that $(|H| : p) = 1$ and $M(G/H) = 0$.*

Proof. Since $\mathbb{Z}_p^\wedge G/H$ is a projective $\mathbb{Z}_p G$ -module, there exists some $n \in \mathbb{N}$ and

$$f \in \operatorname{Hom}_{\mathbb{Z}_p^\wedge G}(\mathbb{Z}_p^\wedge G/H, (\mathbb{Z}_p^\wedge G)^{\oplus n}), \quad g \in \operatorname{Hom}_{\mathbb{Z}_p^\wedge G}((\mathbb{Z}_p^\wedge G)^{\oplus n}, \mathbb{Z}_p^\wedge G/H)$$

such that

$$g \circ f = 1_{\mathbb{Z}_p^\wedge G/H} \in \operatorname{Hom}_{\mathbb{Z}_p^\wedge G}(\mathbb{Z}_p^\wedge G/H, \mathbb{Z}_p^\wedge G/H).$$

Then, arguing as the proof of our main theorem, we see that $M(G/H)$ is a direct summand of $M(n \cdot G/\{e\}) = M(G/\{e\})^{\oplus n} = 0$. $\square$

Theorem 5.5. *Let M be a cohomotopical Mackey functor over $\mathbb{Z}_p^\wedge$ such that $M(G/\{e\}) = 0$. Then, under the same assumption and the notation of Theorem 5.1, we have a formal isomorphism of continuous $\mathbb{Z}_p^\wedge$ -modules:*

$$M(G/G) = \sum_{\sigma \in \Delta/G} (-1)^{\dim(\sigma)} M(G/G_\sigma).$$

Proof. This is a direct consequence of Theorem 5.3, the Main Theorem, and Lemma 5.4. $\square$

Similarly, Theorem 5.3, with the help of the Main Theorem, implies the following:

Theorem 5.6. *Let M be a cohomotopical Mackey functor over $\mathbb{Z}_p^\wedge$. Then, under the same assumption and the notation of Theorem 5.3, we have a formal isomorphism of continuous $\mathbb{Z}_p^\wedge$ -modules:*

$$M(G/G) = \sum_{H \in \mathcal{X}} \frac{\mu(H)}{[G:H]} M(G/H). \quad \square$$

6. TOPOLOGICAL APPLICATIONS

In this section we give some topological applications of our algebraic investigations. We first recall that the Eilenberg-MacLane space $K(M, n)$ [5, 12] associated with a Mackey functor M becomes an infinite loop G -space and they form an $RO(G)$ -graded cohomology theory $\tilde{H}_G^*(?, M)$ [22]. It's easy to see, for any G -space X and $* \in RO(G)$, that the functor $S \mapsto \tilde{H}_G^*(X_+ \wedge S_+, M)$ becomes a cohomotopical Mackey functor over $\mathbb{Z}_p^\wedge$ whenever M is. As in the previous section, we assume G is a finite group. Thus we have the following corollaries of Theorems 5.5 and 5.6.

Corollary 6.1. *Let X be a G -space, $* \in RO(G)$, and M a cohomotopical Mackey functor such that $\tilde{H}_G^*(X_+ \wedge G/\{e\}_+, M) \cong H^*(X, M(G/\{e\})) = 0$, where $\bar{*}$ is the dimension of $*$. Then under the same assumption and the notation of Theorem 5.1, we have a formal isomorphism of continuous $\mathbb{Z}_p^\wedge$ -modules:*

$$\tilde{H}_G^*(X_+, M) = \sum_{\sigma \in \Delta/G} (-1)^{\dim(\sigma)} \tilde{H}_{G_\sigma}^*(X_+, M). \quad \square$$

Corollary 6.2. *Let X be a G -space, $* \in RO(G)$, and M a cohomotopical Mackey functor. Then, under the same assumption and the notation of Theorem 5.3, we have a formal isomorphism of continuous $\mathbb{Z}_p^\wedge$ -modules:*

$$\tilde{H}_G^*(X_+, M) = \sum_{H \in \mathcal{X}} \frac{\mu(H)}{[G:H]} \tilde{H}_H^*(X_+, M). \quad \square$$

In particular, when we take $M = \mathbb{Z}_p^\wedge$, the $\mathbb{Z}_p^\wedge$ -constant Mackey functor [22] [24, v.9.10], we have the following corollaries.

Corollary 6.3. *Let a G -space X and $i \in \mathbb{N}$ be such that $H^i(X, \mathbb{Z}_p^\wedge) = 0$. Then, under the same assumption and the notation of Theorem 5.1, we have a formal isomorphism of continuous $\mathbb{Z}_p^\wedge$ -modules:*

$$H^i(X/G, \mathbb{Z}_p^\wedge) = \sum_{\sigma \in \Delta/G} (-1)^{\dim(\sigma)} H^i(X/G_\sigma, \mathbb{Z}_p^\wedge). \quad \square$$

Corollary 6.4. *Let X be a G -space and $i \in \mathbb{N}$. Then, under the same assumption and the notation of Theorem 5.3, we have a formal isomorphism of continuous $\mathbb{Z}_p^\wedge$ -modules:*

$$H^i(X/G, \mathbb{Z}_p^\wedge) = \sum_{H \in \mathcal{X}} \frac{\mu(H)}{[G:H]} H^i(X/H, \mathbb{Z}_p^\wedge). \quad \square$$

Actually, these two results are immediate consequences of Yoshida's Theorem [46] (see Theorem 3.7); just check that the classical transfer for the (possibly ramified) covering forms a cohomological Mackey functor (in the original sense).

Theorem 6.5. *Let X be a free G -space such that $(\Sigma^\infty X)_p^\wedge$ is contractible. Then under the same assumption and the notation of Theorem 5.1, we have a formal equivalence of spectra:*

$$(\Sigma^\infty X/G)_p^\wedge = \sum_{\sigma \in \Delta/G} (-1)^{\dim(\sigma)} (\Sigma^\infty X/G_\sigma)_p^\wedge.$$

Proof. By Corollary 3.11, the correspondence

$$S \mapsto (\Sigma^\infty X \times_G S_+)_p^\wedge$$

is a cohomotopical Mackey functor over $\mathbb{Z}_p^\wedge$. From the assumption and the usual transfer argument, we see that

$$(\Sigma^\infty X \times_G G/H_+)_p^\wedge = (\Sigma^\infty X/H_+)_p^\wedge \simeq (\Sigma^\infty S^0)_p^\wedge$$

for any subgroup H of G with $(p, |H|) = 1$. So, applying Theorem 5.1 and the Main Theorem to the above cohomotopical Mackey functor, we see that

$$(\Sigma^\infty X/G_+)_p^\wedge = \sum_{\sigma \in \Delta/G} (-1)^{\dim(\sigma)} (\Sigma^\infty X/G_{\sigma_+})_p^\wedge,$$

modulo finitely many wedge sums of $(\Sigma^\infty S^0)_p^\wedge$. However, $(\Sigma^\infty X/K)_p^\wedge$ is connected for any subgroup K of G , by the assumption. Thus the above equivalence implies the equality

$$(\Sigma^\infty X/G)_p^\wedge = \sum_{\sigma \in \Delta/G} (-1)^{\dim(\sigma)} (\Sigma^\infty X/G_\sigma)_p^\wedge,$$

as was desired. $\square$

Theorem 6.6. *Let X be a free G -space, Then, under the same assumption and the notation of Theorem 5.3, we have a formal isomorphism of spectra:*

$$(\Sigma^\infty X/G)_p^\wedge = \sum_{H \in \mathcal{X}} \frac{\mu(H)}{[G:H]} (\Sigma^\infty X/H)_p^\wedge.$$

Proof. As in the proof of Theorem 6.5, we obtain an equality

$$(\Sigma^\infty X/G_+)_p^\wedge = \sum_{H \in \mathcal{X}} \frac{\mu(H)}{[G:H]} (\Sigma^\infty X/H_+)_p^\wedge,$$

by applying Theorem 5.6 this time. However, it is easy to see that

$$1 = \sum_{H \in \mathcal{X}} \frac{\mu(H)}{[G:H]}$$

from Theorem 5.3 (cf. [44, Prop.8.4(ii)]). This means we can eliminate the extra base points from the above formal equality of spectra, which completes the proof. $\square$

Remark 6.7. For $X = EG$, Theorem 6.6 was called the “Minami-Webb formula” in [27, 28]. There is a recent interesting preprint of Martino and Priddy [29] on this subject.

We now offer another proof of Theorem 6.5 and Theorem 6.6 without using our Main Theorem; this proof is the simplest and most elementary one for these two results. For this purpose, it suffices to provide an elementary proof of the following fact (which is as a matter of course an immediate consequence of the Main Theorem):

Lemma 6.8. *Let G be a (possibly infinite) discrete group, and let S and T be finite G -sets such that*

$$\mathbb{F}_p S \cong \mathbb{F}_p T$$

as permutation $\mathbb{F}_p G$ -modules. Then for any free G -space X , there is an equivalence of spectra:

$$(\Sigma^\infty X \times_G S)_p^\wedge = (\Sigma^\infty X \times_G T)_p^\wedge.$$

Proof. From the assumption, there are

$$f \in \text{Hom}_{\mathbb{F}_p G}(\mathbb{F}_p S, \mathbb{F}_p T) \quad \text{and} \quad g \in \text{Hom}_{\mathbb{F}_p G}(\mathbb{F}_p T, \mathbb{F}_p S)$$

such that

$$(6.1) \quad f \circ g = 1_{\mathbb{F}_p T}, \quad g \circ f = 1_{\mathbb{F}_p S}.$$

Then let $f' \in \text{Hom}_{\mathbb{Z} G}(\mathbb{Z} S, \mathbb{Z} T)$ and $g' \in \text{Hom}_{\mathbb{Z} G}(\mathbb{Z} T, \mathbb{Z} S)$ be arbitrary lifts of $f \in \text{Hom}_{\mathbb{F}_p G}(\mathbb{F}_p S, \mathbb{F}_p T)$ and $g \in \text{Hom}_{\mathbb{F}_p G}(\mathbb{F}_p T, \mathbb{F}_p S)$, respectively.

We now claim that lifts $\tilde{f} \in \text{Mor}_{\mathcal{M}}(S, T)$ and $\tilde{g} \in \text{Mor}_{\mathcal{M}}(T, S)$ of f' and g' respectively exist, and that $\tilde{f}$ and $\tilde{g}$ respectively induce the corresponding spectra maps

$$\begin{aligned} F : \Sigma^\infty X \times_G S_+ &\rightarrow \Sigma^\infty X \times_G T_+, \\ G : \Sigma^\infty X \times_G T_+ &\rightarrow \Sigma^\infty X \times_G S_+. \end{aligned}$$

Although the existence of such lifts and the resulting spectra maps are more or less immediate consequences of Corollary 2.36 and Example 3.3(ii), we now make a

little detour by constructing such lifts and spectra maps explicitly for the reader's convenience, especially for those readers who are simply interested in this result on spectra and have decided to skip the cohomotopical development.

For this purpose, we first set some notation. Given a sequence of finite index subgroups $K \subseteq H \subseteq G$ and an element $g \in G$, we let

$$\begin{aligned} i_K^H &: (X \times_G G/K)_+ \rightarrow (X \times_G G/H)_+, \\ c_g &: (X \times_G G/K^g)_+ \rightarrow (X \times_G G/K)_+ \end{aligned}$$

denote maps induced by the inclusion $K \subseteq H$ and the conjugation $K^g \rightarrow K$; $g^{-1}kg \rightarrow k$, respectively. Furthermore, we let

$$t_K^H : \Sigma^\infty(X \times_G G/H)_+ \rightarrow \Sigma^\infty(X \times_G G/K)_+$$

denote the stable transfer map induced by the inclusion $K \subseteq H$. Notice that these maps respectively correspond to Res_K^H , c_g , Ind_K^H in Definition 2.2.

Next, let L and M be two finite index subgroups of G . Through the isomorphisms

$$\begin{aligned} \text{Mor}_{\mathcal{H}}(G/L, G/M) &= \text{Hom}_{\mathbb{Z}G}(\mathbb{Z}G/L, \mathbb{Z}G/M) \\ &\cong \text{Hom}_{\mathbb{Z}L}(\mathbb{Z}, \mathbb{Z}G/M) \cong (\mathbb{Z}G/M)^L \cong \mathbb{Z}[L \backslash G/M], \end{aligned}$$

we see that $\text{Mor}_{\mathcal{H}}(G/L, G/M)$ is a free abelian group with elements of the form $LxM \in \mathbb{Z}L \backslash G/M$ as its basis. Furthermore, we see that this basis element LxM corresponds to

$$\left(gL \mapsto g \sum_{l \in L/L \cap M^{x^{-1}}} l x M \right) \in \text{Hom}_{\mathbb{Z}G}(\mathbb{Z}G/L, \mathbb{Z}G/M)$$

through the above isomorphism. But it follows easily from Definition 2.3 that this element is the Hurewicz functor image of

$$\begin{aligned} &c_{x^{-1}} \circ \text{Res}_{L \cap M^{x^{-1}}}^{M^{x^{-1}}} \circ \text{Ind}_{L \cap M^{x^{-1}}}^L \\ &= \left[\begin{array}{ccc} & G/L \cap M^{x^{-1}} & \\ p \swarrow & & \searrow rp \\ G/L & & G/M \end{array} \right] \in \text{Mor}_{\mathcal{M}}(G/L, G/M), \end{aligned}$$

where p is the obvious canonical projection and $rp : G/L \cap M^{x^{-1}} \rightarrow G/M$ is the composite of the canonical projection onto $G/M^{x^{-1}}$ and $r_x : G/M^{x^{-1}} \rightarrow G/M$; $gM^{x^{-1}} = gxMx^{-1} \mapsto gxM$, the right multiplication by x .

Motivated by this observation, we define a map

$$\begin{aligned} \alpha_X(G/L, G/M) &: \text{Hom}_{\mathbb{Z}G}(\mathbb{Z}G/L, \mathbb{Z}G/M) \\ &\rightarrow \{\Sigma^\infty(X \times_G G/L)_+, \Sigma^\infty(X \times_G G/M)_+\} \end{aligned}$$

by

$$LxM \mapsto c_{x^{-1}} \circ i_{L \cap M^{x^{-1}}}^{M^{x^{-1}}} \circ t_{L \cap M^{x^{-1}}}^L.$$

To generalize this definition, recall the canonical decompositions:

$$\begin{aligned} \operatorname{Hom}_{\mathbb{Z}G}(\mathbb{Z}S, \mathbb{Z}T) &= \bigoplus_{i,j} \operatorname{Hom}_{\mathbb{Z}G}(\mathbb{Z}G/L_i, \mathbb{Z}G/M_j), \\ \{\Sigma^\infty(X \times_G S)_+, \Sigma^\infty(X \times_G T)_+\} \\ &= \bigoplus_{i,j} \{\Sigma^\infty(X \times_G G/L_i)_+, \Sigma^\infty(X \times_G G/M_j)_+\}, \end{aligned}$$

where $S = \coprod_i G/L_i$, $T = \coprod_j G/M_j$.

Then the above definition is generalized to

$$\alpha_X(S, T) : \operatorname{Hom}_{\mathbb{Z}G}(\mathbb{Z}S, \mathbb{Z}T) \rightarrow \{\Sigma^\infty(X \times_G S)_+, \Sigma^\infty(X \times_G T)_+\},$$

by setting the (i, j) -component to be $\alpha_X(G/L_i, G/M_j)$.

Having done this, we may simply set

$$\begin{aligned} F &= \alpha_X(S, T)(\tilde{f}) : \Sigma^\infty X \times_G S_+ \rightarrow \Sigma^\infty X \times_G T_+, \\ G &= \alpha_X(T, S)(\tilde{f}) : \Sigma^\infty X \times_G T_+ \rightarrow \Sigma^\infty X \times_G S_+, \end{aligned}$$

which finishes our little detour.

Now we are ready to finish the proof. Although

$$F \circ G \neq 1_{\Sigma^\infty X \times_G T_+}, \quad G \circ F \neq 1_{\Sigma^\infty X \times_G S_+}$$

in general, they are the same at the level of mod- p homology. In fact, this follows from (6.1) by noticing that the correspondence

$$S \mapsto \tilde{H}_*(\Sigma^\infty X \times_G S_+; \mathbb{F}_p)$$

defines a cohomological Mackey functor over $\mathbb{F}_p$. Of coures, this completes the proof. $\square$

Remark 6.9. 1. The reader might wonder why we should consider all the mod- p cyclic subgroups even though the mod- p cohomology of a group is detected by its p -Sylow subgroup. Now the following example by [35] negatively answers this question:

$$B(\mathbb{Z}/2 \times \mathbb{Z}/2) \simeq BA_4 \vee L(2) \vee L(2) \vee B\mathbb{Z}/2 \vee B\mathbb{Z}/2, \text{ as } 2\text{-local spectra.}$$

In this example, no matter how we try to express BA_4 using 2-subgroups $\mathbb{Z}/2 \times \mathbb{Z}/2$, $\mathbb{Z}/2$, we can never do it, since we can't handle $L(2)$.

2. In [44], the following formal isomorphism as $\mathbb{Z}_2^\wedge \Sigma_4$ -modules was given as an example:

$$\mathbb{Z}_2^\wedge \Sigma_4 / \Sigma_4 = \frac{1}{2} \mathbb{Z}_2^\wedge \Sigma_4 / A_4 + \mathbb{Z}_2^\wedge \Sigma_4 / D_8 - \frac{1}{2} \mathbb{Z}_2^\wedge \Sigma_4 / (\mathbb{Z}/2)^2.$$

This is consistent with the following 2-local stable splittings due to [35]:

$$\begin{aligned} B\Sigma_4 &\simeq BPSL_2(\mathbb{F}_7) \vee L(2) \vee B\mathbb{Z}/2, \\ BD_8 &\simeq BPSL_2(\mathbb{F}_7) \vee L(2) \vee L(2) \vee B\mathbb{Z}/2 \vee B\mathbb{Z}/2, \end{aligned}$$

which together imply

$$B\Sigma_4 \simeq \frac{1}{2}BA_4 + BD_8 - \frac{1}{2}B(\mathbb{Z}/2)^2.$$

This coincidence led Stewart Priddy to conjecture Theorem 6.6 for $X = EG$, which in turn became the starting point of this research.

3. In view of [33], our main theorem may be applied to study the classifying spaces of more general infinite discrete groups.
4. While our results do not say anything about (non-finite) compact Lie groups, the Mackey functor (and even the Hecke category) for compact Lie groups was defined and studied in [24, 43, 21]. Of course, the problem is how to define the cohomotopical Mackey category in this context (cf. [43]) and (if so) whether it is useful (like the finite group case or not). As a related subject, [19] and [34] may be regarded as studies of the so-called Burnside category for compact Lie groups.

REFERENCES

1. J. F. Adams, *On the structure and applications of the Steenrod algebra*, Comment. Math. Helvetici **32** (1958), 180–214. MR **20**:2711
2. J. F. Adams, *Prerequisites (on equivariant theory) for Carlsson's lecture*, Lecture Notes in Math. **1051** Springer-Verlag, Berlin-New York. 1984, 483–532. MR **86f**:57037
3. A. Adem, R. L. Cohen and W. G. Dwyer, *Generalized Tate homology, homotopy fixed points and the transfer*, Contemporary Mathematics **96**, American Mathematical Society (1989), 1–13. MR **90k**:55012
4. F. W. Anderson and K. R. Fuller, *Rings and Categories of Modules*, 2nd ed., Graduate Texts in Mathematics 13, Springer-Verlag, New York-Berlin-Heidelberg (1992). MR **94i**:16001
5. G. E. Bredon, *Equivariant cohomology theories*, Lecture Notes in Math. **34** Springer-Verlag, Berlin-New York. (1967). MR **35**:4914
6. G. Carlsson, *Equivariant stable homotopy and Segal's Burnside ring conjecture*, Annals of Math. **120** (1984), 189–224. MR **86f**:57036
7. E. Cline, B. Parshall and L. Scott, *Cohomology of finite groups of Lie type, I*, Publ. Math. I.H.E.S. **45** (1975) 169–191. MR **53**:3134
8. S. B. Conlon, *Decompositions induced from the Burnside Algebra*, J.Algebra **10** (1968), 102–122. MR **38**:5945
9. C. Curtis and I. Reiner, *Representation theory of finite groups and associative algebras*, Interscience Publishers, New York (1962). MR **26**:2519
10. A. Dress, *Contributions to the theory of induced representations*, Lecture Notes in Math. **342** Springer-Verlag, Berlin-New York. 1973, 183–240. MR **52**:5787
11. A. Dress, *Induction and structure theorems for orthogonal representations of finite groups*, Annals of Math. **102** (1975), 291–325. MR **52**:8235
12. A. Elmendorf, *Systems of fixed point sets*, Trans. Amer. Math. Soc. **277** (1983), 275–284. MR **84f**:57029
13. D. Gluck, *Idempotent formula for the Burnside algebra with applications to the p -subgroup simplicial complex*, Ill. J. Math. **25** (1981), 63–67. MR **82c**:20005
14. J. A. Green, *Axiomatic representation theory for finite groups*, J. Pure and Applied Algebra **1** (1971), 41–77. MR **43**:4931
15. J. P. C. Greenlees, *Representing Tate cohomology of G -spaces*, Proc. Edinburgh Math. Soc. **30** (1987), 435–443. MR **88m**:57055
16. J. P. C. Greenlees and J. P. May, *Generalized Tate cohomology*, Mem. Amer. Math. Soc. **113** (1995), no. 543, viii+178 pp. MR **96e**:55006
17. C. Kosniowski, *Equivariant cohomology and stable cohomotopy*, Math. Ann. **210** (1974), 83–104. MR **54**:1202
18. E. Laitinen, *On the Burnside ring and stable cohomotopy of a finite group*, Math. Scand. **44** (1979), 37–72. MR **80k**:55030
19. C-N. Lee and N. Minami, *Segal's Burnside ring conjecture for compact Lie groups*, pp.133–161, Algebraic topology and Its Applications (1994) Springer-Verlag. MR **95b**:55001
20. L. G. Lewis, Jr., *An introduction to Mackey functors*, a draft.
21. L. G. Lewis, Jr., *The Category of Mackey Functors for a Compact Lie Group*, Group Representations: Cohomology, Group Actions and Topology, Proc. Sympos. Pure Math., vol. 63, Amer. Math. Soc., Providence, RI, 1998, pp. 301–354. CMP 98:08
22. L. G. Lewis, Jr., J. P. May and J. E. McClure, *Ordinary $RO(G)$ -graded cohomology*, Bull. Amer. Math. Soc. **4** (1981), 208–212. MR **82e**:55008

23. L. G. Lewis, Jr., J. P. May and J. E. McClure, *Classifying G -spaces and the Segal conjecture*, Canadian Math. Soc. Conference Proceedings Vol 2, part 1 (1982), 165–179. MR **84d**:55007a
24. L. G. Lewis, Jr., J. P. May and M. Steinberger (with contributions by J. E. McClure), *Equivariant Stable Homotopy Theory*, Lecture Notes in Math. **1213** Springer-Verlag, Berlin-New York. 1986. MR **88e**:55002
25. H. Linder, *A remark on Mackey functors*, Manuscripta Math. **18** (1976), 273–278. MR **53**:5691
26. F. Luca, *The algebra of Green and Mackey functors*, The University of Alaska Fairbanks Ph. D. Thesis. (1996).
27. J. Martino and S. Priddy, *A classification of the stable type of BG* , Bull. Amer. Math. Soc. (N.S.) **27** (1992), 165–170. MR **93b**:55019
28. J. Martino and S. Priddy, *The complete stable splitting for the classifying space of a finite group*, Topology **31** (1992), 143–156. MR **93d**:55012
29. J. Martino and S. Priddy, *Applications of the Minami-Webb formula to splitting classifying spaces*, preprint.
30. J. P. May, *Stable maps between classifying spaces*, Contemporary Mathematics **37** (1985), 121–129. MR **86m**:55022
31. N. Minami, *On the $I(G)$ -adic topology of the Burnside rings of compact Lie groups*, Publ. Res. Inst. Math. Sci. **20** (1984), 447–460. MR **86g**:57029
32. N. Minami, “*A Hecke Algebra*” for cohomotopical Mackey functors, the stable homotopy of orbit spaces, and a Möbius function, MSRI preprint series 00425-91 (1990).
33. N. Minami, *On the classifying spaces of $SL_3(\mathbb{Z})$, $St_3(\mathbb{Z})$ and finite Chevalley groups*, Contemp. Math., vol. 158 (1994), 175–185. MR **95b**:55015
34. N. Minami, *The relative Burnside module and stable maps between classifying spaces of compact Lie groups*, Trans. Amer. Math. Soc., **347** (1995), 461–498. MR **95i**:55008
35. S. Mitchell and S. Priddy, *Symmetric product spectra and splittings of classifying spaces*, Amer. J. Math. **106** (1984), 219–232. MR **86g**:55009
36. G. Nishida, *The nilpotency of elements of the stable homotopy groups of spheres*, J. Math. Soc. Japan **25** (1973), 707–732. MR **49**:6236
37. S. Priddy, *Recent progress in stable splittings*, London Mathematical Society Lecture Notes Series **117**, 149–174, Cambridge University Press 1987. MR **89f**:55017
38. D. Quillen, *Homotopical algebra*, Lecture Notes in Math. **43** Springer-Verlag, Berlin-New York. 1967. MR **36**:6480
39. D. Quillen, *Homotopy properties of the poset of nontrivial p -subgroups of a group*, Advances in Math. **28** (1978), 101–128. MR **80k**:20049
40. G. Shimura, *Introduction to the arithmetic theory of automorphic functions*, Iwanami Shoten and Princeton University Press, 1971. MR **47**:3318
41. J. Thévenaz and P. Webb, *The structure of Mackey functors*, Trans. Amer. Math. Soc. **347** (1995), 1865–1961. MR **95i**:22018
42. R. W. Thomason, *The homotopy limit problem*, Contemporary Mathematics **19**, American Mathematical Society 1983. MR **84j**:18012
43. S. Waner, *Mackey functors and G -cohomology*, Proc. Amer. Math. Soc., **85** (1982), 641–648. MR **85e**:55013
44. P. J. Webb, *A Local Method in Group Cohomology*, Comment. Math. Helvetici **62** (1987), 135–167. MR **88h**:22065
45. K. Wirthmüller, *Equivariant homology and duality*, Manuscripta Math. **11** (1974), 373–390. MR **49**:8004
46. T. Yoshida, *On G -functors (II): Hecke operators and G -functors*, J. Math. Soc. Japan **35** (1983), 179–190. MR **84b**:20010
47. T. Yoshida, *Idempotents of the Burnside rings and Dress induction theorem*, J. Algebra **80** (1983), 90–105. MR **85d**:20004

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF ALABAMA, BOX 870350, TUSCALOOSA ALABAMA 35487-0350

E-mail address: norihiko@gp.as.ua.edu