LINK COMPLEMENTS AND THE BIANCHI MODULAR GROUPS

MARK D. BAKER

ABSTRACT. We determine the values of m for which the Bianchi modular group $\mathrm{PSL}_2(\mathcal{O}_m)$ contains a link group.

1. INTRODUCTION AND STATEMENT OF RESULTS

Let $\mathcal{O}_m$ be the integers of the imaginary quadratic number field $\mathbb{Q}(\sqrt{-m})$. The Bianchi modular group, $\mathrm{PSL}_2(\mathcal{O}_m)$, is a discrete subgroup of $\mathrm{PSL}_2(\mathbb{C})$ the orientation preserving isometries of hyperbolic 3-space, $\mathbb{H}^3$.

In this paper we determine those values of m for which $\mathrm{PSL}_2(\mathcal{O}_m)$ contains a link group, that is, a torsion-free subgroup of finite index, Γ , such that $\mathbb{H}^3/\Gamma$ is homeomorphic to a link complement in S^3 .

Results on the cohomology of $\mathrm{PSL}_2(\mathcal{O}_m)$ (see [V]) limit the values of m for which this is possible to the following list, $\mathcal{L}$:

$$\mathcal{L} = \{1, 2, 3, 5, 6, 7, 11, 15, 19, 23, 31, 39, 47, 71\}.$$

Over the last 20 years, numerous link groups have been found in $\mathrm{PSL}_2(\mathcal{O}_m)$ for the seven cases $m = 1, 2, 3, 7, 11, 15, 23$ (see [B], [H1], [R1], [T], [Wi]). Furthermore, many of the corresponding links (figure eight knot, Whitehead link, Borromean rings, . . .) have been central to the study of 3-manifolds.

We prove that the seven remaining Bianchi groups from the above list, $\mathcal{L}$, also contain link groups; hence:

Theorem. *The Bianchi group $\mathrm{PSL}_2(\mathcal{O}_m)$ contains a link group if and only if $m \in \mathcal{L}$.*

2. PRELIMINARIES AND OUTLINE OF PROOF

Henceforth, let Ω_m denote the orbifold $\mathbb{H}^3/\mathrm{PSL}_2(\mathcal{O}_m)$. We prove our result by:

i) Embedding Ω_m in S^3 as the complement of a link of circles. The orbifolds Ω_m for $m = 5, 6, 15, 19, 23, 31, 39, 47, 71$ are drawn in Figure 1.

ii) Using the embeddings in (i) to prove that Ω_m admits a finite sheeted orbifold covering $\tilde{\Omega}_m \rightarrow \Omega_m$ such that $\tilde{\Omega}_m$ is a nonsingular hyperbolic link complement.

Thus $\tilde{\Omega}_m \cong \mathbb{H}^3/\Gamma$, where Γ is a torsion-free finite index subgroup of $\mathrm{PSL}_2(\mathcal{O}_m)$ and hence Γ is a link group (see [T], Chapter 13). The existence of $\tilde{\Omega}_m$ follows from an analysis of the singular locus of Ω_m .

In Section 3 we construct the link complements, $\tilde{\Omega}_m$. We also apply our methods to construct link complements corresponding to subgroups of $\mathrm{PSL}_2(\mathcal{O}_{15})$ and

Received by the editors June 3, 1998 and, in revised form, March 8, 1999.

2000 *Mathematics Subject Classification.* Primary 57M25; Secondary 11F06.

Key words and phrases. Link complements, Bianchi modular groups.

FIGURE 1. The orbifolds Ω_m

$\mathrm{PSL}_2(\mathcal{O}_{23})$ giving further examples in these two cases (see [B] for previous examples). Section 4 is devoted to a discussion of the orbifolds Ω_m and their embeddings in S^3 . In Section 5 we give the volumes of $\tilde{\Omega}_m$ calculated using SnapPea. Comparing these volumes with expected theoretical values provides a good check of our results.

3. LINK COMPLEMENTS: THE COVERS $\tilde{\Omega}_m$

Consider the orbifolds Ω_m drawn in Figure 1. Each Ω_m is a singular link complement. The components of the link are labelled by ∞ , while segments of the singular locus of cone angle π (resp. $2\pi/3$) are labelled by 2 (resp. 3).

3.1. The cases Ω_m , $m = 15, 23, 47, 71$. Notice that the singular locus of these four orbifolds consists of two unknotted circles—one of cone angle π , the other of cone angle $2\pi/3$ —linked as shown:

Now we simply use the fact that the n -fold cyclic cover of S^3 branched over an unknotted circle is homeomorphic to S^3 . Let $\Omega'_m \rightarrow \Omega_m$ be the 3-fold cyclic cover branched over the circle of cone angle $2\pi/3$. Then the orbifold Ω'_m is a link complement with singular locus an unknotted circle Σ of cone angle π . Hence the 2-fold cover $\tilde{\Omega}_m \rightarrow \Omega'_m$ branched over Σ gives the desired nonsingular link complement.

We have drawn these links for $m = 15, 23, 47$ in Figures 2–4. As for $m = 71$, the complexity of the resulting link is such that we leave this as an exercise for the reader!

FIGURE 2.

FIGURE 3.

FIGURE 4.

3.2. **The case Ω_{31} .** The singular locus of Ω_{31} is:

The edges of cone angle π form an unknotted circle, Σ_2 . Let $\Omega'_{31} \rightarrow \Omega_{31}$ be the 2-fold cyclic cover, branched over Σ_2 . Thus the orbifold Ω'_{31} , is again a link complement, with singular locus an unknotted circle, Σ_3 , of cone angle $2\pi/3$. Hence the 3-fold cyclic cover of Ω'_{31} branched over Σ_3 is a nonsingular link complement, drawn in Figure 5.

3.3. **The cases Ω_m , $m = 5, 6, 39$.** We construct the covers $\tilde{\Omega}_m \rightarrow \Omega_m$ using, in addition to the above techniques, the following lemma.

FIGURE 5.

FIGURE 6a.

Lemma. (a) The orbifold $\left(\begin{array}{c} \infty \\ \text{---} \\ 2 \end{array} \right)_2$ is 2-fold branch covered by $\left(\begin{array}{c} \infty \\ \text{---} \\ \infty \end{array} \right)_\infty$.

(b) The orbifold $\left(\begin{array}{c} \infty \\ \text{---} \\ 2 \end{array} \right)_3$ is 3-fold (irregular) branch covered by $\left(\begin{array}{c} \infty \\ \text{---} \\ 2 \end{array} \right)_\infty$.

Proof. We prove (b). The proof of (a) is analogous. We begin by compactifying the orbifold (b) by removing a tubular neighborhood of its link. Now consider its 3-fold cyclic cover, branched over Σ_3 pictured in Figure 6a. The preimage of Σ_2 consists of three unknotted circles of cone angle π . We further identify two of these singular circles by cutting along their (punctured) bounding disks and then identifying D_1^+ with D_2^- (resp. D_1^- with D_2^+) by pushing everything through the tube, T (Figures 6b–6c). $\square$

Notice that in these three cases, Ω_m contains the configuration of lemma (b):

Let $\Omega'_m \rightarrow \Omega_m$ be the corresponding 3-fold irregular branched cover.

3.3.1. *The case Ω_5 .* The construction of Ω'_5 is illustrated in Figure 7a. The singular locus of Ω'_5 is $\left(\begin{array}{c} 2 \\ \text{---} \\ 2 \end{array} \right)_2$ which we desingularize by taking cyclic covers as illustrated in Figure 7b. We obtain $\tilde{\Omega}_5$ an 8-circle nonsingular link complement.

3.3.2. *The case Ω_6 .* The 3-fold irregular cover $\Omega'_6 \rightarrow \Omega_6$ is drawn in Figure 8a. Now Ω'_6 contains the configuration of lemma (a); hence we obtain the 2-fold branched cover $\Omega''_6 \rightarrow \Omega'_6$ pictured in Figure 8b. The singular locus of Ω''_6 is one circle of cone angle π , so that its 2-fold branched cover yields the 12-circle link complement in Figure 8c.

3.3.3. *The case Ω_{39} .* The 3-fold irregular branched cover $\Omega'_{39} \rightarrow \Omega_{39}$ is drawn in Figure 9. The singular locus of Ω'_{39} is:

Thus (by applying verbatim the steps in §3.2) we know that Ω'_{39} is 6-fold covered by a nonsingular link complement. Given the complexity of this link, we do not draw it here.

FIGURE 6b.

FIGURE 6c.

3.4. The case Ω_{19} . The orbifold Ω_{19} , owing to its complicated singular locus, was the hardest to deal with. We began by considering the quotient group

$$\mathrm{PSL}_2(\mathcal{O}_{19}) / \left\langle \left\langle \begin{bmatrix} 1 & \omega \\ 0 & 1 \end{bmatrix} \right\rangle \right\rangle, \quad \omega = \frac{-1 + \sqrt{-19}}{2},$$

which the computer program GAP ([G]) found to be of order 60. This implies the following:

Proposition. *The congruence subgroup $\Gamma(I) \subset \mathrm{PSL}_2(\mathcal{O}_{19})$, $I = (\omega)$, is the fundamental group of a link complement in a homotopy 3-sphere.*

Proof. Recall that $\Gamma(I)$ is the kernel of $\mathrm{PSL}_2(\mathcal{O}_{19}) \rightarrow \mathrm{PSL}_2(\mathcal{O}_{19}/I)$ (see [S] for details). Since $\mathcal{O}_{19}/I \cong \mathbb{Z}/5\mathbb{Z}$, it follows that $\Gamma(I)$ is a normal, torsion-free subgroup of index 60 in $\mathrm{PSL}_2(\mathcal{O}_{19})$ containing the matrix $\begin{bmatrix} 1 & \omega \\ 0 & 1 \end{bmatrix}$. Thus $\Gamma(I) = \langle \langle \begin{bmatrix} 1 & \omega \\ 0 & 1 \end{bmatrix} \rangle \rangle$ so that $\Gamma(I)$ is generated by $\mathrm{PSL}_2(\mathcal{O}_{19})$ -conjugates of the parabolic element $\begin{bmatrix} 1 & \omega \\ 0 & 1 \end{bmatrix}$. Hence one can perform Dehn filling on the ends of $\mathbb{H}^3/\Gamma(I)$ so as to obtain a simply connected 3-manifold. $\square$

We will show that $\mathbb{H}^3/\Gamma(I)$ is in fact a link complement in S^3 (which we do not draw) as well as give explicitly a 4-component link complement $\tilde{\Omega}_{19}$ (Figure 10c), that is a 12-sheeted orbifold cover of Ω_{19} .

Let X denote Ω_{19} with its link circle put back (Dehn fill Ω_{19} with respect to the meridian $\begin{bmatrix} 1 & \omega \\ 0 & 1 \end{bmatrix}$). Note that X is topologically S^3 . Both of the above results follow from the fact that X admits a 6-fold irregular orbifold cover, X' , that is

FIGURE 7a. Construction of Ω'_5

homeomorphic to S^3 with singular locus a figure eight knot of cone angle π . The construction of X' is carried out in §3.4.1 below (see Figures 10d–10j).

The existence of X' implies that X is 60-fold orbifold covered by a nonsingular S^3 , for X' is 2-fold branch covered by the lens space $L(5, 2)$ which is in turn 5-fold covered by S^3 . Now, removing the link circle from X and its preimages from S^3 yields a 60-fold orbifold cover of Ω_{19} by a link complement which, we claim, corresponds to the group $\Gamma(I)$. Indeed, by construction the group Γ of the link complement is generated by conjugates of the matrix $\begin{bmatrix} 1 & \varphi \\ 0 & 1 \end{bmatrix}$ so that $\Gamma \subset \Gamma(I)$ and thus $\Gamma = \Gamma(I)$ since both are of index 60 in $\text{PSL}_2(\mathcal{O}_{19})$.

Actually constructing this link seemed rather difficult so we obtained the 4-component link complement $\tilde{\Omega}_{19}$ (Figure 10c) as follows. Removing the link circle from X and its preimages from X' yields a 6-fold irregular orbifold cover, Ω'_{19} , of Ω_{19} pictured in Figure 10a. Explicitly, one obtains Ω'_{19} by removing the link circle from X and performing the same cutting and pasting as in the construction of X'

FIGURE 7b. Construction of $\tilde{\Omega}_5$

given below. We omit the details. Now one easily obtains $\tilde{\Omega}_{19}$ as illustrated in Figures 10b–10c: first twist in the top component of the link in Ω'_{19} so as to unknot the singular circle, then take the appropriate 2-fold branched cover.

3.4.1. We conclude by constructing a 6-fold irregular orbifold cover, X' , of X homeomorphic to S^3 with singular locus a figure eight knot of cone angle π . This is done in steps 1–5 below (see Figures 10d–10j).

The orbifold X is drawn in Figure 10d. Remember that $\frac{n}{-}$ represents an edge of cone angle $2\pi/n$. We have added arrows to the edges of the singular locus to aid

FIGURE 8a.

FIGURE 8b.

in following the construction. The orbifold X' is obtained by cutting and pasting together 6 copies of X along disks which are bounded by loops in the singular locus of X .

1. Redraw X as in Figure 10e. In Figure 10f we have split S^3 open along a disk D bounded by a loop in the singular locus of X . Note that the segment $\overset{3}{\rightarrow}$ has been split into two segments of cone angle $\pi/3$, one on each copy of D .

2. Glue together two copies of X along D as shown in Figure 10g to obtain the cone manifold X_2 . Note that X_2 is not an orbifold cover of X , since the labels on its singular locus are not all integers.

FIGURE 8c.

3. Glue a copy of X to X_2 from the “back” of X_2 (cut and paste along disk E) to get X_3 as pictured in Figure 10h. Note cancellation of singular loci 3 and $3/2$ since $2\pi/3 + 2\pi/(3/2) = 2\pi$.

4. Now glue a copy of X to X_2 from the “front” (cut and paste along disk F) to get X'_3 as pictured in Figure 10i.

5. Finally, gluing X_3 to X'_3 (along disk G) gives the desired cover X' of X drawn in Figure 10j.

4. THE ORBIFOLDS Ω_m

The orbifold $\Omega_m = \mathbb{H}^3/\mathrm{PSL}_2(\mathcal{O}_m)$ is topologically a noncompact 3-manifold with cusps of the form $T^2 \times [0, \infty)$ for $m \neq 1, 3$ (resp. $S^2 \times [0, \infty)$ for $m = 1, 3$). In all cases, the number of cusps of Ω_m is equal to the class number of $\mathcal{O}_m$ (see [S]).

It is precisely for the 14 values of m in the list $\mathcal{L}$ that Ω_m embeds in S^3 . The orbifolds for $m = 5, 6, 15, 19, 23, 31, 39, 47, 71$ are drawn in Figure 1.

We obtained these embeddings as follows.

4.1. The cases $m = 5, 6, 15, 19, 23, 31$. Let $\Gamma_m = \mathrm{PSL}_2(\mathcal{O}_m)$ and D_m be a Ford fundamental polyhedron for the action of Γ_m on $\mathbb{H}^3$ (see [Sw] for details of this construction. The domains D_m , $m \leq 19$, are also explicitly given). R. Riley has written a computer program (described in [R2]) that computes the Ford domains

FIGURE 9.

FIGURE 10a.

and face pairings for $\mathrm{PSL}_2(\mathcal{O}_m)$ and $\mathrm{PGL}_2(\mathcal{O}_m)$. Copies of the program/domains are available from him upon request.

Let D_m^* be the compactification of D_m obtained by removing a horoball neighborhood about each of the cusps.

FIGURE 10b.

FIGURE 10c.

FIGURE 10d.

FIGURE 10e.

FIGURE 10f.

 FIGURE 10g. Construction of X_2

For our purposes, the key facts concerning D_m^* ($m \neq 1, 3$) are:

- i) D_m^*/Γ_m is topologically a compact 3-manifold with interior homeomorphic to $\Omega_m = D_m/\Gamma_m$ and boundary a disjoint union of tori.

FIGURE 10h. Construction of X_3 FIGURE 10i. Construction of X'_3 FIGURE 10j. Construction of X'

ii) Letting Γ_m^∞ denote the parabolic subgroup of Γ_m fixing $\{\infty\}$, then D_m^*/Γ_m^∞ is homeomorphic to $T^2 \times [0, 1]$ with $T^2 \times \{1\}$ the compactification of the cusp at $\{\infty\}$ and $T^2 \times \{0\}$ containing the singular locus of Ω_m as well as the remaining unpaired faces of D_m^* .

Thus the first step in obtaining Ω_m is to embed $D_m^*/\Gamma_m^\infty \cong T^2 \times [0, 1]$ in $S^3 = \mathbb{R}^3 \cup \{\infty\}$ so that:

1) $T^2 \times \{1\}$ is the boundary of an open tubular neighborhood of the circle, C , formed by the z -axis of $\mathbb{R}^3$ and the point $\{\infty\}$.

2) The loop on $T^2 \times \{1\}$ corresponding to the element $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ (resp. the element $\begin{bmatrix} 1 & \varphi \\ 0 & 1 \end{bmatrix}$) of Γ_m is parallel to C (resp. links C).

FIGURE 11. Embedding Ω_6

FIGURE 12. Obtaining Ω_6 from X_6

The next step is to identify the remaining faces on $T^2 \times \{0\}$ from the inside. As an example, the embedding of D_6^*/Γ_6 is illustrated in Figure 11.

4.2. The cases $m = 39, 47, 71$. After doing the above cases, we were motivated to study Allen Hatcher's work on $\mathrm{PGL}_2(\mathcal{O}_m)$ orbifolds, $X_m = \mathbb{H}^3/\mathrm{PGL}_2(\mathcal{O}_m)$ (see [H2]). Using Riley's $\mathrm{PGL}_2(\mathcal{O}_m)$ Ford domains, Hatcher draws the nineteen X_m that embed in S^3 . These orbifolds have cusps of the form $S^2 \times [0, \infty)$, and (for $m \neq 1, 3$) Ω_m is a 2-fold branched cover of X_m branched in such a way so that the cusp spheres of X_m are covered by cusp tori in Ω_m . Indeed, $\mathrm{PGL}_2(\mathcal{O}_m)/\mathrm{PSL}_2(\mathcal{O}_m) \cong \mathbb{Z}/2\mathbb{Z}$, and one obtains $\mathrm{PGL}_2(\mathcal{O}_m)$ from $\mathrm{PSL}_2(\mathcal{O}_m)$ by adjoining the involution $\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$. It is this involution and its conjugates that act on the cusp tori of Ω_m transforming them into cusp spheres.

In each case ($m \neq 1, 3$) one can find an unknotted circle, Σ , in X_m consisting of edges of cone angle π and two disjoint arcs on each cusp sphere such that Σ contains all singular edges with one or both vertices on cusp spheres. One obtains Ω_m as the 2-fold cover of X_m branched over Σ . In addition to obtaining Ω_m for $m = 39, 47, 71$ in this manner, we also constructed Ω_m for $m = 5, 6, 15, 19, 23, 31$ and found that they agreed with the Ω_m constructed in §4.1. We illustrate this method in Figure 12 by obtaining Ω_6 from X_6 . The reader can check that the orbifolds Ω_6 in Figures 11 and 12 are equivalent.

4.3. Modification of embeddings: Twist homeomorphisms. In certain cases, we have modified our embeddings of Ω_m in S^3 by:

- 1) Twisting about unknotted link components

- 2) Twisting about parallel link components

These twist homeomorphisms allow us to unknot singular circles as well as to link and unlink different components of a singular locus (see [Ro], Chapter 9, for a discussion of twist homeomorphisms).

The orbifolds Ω_m pictured in Figure 1 are embedded in a way that is most convenient for constructing the covers $\tilde{\Omega}_m$, and hence may differ from the embeddings appearing elsewhere.

5. VOLUME OF $\tilde{\Omega}_m$

A nice property of the Bianchi orbifolds is the existence of a number theoretical formula, due to Humbert, for their volume (see [T], Chapter 7):

$$\text{vol}(\Omega_m) = \frac{(D)^{3/2} \zeta_{\mathbb{Q}\sqrt{-m}}(2)}{24 \zeta_{\mathbb{Q}}(2)}, \quad D = \begin{cases} 4m, & m \equiv 1, 2 \pmod{4}, \\ m, & m \equiv 3 \pmod{4}. \end{cases}$$

Thus the volume of Ω_m and hence that of $\tilde{\Omega}_m$ can be computed to any degree of accuracy desired using the number theory program Pari ([P]).

As a check of our results, we used SnapPea ([W]) to analyze $\tilde{\Omega}_m$, Ω'_{39} and Ω_{71} in Figures 1–10 and found that their volumes agreed with those obtained from

Humbert's formula. We give these volumes below:

$$\text{vol}(\tilde{\Omega}_5) = 50.4476311137 \dots$$

$$\text{vol}(\tilde{\Omega}_6) = 62.1860747738 \dots$$

$$\text{vol}(\tilde{\Omega}_{15}) = 18.8316833668 \dots$$

$$\text{vol}(\tilde{\Omega}_{19}) = 31.8377775733 \dots$$

$$\text{vol}(\tilde{\Omega}_{23}) = 38.6951532243 \dots$$

$$\text{vol}(\tilde{\Omega}_{31}) = 54.0663025806 \dots$$

$$\text{vol}(\Omega''_{39}) = 82.787851850 \dots$$

(Ω''_{39} is the 2-fold branched cover of Ω'_{39} . Its singular locus is an unknotted circle of cone angle $2\pi/3$)

$$\text{vol}(\tilde{\Omega}_{47}) = 116.606541915 \dots$$

$$\text{vol}(\Omega_{71}) = 37.533306130 \dots$$

ACKNOWLEDGMENTS

I thank F. Bonahon for helpful correspondence which led me to consider the orbifolds Ω_m . I also thank A. Hatcher for providing me with his $\text{PGL}_2(\mathcal{O}_m)$ orbifold drawings which greatly facilitated my treatment of the cases $m = 39, 47, 71$.

REFERENCES

- [B] M. Baker, *Link complements and integer rings of class number greater than one*, Topology '90 (Proceedings of the research semester in Low Dimensional Topology at Ohio State University), Walter de Gruyter, Berlin, New York (1992), 55–59. MR **93k**:57007
- [G] GAP (Groups, algorithms, and programming). Available from: <http://www.gap.dcs.st-and.ac.uk/~gap/>.
- [H1] A. Hatcher, *Hyperbolic structures of arithmetic type on some link complements*, J. London Math. Soc. (2), **27** (1983), 345–355. MR **84m**:57005
- [H2] A. Hatcher, *Bianchi orbifolds of small discriminant*, preprint (available from author's web page).
- [P] Pari, Available from Université de Bordeaux: <http://www.math.u-bordeaux.fr/A2X>.
- [R1] R. Riley, *A quadratic parabolic group*, Math. Proc. Cambridge Philos. Soc. **77** (1975), 281–288. MR **54**:542
- [R2] R. Riley, *Applications of a computer implementation of Poincaré's theorem on fundamental polyhedra*, Math. Comp. **40** (1983), 607–632. MR **85b**:20064
- [Ro] D. Rolfsen, *Knots and Links*, Publish or Perish, Inc., Berkeley, (1976). MR **58**:24236; corrected reprint MR **95c**:57018
- [S] J.-P. Serre, *Le problème des groupes de congruence pour SL_2* , Ann. of Math. **92** (1972), 489–527. MR **42**:7671
- [Sw] R. Swan, *Generators and relations for certain special linear groups*, Adv. in Math. **6** (1971), 1–77. MR **44**:1741
- [T] W. Thurston, *The geometry and topology of 3-manifolds*, Mimeographed notes, Princeton University, 1978.
- [V] K. Vogtmann, *Rational homology of Bianchi groups*, Math. Ann. **272** (1985), 399–419. MR **87a**:22025
- [W] J. Weeks, SnapPea 2.4 PPC. Available from: weeks@geom.umn.edu.
- [Wi] N. Wielenberg, *The structure of certain subgroups of the Picard group*, Math. Proc. Cambridge Philos. Soc. **84** (1978), 427–436. MR **80b**:57010