SPIN STRUCTURES AND CODIMENSION TWO EMBEDDINGS OF 3-MANIFOLDS UP TO REGULAR HOMOTOPY

OSAMU SAEKI AND MASAMICHI TAKASE

ABSTRACT. We clarify the structure of the set of regular homotopy classes containing embeddings of a 3-manifold into 5-space inside the set of all regular homotopy classes of immersions with trivial normal bundles. As a consequence, we show that for a large class of 3-manifolds M^3 , the following phenomenon occurs: there exists a codimension two immersion of the 3-sphere whose double points cannot be eliminated by regular homotopy, but can be eliminated after taking the connected sum with a codimension two embedding of M^3 . This involves introducing and studying an equivalence relation on the set of spin structures on M^3 . Their associated μ -invariants also play an important role.

1. INTRODUCTION

Let $f: M^3 \looparrowright \mathbf{R}^4$ be an immersion of an oriented 3-manifold M^3 into oriented $\mathbf{R}^4$. Then, via the bundle isomorphism $TM^3 \oplus \varepsilon^1 \cong f^*T\mathbf{R}^4$ (ε^1 being the trivial line bundle), the unique spin structure on $\mathbf{R}^4$ induces a spin structure on M^3 that is clearly a regular homotopy invariant of f .

Now let $F: M^3 \looparrowright \mathbf{R}^5$ be an immersion with trivial normal bundle. By taking a normal framing for F , we obtain a spin structure on M^3 that usually depends on the choice of the normal framing. Then, as has been seen in [9, Section 3], we are naturally led to an equivalence relation on the set of spin structures on M^3 , regarded as an affine space over $H^1(M^3; \mathbf{Z}_2)$. We say that two spin structures are *equivalent modulo* $\text{Im } \rho$ if their difference lies in the image of the modulo two reduction map $\rho: H^1(M^3; \mathbf{Z}) \rightarrow H^1(M^3; \mathbf{Z}_2)$. The spin structure modulo $\text{Im } \rho$ associated to F is independent of the choice of the normal framing and is a regular homotopy invariant of F .

In [9], a geometric characterisation of regular homotopy classes of immersions of M^3 into $\mathbf{R}^5$ has also been given. As a consequence, we have encountered a rather interesting situation for embeddings of the 3-torus T^3 up to regular homotopy: *there exists an immersion $g: S^3 \looparrowright \mathbf{R}^5$ such that (1) g is not regularly homotopic to an embedding $S^3 \hookrightarrow \mathbf{R}^5$, but (2) for any embedding $E: T^3 \hookrightarrow \mathbf{R}^5$, the connected sum $E \sharp g: T^3 \looparrowright \mathbf{R}^5$ is again regularly homotopic to an embedding $T^3 \hookrightarrow \mathbf{R}^5$.* In other words, the double points of the immersion g cannot be eliminated by regular homotopy, but one can eliminate them after taking the connected sum with an embedding of T^3 . We call such an immersion $g: S^3 \looparrowright \mathbf{R}^5$ a *T^3 -pseudo-embedding* of S^3 (see Definition 3.1). This phenomenon is closely related to the diversity of

Received by the editors May 25, 2001.

2000 *Mathematics Subject Classification.* Primary 57R42, 57M50; Secondary 57R40, 57M27.

The first author was partially supported by Grant-in-Aid for Scientific Research No. 13640076, Ministry of Education, Science and Culture, Japan.

spin structures of T^3 , and, in the proof given in [9], spin structures modulo $\text{Im } \rho$ associated to embeddings of T^3 play an important role.

In this paper, we prove the existence of M^3 -pseudo-embeddings $S^3 \looparrowright \mathbf{R}^5$ for a large class of 3-manifolds M^3 (see Proposition 3.2 and Theorem 3.7). This involves a study of spin structures modulo $\text{Im } \rho$ on M^3 , and of their associated μ -invariants (see Section 3). In Section 4, we study how the regular homotopy classes of embeddings are distributed in the set of all regular homotopy classes of immersions, and give a result which implies that the cases of S^3 and T^3 are typical (Theorem 4.2). We also give an example of a 3-manifold for which the classes of embeddings are distributed in a complicated manner (Proposition 4.4). In Section 5, we generalise the above study, replacing S^3 by an arbitrary 3-manifold N^3 . Finally, in the last section, we introduce the notion of virtual homotopy: we say that two immersions are *virtually homotopic* if they are regularly homotopic after taking connected sums with embeddings. Virtual homotopy classes of immersions of closed connected oriented 3-manifolds into $\mathbf{R}^5$ with trivial normal bundles form a group under connected sum. We show that this group is isomorphic to $\mathbf{Z}/12\mathbf{Z}$.

Throughout the paper, manifolds are of class C^∞ and M^3 is a closed oriented 3-manifold. We write “ $A \approx B$ ” if there is a bijective correspondence between the sets A and B ; we use the symbol “ $\cong$ ” for a group isomorphism. The symbol “ $\sim_r$ ” means “regularly homotopic”. We often do not distinguish between an immersion f and its regular homotopy class, which we also denote by f .

The authors are extremely grateful to András Szűcs for his useful advice during the preparation of this paper. In particular, the term M^3 -pseudo-embedding is due to him, and Section 5 has been motivated by his question.

2. PRELIMINARIES

In this section, we recall some known results. We review them only in our case — that of immersions of 3-manifolds into $\mathbf{R}^5$ — although some of them were originally stated in more general contexts.

2.1. Immersions of the 3-sphere. Let $\text{Imm}[X, Y]$ denote the set of regular homotopy classes of immersions of a manifold X into a manifold Y . Denote by $\text{Emb}[X, Y]$ the subset of regular homotopy classes containing an embedding. Note that the set $\text{Imm}[S^3, \mathbf{R}^5]$ has a group structure given by the connected sum, and that the Smale invariant

$$\Omega: \text{Imm}[S^3, \mathbf{R}^5] \rightarrow \mathbf{Z}$$

gives a group isomorphism (see [6, 3]). Hughes and Melvin [4] have shown that $\text{Emb}[S^3, \mathbf{R}^5]$ forms the subgroup isomorphic to $24\mathbf{Z}$. Furthermore, for an embedding $f: S^3 \hookrightarrow \mathbf{R}^5$ we have $\Omega(f) = 3\sigma(W_f^4)/2 \in 24\mathbf{Z}$, where W_f^4 is a Seifert surface for f and $\sigma(W_f^4)$ is its signature.

This last result has been extended by Ekholm and Szűcs [1] as follows. For an immersion $f: S^3 \looparrowright \mathbf{R}^5$, consider a *singular Seifert surface* for f , that is, a generic map $\tilde{f}: W_f^4 \rightarrow \mathbf{R}^5$ that is bounded by f and has no singularity near the boundary $S^3 = \partial W_f^4 \subset W_f^4$. If we denote by $\#\Sigma^{1,1}(\tilde{f})$ the algebraic number of cusp points of $\tilde{f}$, then

$$\Omega(f) = \frac{1}{2}(3\sigma(W_f^4) + \#\Sigma^{1,1}(\tilde{f})).$$

Note that for an embedding we can consider a usual nonsingular Seifert surface and obtain the result of Hughes and Melvin as a corollary.

2.2. Immersions of 3-manifolds with trivial normal bundles. Let M^3 be a closed oriented 3-manifold. Denote by $\text{Imm}[M^3, \mathbf{R}^5]_0$ the subset of $\text{Imm}[M^3, \mathbf{R}^5]$ consisting of the regular homotopy classes of immersions with trivial normal bundles. Note that $\text{Emb}[M^3, \mathbf{R}^5] \subset \text{Imm}[M^3, \mathbf{R}^5]_0$. In this subsection, we recall the results in [9], which give a geometric description of the set $\text{Imm}[M^3, \mathbf{R}^5]_0$.

First, we introduce an equivalence relation on the set of spin structures on an oriented manifold.

Definition 2.1 (Spin structures modulo $\text{Im } \rho$). Denote by $\text{Spin}(M^n)$ the set of all spin structures on an oriented n -manifold M^n . Recall that $\text{Spin}(M^n)$ is an affine space over $H^1(M^n; \mathbf{Z}_2)$. Let

$$\rho: H^1(M^n; \mathbf{Z}) \rightarrow H^1(M^n; \mathbf{Z}_2)$$

be the modulo two reduction.

Two spin structures ω and $\omega' \in \text{Spin}(M^n)$ are said to be *equivalent modulo $\text{Im } \rho$* if their difference lies in $\text{Im } \rho \subset H^1(M^n; \mathbf{Z}_2)$. We denote the equivalence class containing ω by $[\omega]$ and call it a *spin structure modulo $\text{Im } \rho$* . Let

$$\begin{array}{ccc} [\cdot]: & \text{Spin}(M^n) & \rightarrow \text{Spin}(M^n)/\text{Im } \rho \\ & \omega & \mapsto [\omega] \end{array}$$

be the natural projection. Note that $\text{Spin}(M^n)/\text{Im } \rho \approx H^1(M^n; \mathbf{Z}_2)/\text{Im } \rho$ (non-canonically).

As has been shown in [9], an element in the set $\text{Imm}[M^3, \mathbf{R}^5]_0$ is characterised by two invariants c and i defined as follows.

Let $F: M^3 \looparrowright \mathbf{R}^5$ be an immersion with trivial normal bundle. If we take a normal framing for F , then the unique spin structure on $T\mathbf{R}^5$ induces a spin structure ω_F of M^3 . If we take another normal framing, then we get another spin structure ω'_F of M^3 . We have shown in [9, Section 3] that the difference of two such spin structures always lies in $\text{Im } \rho \subset H^1(M^3; \mathbf{Z}_2)$. Thus to F we can associate $[\omega_F]$ ($= [\omega'_F]$) — the spin structure modulo $\text{Im } \rho$ — which is a regular homotopy invariant of F . Let

$$c: \text{Imm}[M^3, \mathbf{R}^5]_0 \rightarrow \text{Spin}(M^3)/\text{Im } \rho$$

be this correspondence.

The second invariant $i: \text{Imm}[M^3, \mathbf{R}^5]_0 \rightarrow \mathbf{Z}$ is an analogue of a geometric formula for the Smale invariant given in [1]. Denote by $\tau H^1(M^3; \mathbf{Z})$ the torsion subgroup of $H^1(M^3; \mathbf{Z})$, and let $\alpha = \alpha(M^3)$ be the dimension of $\tau H^1(M^3; \mathbf{Z}) \otimes \mathbf{Z}_2$ over $\mathbf{Z}_2$. Note that the number of elements in the set $\text{Spin}(M^3)/\text{Im } \rho$ is equal to 2^α . Then, for an immersion $F: M^3 \looparrowright \mathbf{R}^5$ with trivial normal bundle, we define

$$i(F) := \frac{1}{2}(3(\sigma(W_F^4) - \alpha(M^3)) + \#\Sigma^{1,1}(\tilde{F})),$$

where $\tilde{F}: W_F^4 \rightarrow \mathbf{R}^5$ is a singular Seifert surface for F and $\#\Sigma^{1,1}(\tilde{F})$ is the algebraic number of cusp points of $\tilde{F}$. Note that $i(F)$ is always an integer.

The following results were obtained in [9]. Put $\text{Imm}[M^3, \mathbf{R}^5]_0^C := c^{-1}(C)$ for $C \in \text{Spin}(M^3)/\text{Im } \rho$.

- (a) For any $[\omega] \in \text{Spin}(M^3)/\text{Im } \rho$, $\text{Imm}[M^3, \mathbf{R}^5]_0^{[\omega]}$ contains a class represented by an embedding.

- (b) The map $(c, i): \text{Imm}[M^3, \mathbf{R}^5]_0 \rightarrow (\text{Spin}(M^3)/\text{Im } \rho) \times \mathbf{Z}$ gives a bijection. As a corollary, two embeddings in $\text{Imm}[M^3, \mathbf{R}^5]_0^{[\omega]} \approx \mathbf{Z}$ ($[\omega] \in \text{Spin}(M^3)/\text{Im } \rho$) are regularly homotopic if and only if they have Seifert surfaces with the same signature.
- (c) By taking connected sums of immersions, we can define an action of the group $\text{Imm}[S^3, \mathbf{R}^5]$ on the set $\text{Imm}[M^3, \mathbf{R}^5]_0$. This action is effective, and each orbit coincides with $\text{Imm}[M^3, \mathbf{R}^5]_0^{[\omega]}$ for some $[\omega] \in \text{Spin}(M^3)/\text{Im } \rho$.

Remark 2.2. The above items (a) and (b) imply that each “ $\mathbf{Z}$ -component” of

$$(2.1) \quad \text{Imm}[M^3, \mathbf{R}^5]_0 \approx (\text{Spin}(M^3)/\text{Im } \rho) \times \mathbf{Z} = \mathbf{Z} \amalg \cdots \amalg \mathbf{Z}$$

contains a class represented by an embedding. Furthermore, (c) implies that if we take an embedding F in $\text{Imm}[M^3, \mathbf{R}^5]_0^{[\omega]} \approx \mathbf{Z}$, then we can define the bijection

$$\begin{array}{ccc} \sharp_F: \text{Imm}[S^3, \mathbf{R}^5] & \rightarrow & \text{Imm}[M^3, \mathbf{R}^5]_0^{c(F)} \\ g & \mapsto & F \sharp g. \end{array}$$

From now on, we always consider each $\mathbf{Z}$ -component of (2.1) to be endowed with such a group structure, induced from that of $\text{Imm}[S^3, \mathbf{R}^5]$. In other words, whenever we mention a $\mathbf{Z}$ -component of (2.1), we fix in it an embedding F , which determines the group structure via the bijection $\sharp_F$ above.

Remark 2.3. According to Wu’s computation [11] of $\text{Imm}[M^3, \mathbf{R}^5]$, the set $\text{Imm}[M^3, \mathbf{R}^5]_0$ corresponds bijectively to $\Gamma_2(M^3) \times \mathbf{Z}$, where $\Gamma_2(M^3)$ denotes the set of order two elements of $H^2(M^3; \mathbf{Z})$ together with the zero element. The Bockstein homomorphism $H^1(M^3; \mathbf{Z}_2) \rightarrow H^2(M^3; \mathbf{Z})$ induces a bijection between $H^1(M^3; \mathbf{Z}_2)/\text{Im } \rho$ and $\Gamma_2(M^3)$ (see [9, Remark 3.6]).

3. M^3 -PSEUDO-EMBEDDINGS OF THE 3-SPHERE

As mentioned in the introduction, we have observed in [9, Section 6] a surprising situation for embeddings of T^3 up to regular homotopy. We first give a definition to describe this phenomenon.

Definition 3.1. Let M^n and N^n be closed oriented n -manifolds and $g: N^n \looparrowright \mathbf{R}^{n+k}$ be an immersion. We say that g is an M^n -pseudo-embedding if (1) g is not regularly homotopic to any embedding, and (2) for some embedding $F: M^n \hookrightarrow \mathbf{R}^{n+k}$, the connected sum $F \sharp g: M^n \sharp N^n \looparrowright \mathbf{R}^{n+k}$ is regularly homotopic to an embedding.

In this paper, we consider only the case of codimension two immersions of 3-manifolds, i.e., $n = 3$ and $k = 2$.

Under this notion of “ M^n -pseudo-embedding”, the curious phenomenon mentioned in the introduction (see [9, Section 6]) can be described as follows: there exists a T^3 -pseudo-embedding $h: S^3 \looparrowright \mathbf{R}^5$. Here we show that for a large class of 3-manifolds M^3 , there exists an M^3 -pseudo-embedding $S^3 \looparrowright \mathbf{R}^5$.

The following is a necessary and sufficient condition on spin structures of M^3 for the existence of an M^3 -pseudo-embedding $S^3 \looparrowright \mathbf{R}^5$. Let M^3 be a closed oriented 3-manifold. If ω is a spin structure on M^3 , then we denote by $\mu(M^3, \omega)$ the μ -invariant (or Rohlin invariant) of the spin manifold (M^3, ω) , i.e., the signature modulo 16 of a spin 4-manifold that is spin bounded by (M^3, ω) (for example, see [7]).

Proposition 3.2. *There exists an M^3 -pseudo-embedding $S^3 \looparrowright \mathbf{R}^5$ if and only if M^3 admits two spin structures ω and ω' such that*

- (1) $[\omega] = [\omega']$, i.e., $\omega - \omega' \in \text{Im } \rho \subset H^1(M^3; \mathbf{Z}_2)$, and
- (2) $\mu(M^3, \omega) \neq \mu(M^3, \omega')$.

Proof. Suppose that M^3 has two spin structures ω and ω' satisfying the two conditions (1) and (2). There exist compact spin 4-manifolds W^4 and W'^4 that are spin bounded by (M^3, ω) and (M^3, ω') , respectively, such that each of them has a handlebody decomposition with one 0-handle and some 2-handles with even framings (see [5]). The condition $\mu(M^3, \omega) \neq \mu(M^3, \omega')$ implies that $\sigma(W^4) \not\equiv \sigma(W'^4) \pmod{16}$. Furthermore, by the same argument as in [9, Section 6], we can take embeddings $\tilde{F}: W^4 \hookrightarrow \mathbf{R}^5$ and $\tilde{G}: W'^4 \hookrightarrow \mathbf{R}^5$, and then their restrictions $F := \tilde{F}|_{M^3}$ and $G := \tilde{G}|_{M^3}$ to the boundaries have the same associated spin structure modulo $\text{Im } \rho$, that is, $c(F) = [\omega] = [\omega'] = c(G)$.

Then, from the fact that

$$\sharp_F: \text{Imm}[S^3, \mathbf{R}^5] \longrightarrow \text{Imm}[M^3, \mathbf{R}^5]_0^{[\omega]}$$

is a bijection (see Remark 2.2), there exists an immersion $g: S^3 \looparrowright \mathbf{R}^5$ with $F\sharp g \sim_r G$.

Let us assume that g is regularly homotopic to an embedding. Then, by the result of Hughes and Melvin [4], g extends to an immersion of a compact oriented 4-manifold V^4 with signature $16k$ ($k \in \mathbf{Z}$). Hence,

$$i(F\sharp g) = \frac{3}{2}(\sigma(W^4 \natural V^4) - \alpha(M^3)) = \frac{3}{2}(\sigma(W^4) - \alpha(M^3)) + 24k,$$

where $W^4 \natural V^4$ stands for the boundary connected sum of W^4 and V^4 . This contradicts the fact that $F\sharp g \sim_r G$, since

$$i(G) = \frac{3}{2}(\sigma(W'^4) - \alpha(M^3))$$

and $\sigma(W^4) \not\equiv \sigma(W'^4) \pmod{16}$.

Conversely, if there is an M^3 -pseudo-embedding $g: S^3 \looparrowright \mathbf{R}^5$, then we have two embeddings F and $G: M^3 \hookrightarrow \mathbf{R}^5$ such that $F\sharp g \sim_r G$. Denote by ω_F and ω_G the spin structures determined by F and G respectively. Then, $[\omega_F] = c(F) = c(G) = [\omega_G]$, since F and G restricted to $M^3 \setminus \text{Int } D^3$ are regularly homotopic (see [9, Section 3]), where $\text{Int } D^3$ denotes the interior of a 3-disc embedded in M^3 . We also see from (b) in Subsection 2.2 that $\mu(M^3, \omega_F) \neq \mu(M^3, \omega_G)$, since $i(g) = i(G) - i(F)$ and g is not regularly homotopic to an embedding.

This completes the proof. $\square$

Example 3.3. Let T^3 be the 3-torus. Since all the classes in $H^1(M^3; \mathbf{Z}_2) \cong \mathbf{Z}_2 \oplus \mathbf{Z}_2 \oplus \mathbf{Z}_2$ are modulo two reductions of integral classes, we see that $\text{Spin}(M^3)/\text{Im } \rho = 0$ and $\text{Imm}[T^3, \mathbf{R}^5]_0 \approx \mathbf{Z}$.

The 3-torus T^3 has eight distinct spin structures, which all belong to the unique equivalence class in $\text{Spin}(M^3)/\text{Im } \rho$. It is well known that seven of them are spin bounded by a spin 4-manifold with signature 0 modulo 16 and the remaining one is spin bounded by a spin 4-manifold with signature 8 modulo 16.

Thus, T^3 satisfies the conditions in Proposition 3.2. Actually, $\text{Emb}[T^3, \mathbf{R}^5]$ forms the subgroup isomorphic to $12\mathbf{Z}$ of $\text{Imm}[T^3, \mathbf{R}^5]_0 \approx \mathbf{Z}$. Then, an immersion $g: S^3 \looparrowright \mathbf{R}^5$ with Smale invariant $12 + 24k$ ($k \in \mathbf{Z}$) is not regularly homotopic to any embedding, but the connected sum of g and an arbitrary embedding $T^3 \hookrightarrow \mathbf{R}^5$ is regularly homotopic to an embedding (for details, see [9, Section 6]).

The following theorem, due to Kaplan [5], together with Lemma 3.5 below, provides some 3-manifolds with two spin structures that satisfy the two conditions in Proposition 3.2.

Theorem 3.4 (Kaplan [5, Theorem 6.12]). *Suppose that M^3 is a closed connected oriented 3-manifold that bounds a spin 4-manifold of signature k . Suppose further that there exist elements $x_i \in H^1(M^3; \mathbf{Z})$, $i = 1, 2, 3$, such that $x_1 \smile x_2 \smile x_3$ is an odd multiple of the generator of $H^3(M^3; \mathbf{Z})$. Then M^3 bounds a spin 4-manifold of signature $k + 8$. $\square$*

Actually, in the proof given in [5] of the above theorem, two spin structures of M^3 for which the μ -invariants differ by 8 are specified. We have:

Lemma 3.5. *These two spin structures satisfy the condition (1) in Proposition 3.2.*

Before proving this lemma, we need a definition.

Definition 3.6. Let $L = \{K_1, K_2, \dots, K_n\}$ be a framed link in S^3 . A sublink $L' \subset L$ is said to be *characteristic*, if $lk(L', K_i) \equiv lk(K_i, K_i) \pmod{2}$ for $1 \leq i \leq n$.

Proof of Lemma 3.5. Consider $M^3 = M_L^3$ as the boundary of the 4-manifold W_L^4 for a framed link $L = \{K_1, K_2, \dots, K_n\}$ with even framings, that is, the boundary of the spin 4-manifold obtained by attaching a 2-handle to D^4 along each component of $L \subset S^3 = \partial D^4$ with the given framing.

Then, characteristic sublinks L' of L bijectively correspond to spin structures $\omega_{L'}$ of M^3 . Note that each component K_i of L corresponds to an element $[K_i] \in H_2(W_L^4; \mathbf{Z})$, which is represented by the union of the core of the handle over K_i and the cone on K_i in D^4 . Note also that the classes $[K_i]$ form a basis of $H_2(W_L^4; \mathbf{Z})$ that is free abelian. Furthermore, each characteristic sublink L' represents the Poincaré dual of the relative obstruction to extending the spin structure $\omega_{L'}$ over W_L^4 , $w_2(W_L^4, \omega_{L'}) \in H^2(W_L^4, M^3; \mathbf{Z}_2) \cong H_2(W_L^4; \mathbf{Z}_2)$, which maps to the second Stiefel-Whitney class $w_2(W_L^4)$ of W_L^4 under the map

$$j^*: H^2(W_L^4, M^3; \mathbf{Z}_2) \rightarrow H^2(W_L^4; \mathbf{Z}_2),$$

where $j: W_L^4 \rightarrow (W_L^4, M^3)$ is the inclusion (see the diagram below). Obviously, the empty (characteristic) sublink corresponds to the spin structure that extends over W_L^4 .

Therefore, in order to obtain Theorem 3.4, we need to find a characteristic sublink corresponding to a spin structure that provides the μ -invariant $\sigma(W_L^4) + 8 \pmod{16}$. In fact, in the proof of Theorem 3.4, Kaplan [5] has found in a suitable L such a characteristic sublink ℓ of one component with the property that $[\ell] \cdot x = 0$ for all $x \in H_2(W_L^4; \mathbf{Z})$. This property of $[\ell]$ implies that $[\ell]$ lies in the kernel of the map $j_*: H_2(W_L^4; \mathbf{Z}) \rightarrow H_2(W_L^4, M^3; \mathbf{Z})$ (see the diagram below), and hence it is the image of a class (ℓ) in $H_2(M^3; \mathbf{Z})$ under the homomorphism $i_*: H_2(M^3; \mathbf{Z}) \rightarrow H_2(W_L^4; \mathbf{Z})$ induced by the inclusion i .

Now let us consider the following diagram, in which the horizontal arrows come from the exact sequences for the pair (W_L^4, M^3) , each vertical arrow upstairs comes from Poincaré-Lefschetz duality, and each vertical arrow downstairs is the modulo

two reduction:

$$\begin{array}{ccccc}
 H_2(M^3; \mathbf{Z}) & \xrightarrow{i_*} & H_2(W_L^4; \mathbf{Z}) & \xrightarrow{j_*} & H_2(W_L^4, M^3; \mathbf{Z}) \\
 \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\
 H^1(M^3; \mathbf{Z}) & \longrightarrow & H^2(W_L^4, M^3; \mathbf{Z}) & \longrightarrow & H^2(W_L^4; \mathbf{Z}) \\
 \downarrow \rho & & \downarrow & & \downarrow \\
 H^1(M^3; \mathbf{Z}_2) & \longrightarrow & H^2(W_L^4, M^3; \mathbf{Z}_2) & \xrightarrow{j^*} & H^2(W_L^4; \mathbf{Z}_2) \\
 \omega_\ell & \longmapsto & w_2(W_L^4, \omega_\ell) & \longmapsto & w_2(W_L^4) = 0.
 \end{array}$$

Then we see that the difference between the two spin structures corresponding to the characteristic sublink ℓ and to the empty sublink corresponds to $\rho((\ell)^*)$, where $(\ell)^* \in H^1(M^3; \mathbf{Z})$ is the Poincaré dual of (ℓ) . This completes the proof. $\square$

Thus Theorem 3.4, together with Lemma 3.5, implies that a large class of 3-manifolds exhibit the same phenomenon as the 3-torus. Namely, we have the following theorem.

Theorem 3.7. *If there exist elements $x_i \in H^1(M^3; \mathbf{Z})$, $i = 1, 2, 3$, such that $x_1 \smile x_2 \smile x_3$ is an odd multiple of the generator of $H^3(M^3; \mathbf{Z})$, then there exists an M^3 -pseudo-embedding $S^3 \looparrowright \mathbf{R}^5$.* $\square$

We see easily that $S^1 \times F^2$ is such a 3-manifold satisfying the conditions of the theorem above, where F^2 is a closed connected orientable surface of positive genus.

4. REGULAR HOMOTOPY CLASSES OF EMBEDDINGS

In all the cases that we have dealt with, $\text{Emb}[M^3, \mathbf{R}^5]$ forms the subgroup $24\mathbf{Z}$ (for S^3 or $\mathbf{Z}_2$ -homology 3-spheres [4, 10]) or $12\mathbf{Z}$ (for T^3 or $S^1 \times F^2$) of a $\mathbf{Z}$ -component in $\text{Imm}[M^3, \mathbf{R}^5]_0$. The following proposition implies that these are all the possibilities.

Proposition 4.1. *Let ω and ω' be spin structures of M^3 such that $[\omega] = [\omega']$, i.e., $\omega - \omega' \in \text{Im } \rho \subset H^1(M^3; \mathbf{Z}_2)$. Then,*

$$\mu(M^3, \omega) \equiv \mu(M^3, \omega') \pmod{8}.$$

Proof. Let W^4 and W'^4 be compact spin 4-manifolds spin bounded by (M^3, ω) and (M^3, ω') , respectively, and put $V^4 := W^4 \cup -W'^4$. By the assumption, we can take an oriented surface $F \subset M^3$ with $[F] \in H_2(M^3; \mathbf{Z})$ being an integral dual to $\omega - \omega' \in H^1(M^3; \mathbf{Z}_2)$. Since $[F] \in H_2(V^4; \mathbf{Z})$ is also an integral dual to the second Stiefel-Whitney class $w_2(V^4) \in H^2(V^4; \mathbf{Z}_2)$, considering that F lies in $M^3 \subset V^4$, we have

$$\sigma(V^4) \equiv [F] \cdot [F] \equiv 0 \pmod{8},$$

where “ $\cdot$ ” stands for the intersection number in V^4 (see [7, p. 25, Lemma 3.4], for example). Thus by Novikov additivity, we have

$$\mu(M^3, \omega) - \mu(M^3, \omega') \equiv \sigma(W^4) - \sigma(W'^4) \equiv \sigma(V^4) \equiv 0 \pmod{8}.$$

This completes the proof. $\square$

Theorem 4.2. *In each $\mathbf{Z}$ -component of $\text{Imm}[M^3, \mathbf{R}^5]_0 \approx \mathbf{Z} \amalg \cdots \amalg \mathbf{Z}$, $\text{Emb}[M^3, \mathbf{R}^5]$ is a subgroup isomorphic either to $24\mathbf{Z}$ or to $12\mathbf{Z}$.*

Proof. In view of the result of Hughes and Melvin and (a), (c) in Subsection 2.2, we see that $\text{Emb}[M^3, \mathbf{R}^5]$ contains the subgroup $24\mathbf{Z}$ in each $\mathbf{Z}$ -component of $\text{Imm}[M^3, \mathbf{R}^5]_0 \approx \mathbf{Z} \amalg \cdots \amalg \mathbf{Z}$.

Suppose that two embeddings F and G : $M^3 \hookrightarrow \mathbf{R}^5$ belong to the same $\mathbf{Z}$ -component of $\text{Imm}[M^3, \mathbf{R}^5]_0 \approx \mathbf{Z} \amalg \cdots \amalg \mathbf{Z}$. Let W_F^4 and W_G^4 be Seifert surfaces for F and G respectively. Then, we can consider spin structures ω_F and ω_G for M^3 induced from the normal framings given by $M^3 \subset W_F^4$ and $M^3 \subset W_G^4$ respectively. Since $c(F) = c(G)$, we have $[\omega_F] = [\omega_G]$. Furthermore, it is clear that $\mu(M^3, \omega_F) \equiv \sigma(W_F^4) \pmod{16}$ and $\mu(M^3, \omega_G) \equiv \sigma(W_G^4) \pmod{16}$. Therefore, by Proposition 4.1, we have $\sigma(W_F^4) - \sigma(W_G^4) \in 8\mathbf{Z}$. Then, by the definitions of the invariant i and the group structure on each $\mathbf{Z}$ -component, we have the desired result. $\square$

Remark 4.3. By the above proof, we see that all the spin structures in a class $C \in \text{Spin}(M^3)/\text{Im } \rho$ give the same μ -invariant if and only if $\text{Emb}[M^3, \mathbf{R}^5]$ forms the subgroup $24\mathbf{Z}$ in $\text{Imm}[M^3, \mathbf{R}^5]_0^C \approx \mathbf{Z}$, and that a class $C \in \text{Spin}(M^3)/\text{Im } \rho$ contains two spin structures that give different μ -invariants if and only if $\text{Emb}[M^3, \mathbf{R}^5]$ forms the subgroup $12\mathbf{Z}$ in $\text{Imm}[M^3, \mathbf{R}^5]_0^C \approx \mathbf{Z}$.

Thus, by Proposition 3.2, we see that there exists an M^3 -pseudo-embedding $S^3 \looparrowright \mathbf{R}^5$ if and only if $\text{Emb}[M^3, \mathbf{R}^5]$ forms the subgroup $12\mathbf{Z}$ in a $\mathbf{Z}$ -component of $\text{Imm}[M^3, \mathbf{R}^5]_0$.

In view of Theorem 4.2, it is natural to ask: given a 3-manifold M^3 , does $\text{Emb}[M^3, \mathbf{R}^5]$ necessarily form the subgroup $24\mathbf{Z}$ (or $12\mathbf{Z}$) simultaneously in all the $\mathbf{Z}$ -components of $\text{Imm}[M^3, \mathbf{R}^5]_0 \approx \mathbf{Z} \amalg \cdots \amalg \mathbf{Z}$? We give an example that answers this question negatively, as follows.

Proposition 4.4. *Let X^3 be the (total space of the) S^1 -bundle over $T^2 = S^1 \times S^1$ with Euler class 2. Then, $\text{Imm}[X^3, \mathbf{R}^5]_0 \approx \mathbf{Z} \amalg \mathbf{Z}$, and $\text{Emb}[X^3, \mathbf{R}^5]$ forms the subset $12\mathbf{Z} \amalg 24\mathbf{Z} \subset \mathbf{Z} \amalg \mathbf{Z}$.*

Proof. We see that $H^2(X^3; \mathbf{Z}) \cong \mathbf{Z} \oplus \mathbf{Z} \oplus \mathbf{Z}_2$ and $H^1(X^3; \mathbf{Z}) \cong \mathbf{Z} \oplus \mathbf{Z}$, which can be computed using the Gysin exact sequence for the S^1 -bundle $X^3 \rightarrow T^2$, for example. Therefore, we have $H^1(X^3; \mathbf{Z}_2) \cong \mathbf{Z}_2 \oplus \mathbf{Z}_2 \oplus \mathbf{Z}_2$. Since two of the three $\mathbf{Z}_2$ factors come from integral classes, we have $H^1(X^3; \mathbf{Z}_2)/\text{Im } \rho \cong \mathbf{Z}_2$. Thus, we have $\text{Spin}(X^3)/\text{Im } \rho \approx \mathbf{Z}_2$, and hence $\text{Imm}[X^3, \mathbf{R}^5]_0 \approx \mathbf{Z} \amalg \mathbf{Z}$. We want to show that in one equivalence class in $\text{Spin}(X^3)/\text{Im } \rho \approx \mathbf{Z}_2$, all spin structures provide the same μ -invariant and that in the other class there are spin structures that provide distinct μ -invariants.

It is known (see [2, p. 198] or [8, p. 714], for example) that X^3 has a framed link representation given by the Borromean rings $L = \{K_1, K_2, K_3\}$ with framings $(0, 0, 2)$ (see Figure 1). This framed link has eight distinct characteristic sublinks — $\emptyset, K_1, K_2, K_3, \{K_1, K_2\}, \{K_2, K_3\}, \{K_3, K_1\}$ and $L = \{K_1, K_2, K_3\}$ — each of which corresponds to a spin structure of X^3 .

Now our strategy is to apply, for each characteristic sublink of our link L , Theorem 4.2 (ii) in [5], which enables us to compute the signature of a spin 4-manifold corresponding to each characteristic sublink of L without actually constructing such a 4-manifold. First, the 4-manifold W_L^4 represented by L is nothing but a spin 4-manifold corresponding to the empty characteristic sublink. Put $k := \sigma(W_L^4)$.

FIGURE 1. A framed link representation of X^3

For characteristic sublinks with one component, we can just apply [5, Theorem 4.2 (ii)]. Then the signature of the spin 4-manifold is $k \bmod 16$ for K_1 and K_2 , and $k - 2 \bmod 16$ for K_3 .

For characteristic sublinks with two components, we first follow the procedure of Case 4 in the proof of Theorem 3.1 in [5], that is, we add a push-off of one component to the other in order to obtain a new characteristic sublink with one component. This move is the so-called handle sliding and does not change the 3-manifold X^3 and the corresponding spin structure. Now for any two components, we get an unknotted component whose framing is 0 when the original characteristic sublink does not contain K_3 , and is 2 when it does contain K_3 . Therefore, applying [5, Theorem 4.2 (ii)], we see that the signature of the spin 4-manifold is $k \bmod 16$ in the former case and $k - 2 \bmod 16$ in the latter case.

When we consider L itself as a characteristic sublink, we first follow the procedure of Case 4 in the proof of Theorem 3.1 in [5] as well. If we add push-offs of K_1 and K_2 to K_3 , then the result is a trefoil knot with framing 2, whose Arf invariant is $1 \in \mathbf{Z}_2$. Therefore, again by [5, Theorem 4.2 (ii)], we see that the signature of the corresponding spin 4-manifold is $k - 10 \bmod 16$.

Summarising the above, we see that among the eight spin structures, four spin structures provide the μ -invariant $k \bmod 16$, three spin structures provide the μ -invariant $k - 2 \bmod 16$, and the last one provides the μ -invariant $k - 10 \bmod 16$. By the same argument as in the last part of the proof of Lemma 3.5, we see that the first four spin structures belong to one equivalence class in $\text{Spin}(X^3)/\text{Im } \rho \approx \mathbf{Z}_2$ and the remaining four belong to the other. Thus, by Remark 4.3, we have the desired conclusion. This completes the proof. $\square$

Remark 4.5. In the case of the 3-torus T^3 , the connected sum of a T^3 -pseudo-embedding $g: S^3 \looparrowright \mathbf{R}^5$ and an *arbitrary* embedding $T^3 \hookrightarrow \mathbf{R}^5$ is regularly homotopic to an embedding. Proposition 4.4 shows, however, that this is not true in general. Namely, for the above X^3 , the connected sum of an X^3 -pseudo-embedding $g: S^3 \looparrowright \mathbf{R}^5$ and an embedding $X^3 \hookrightarrow \mathbf{R}^5$ chosen from $24\mathbf{Z} \subset 12\mathbf{Z} \amalg 24\mathbf{Z} \approx \text{Emb}[X^3, \mathbf{R}^5]$ is not regularly homotopic to an embedding. This shows that we have to choose a correct embedding F for a given M^3 -pseudo-embedding in Definition 3.1 in general.

Remark 4.6. In Proposition 4.4, if we consider the 3-manifold Y^3 whose framed link representation is obtained by replacing the framings $(0, 0, 2)$ for the Borromean rings $L = \{K_1, K_2, K_3\}$ with the framings $(0, 2, 2)$, then we see that $\text{Imm}[Y^3, \mathbf{R}^5]_0 \approx \mathbf{Z} \amalg \mathbf{Z} \amalg \mathbf{Z} \amalg \mathbf{Z}$ and $\text{Emb}[Y^3, \mathbf{R}^5]$ forms the subset $12\mathbf{Z} \amalg 24\mathbf{Z} \amalg 24\mathbf{Z} \amalg 24\mathbf{Z} \subset \mathbf{Z} \amalg \mathbf{Z} \amalg \mathbf{Z} \amalg \mathbf{Z}$, by using a similar argument. These examples show that the distribution of $12\mathbf{Z}$ and $24\mathbf{Z}$ can be complicated in general. For other examples obtained by connected sum, see Section 5.

5. M^3 -PSEUDO-EMBEDDINGS OF GENERAL 3-MANIFOLDS

In this section, we consider a question which naturally arises from the definition of M^3 -pseudo-embeddings: *for a pair of closed oriented 3-manifolds N^3 and M^3 , does there exist an M^3 -pseudo-embedding of N^3 into $\mathbf{R}^5$?* In other words, when does there exist an immersion $g: N^3 \looparrowright \mathbf{R}^5$ such that (1) g is not regularly homotopic to any embedding, and (2) for some embedding $F: M^3 \hookrightarrow \mathbf{R}^5$, the connected sum $F \sharp g$ is regularly homotopic to an embedding $M^3 \sharp N^3 \hookrightarrow \mathbf{R}^5$?

Concerning this question, we have the following.

Proposition 5.1. *Let M^3 and N^3 be closed connected oriented 3-manifolds. Assume that*

$$\begin{aligned} \text{Imm}[M^3, \mathbf{R}^5]_0 &\approx Z(1) \amalg \cdots \amalg Z(i) \amalg \cdots \amalg Z(r), \\ \text{Imm}[N^3, \mathbf{R}^5]_0 &\approx Z(1) \amalg \cdots \amalg Z(j) \amalg \cdots \amalg Z(s), \end{aligned}$$

where $i \in \{1, 2, \dots, r\}$, $j \in \{1, 2, \dots, s\}$ and each $Z(i)$ or $Z(j)$ denotes a copy of $\mathbf{Z}$. Furthermore, assume that

$$\begin{aligned} \text{Emb}[M^3, \mathbf{R}^5] &\approx m_1\mathbf{Z} \amalg \cdots \amalg m_i\mathbf{Z} \amalg \cdots \amalg m_r\mathbf{Z}, \\ \text{Emb}[N^3, \mathbf{R}^5] &\approx n_1\mathbf{Z} \amalg \cdots \amalg n_j\mathbf{Z} \amalg \cdots \amalg n_s\mathbf{Z} \end{aligned}$$

with respect to the above decompositions of $\text{Imm}[M^3, \mathbf{R}^5]_0$ and $\text{Imm}[N^3, \mathbf{R}^5]_0$, respectively, where each m_i or n_j is equal either to 12 or to 24. Then, we have the following.

- (a) *The set $\text{Imm}[M^3 \sharp N^3, \mathbf{R}^5]_0$ corresponds bijectively to the disjoint union of rs copies of $\mathbf{Z}$. More precisely, we have*

$$\text{Imm}[M^3 \sharp N^3, \mathbf{R}^5]_0 \approx Z(1, 1) \amalg \cdots \amalg Z(i, j) \amalg \cdots \amalg Z(r, s),$$

where each $Z(i, j)$ denotes a copy of $\mathbf{Z}$ and each class in $Z(i, j)$ is represented by the connected sum of an immersion in $Z(i) \subset \text{Imm}[M^3, \mathbf{R}^5]_0$ and an immersion in $Z(j) \subset \text{Imm}[N^3, \mathbf{R}^5]_0$.

- (b) *In each $Z(i, j)$, the classes containing an embedding correspond to*

$$\begin{cases} 24\mathbf{Z}, & \text{if } (m_i, n_j) = (24, 24), \\ 12\mathbf{Z}, & \text{if } (m_i, n_j) = (12, 12), (12, 24), \text{ or } (24, 12). \end{cases}$$

- (c) *The set $\text{Imm}[N^3, \mathbf{R}^5]_0$ contains an M^3 -pseudo-embedding $N^3 \looparrowright \mathbf{R}^5$ if and only if the case $(m_i, n_j) = (12, 24)$ occurs. More precisely, if we assume $(m_i, n_j) = (12, 24)$, then an immersion $g: N^3 \looparrowright \mathbf{R}^5$ in $12 + 24\mathbf{Z} \subset Z(j) \subset \text{Imm}[N^3, \mathbf{R}^5]_0$ is not regularly homotopic to an embedding, but the connected sum of g and any embedding $M^3 \hookrightarrow \mathbf{R}^5$ in $12\mathbf{Z} \subset Z(i) \subset \text{Imm}[M^3, \mathbf{R}^5]_0$ is regularly homotopic to an embedding $M^3 \sharp N^3 \hookrightarrow \mathbf{R}^5$ in $12\mathbf{Z} \subset Z(i, j) \subset \text{Imm}[M^3 \sharp N^3, \mathbf{R}^5]_0$.*

Proof. (a) Since a spin structure of M^3 and a spin structure of N^3 determine a spin structure of $M^3 \sharp N^3$, and since $H^1(M^3 \sharp N^3) \cong H^1(M^3) \oplus H^1(N^3)$ (with any coefficients), we have the bijection

$$\sharp: (\text{Spin}(M^3)/\text{Im } \rho) \times (\text{Spin}(N^3)/\text{Im } \rho) \rightarrow \text{Spin}(M^3 \sharp N^3)/\text{Im } \rho.$$

Furthermore, since a normal framing of $F \in \text{Imm}[M^3, \mathbf{R}^5]_0$ and a normal framing of $G \in \text{Imm}[N^3, \mathbf{R}^5]_0$ determine a normal framing of $F \sharp G \in \text{Imm}[M^3 \sharp N^3, \mathbf{R}^5]_0$, and since the associated spin structures are coherent with respect to the connected sum, we see that

$$c(F \sharp G) = \sharp(c(F), c(G)).$$

Thus the result follows.

(b) This follows from the additivity of the μ -invariant with respect to the connected sum together with the above item (a) and Remark 4.3.

(c) This follows immediately from the above items (a) and (b). $\square$

Example 5.2. Let X^3 be the total space of the S^1 -bundle over $T^2 = S^1 \times S^1$ with Euler class 2 as in Proposition 4.4, and Y^3 the 3-manifold as in Remark 4.6. Then, for $M^3 := X^3 \sharp X^3$ or $X^3 \sharp Y^3$ or $Y^3 \sharp Y^3$, the numbers of $12\mathbf{Z}$ - and $24\mathbf{Z}$ -components of $\text{Emb}[M^3, \mathbf{R}^5] \subset \text{Imm}[M^3, \mathbf{R}^5]_0$ are as follows.

M^3	$12\mathbf{Z}$	$24\mathbf{Z}$
$X^3 \sharp X^3$	3	1
$X^3 \sharp Y^3$	5	3
$Y^3 \sharp Y^3$	7	9

In fact, we can show that for the connected sum of m copies of X^3 and n copies of Y^3 , the number of $24\mathbf{Z}$ -components is equal to 3^n and that of $12\mathbf{Z}$ is equal to $2^{m+2n} - 3^n$.

6. VIRTUAL HOMOTOPY CLASSES

In this section, we introduce a new equivalence relation on the set of all immersions of 3-manifolds into 5-space with trivial normal bundles, and study the structure of the quotient set. This relation measures, in a certain sense, the difference between the set of all regular homotopy classes of immersions and those containing embeddings.

Definition 6.1. Let $f: M^n \looparrowright \mathbf{R}^{n+k}$ and $g: N^n \looparrowright \mathbf{R}^{n+k}$ be immersions of closed *connected* oriented n -manifolds with trivial normal bundles. We say that f and g are *virtually homotopic* if there exist embeddings $f_1: M_1^n \hookrightarrow \mathbf{R}^{n+k}$ and $g_1: N_1^n \hookrightarrow \mathbf{R}^{n+k}$ of closed connected oriented n -manifolds such that $M^n \sharp M_1^n$ is orientation-preservingly diffeomorphic to $N^n \sharp N_1^n$ and $f \sharp f_1$ is regularly homotopic to $g \sharp g_1$. This clearly defines an equivalence relation on the set of all immersions of closed connected oriented n -manifolds into $\mathbf{R}^{n+k}$ with trivial normal bundles.

Let $\overline{\text{Imm}}_0(n, n+k)$ denote the set of virtual homotopy classes of immersions of closed connected oriented n -manifolds into $\mathbf{R}^{n+k}$ with trivial normal bundles.

Lemma 6.2. *The set $\overline{\text{Imm}}_0(3, 5)$ is an additive group under connected sum.*

Proof. For an arbitrary dimension pair $(n, n+k)$, it is easy to see that the connected sum (considering the orientation when $k = 1$) gives rise to a well-defined

binary operation on the set $\overline{\text{Imm}}_0(n, n+k)$. Then, clearly $\overline{\text{Imm}}_0(n, n+k)$ becomes an additive semi-group with respect to this operation; the virtual homotopy class containing embeddings is the identity element. When $(n, n+k) = (3, 5)$, for every immersion $f: M^3 \looparrowright \mathbf{R}^5$ of a closed connected oriented 3-manifold M^3 with trivial normal bundle, there exists an immersion $f_1: S^3 \looparrowright \mathbf{R}^5$ such that $f \sharp f_1$ is regularly homotopic to an embedding (see [9] or Subsection 2.2 of the present paper). Thus, every virtual homotopy class of $\overline{\text{Imm}}_0(3, 5)$ has an inverse. This completes the proof. $\square$

We do not know if $\overline{\text{Imm}}_0(n, n+k)$ is a group for every dimension pair $(n, n+k)$.

Proposition 6.3. *The group $\overline{\text{Imm}}_0(3, 5)$ is isomorphic to $\mathbf{Z}/12\mathbf{Z}$.*

Proof. Let $\varphi: \text{Imm}[S^3, \mathbf{R}^5] \rightarrow \overline{\text{Imm}}_0(3, 5)$ be the natural map, which is clearly a group homomorphism. Since every immersion of a closed connected oriented 3-manifold into $\mathbf{R}^5$ with trivial normal bundle is virtually homotopic to an immersion of S^3 (see Subsection 2.2), φ is an epimorphism. Furthermore, the kernel of φ consists of the immersions f of S^3 into $\mathbf{R}^5$ such that $f \sharp g$ is regularly homotopic to an embedding for some embedding g of a closed connected oriented 3-manifold. Thus, using the results in Subsection 2.2 and Theorem 4.2 together with the existence of T^3 -pseudo-embeddings of S^3 ([9]), we see that the kernel of φ consists of those immersions of S^3 into $\mathbf{R}^5$ whose Smale invariants are divisible by 12. Hence the result follows. $\square$

Remark 6.4. Let $f: M^3 \looparrowright \mathbf{R}^5$ be an immersion not regularly homotopic to an embedding. By the very definition of virtual homotopy, f is an N^3 -pseudo-embedding for some 3-manifold N^3 if and only if f is virtually homotopic to an embedding, i.e., if and only if f is an immersion that represents the identity element of the group $\overline{\text{Imm}}_0(3, 5)$.

REFERENCES

- [1] T. Ekholm and A. Szűcs, *Geometric formulas for Smale invariants of codimension two immersions*, to appear in *Topology*.
- [2] R. E. Gompf and A. I. Stipsicz, *4-manifolds and Kirby calculus*, Graduate Studies in Mathematics **20**, American Mathematical Society, Providence, RI, 1999. MR **2000h**:57038
- [3] J. Hughes, *Bordism and regular homotopy of low-dimensional immersions*, *Pacific J. Math.* **156** (1992), 155–184. MR **93j**:57019
- [4] J. Hughes and P. Melvin, *The Smale invariant of a knot*, *Comment. Math. Helv.* **60** (1985), 615–627. MR **87g**:57045
- [5] S. J. Kaplan, *Constructing framed 4-manifolds with given almost framed boundaries*, *Trans. Amer. Math. Soc.* **254** (1979), 237–263. MR **82h**:57015
- [6] M. A. Kervaire, *Sur le fibré normal à une sphère immergée dans un espace euclidien*, *Comment. Math. Helv.* **33** (1959), 121–131. MR **21**:3863
- [7] R. Kirby, *The topology of 4-manifolds*, *Lecture Notes in Math.* **1374**, Springer-Verlag, 1989. MR **90j**:57012
- [8] O. Saeki, *The theory of fibered 3-knots in S^5 and its applications*, *J. Math. Sci. Univ. Tokyo* **6** (1999), 691–756. MR **2001b**:57058
- [9] O. Saeki, A. Szűcs and M. Takase, *Regular homotopy classes of immersions of 3-manifolds into 5-space*, *Manuscripta Math.* **108** (2002), 13–32.

- [10] M. Takase, *Embeddings of $\mathbf{Z}_2$ -homology 3-spheres in $\mathbf{R}^5$ up to regular homotopy*, Pacific J. Math. **193** (2000), 249–256. MR **2001c**:57030
- [11] Wen-Tsün Wu, *On the immersion of C^∞ -3-manifolds in a Euclidean space*, Scientia Sinica **13** (1964), 335–336. MR **30**:2521

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, HIROSHIMA UNIVERSITY,
HIGASHI-HIROSHIMA 739-8526, JAPAN

E-mail address: `saeki@math.sci.hiroshima-u.ac.jp`

Current address: Faculty of Mathematics, Kyushu University, Hakozaki, Fukuoka 812-8581,
Japan

E-mail address: `saeki@math.kyushu-u.ac.jp`

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, 3-8-1 KOMABA,
MEGURO-KU, TOKYO, 153-8914, JAPAN

E-mail address: `takase@ms.u-tokyo.ac.jp`