CAUSAL COMPACTIFICATION OF COMPACTLY CAUSAL SPACES

FRANK BETTEN

ABSTRACT. We give a classification of causal compactifications of compactly causal spaces. Introduced by Ólafsson and Ørsted, for a compactly causal space G/H , these compactifications are given by G -orbits in the Bergman-Šilov boundary of G_1/K_1 , with $G \subset G_1$ and (G_1, K_1, θ) a Hermitian symmetric space of tube type. For the classical spaces an explicit construction is presented.

1. INTRODUCTION

A causal symmetric space is a symmetric space (G, H, τ) consisting of a (connected) Lie group G , an involutive automorphism τ of G , and a closed subgroup H with $G_0^\tau \subset H \subset G^\tau$ (here G_0^τ is the identity component of $G^\tau := \{g \in G \mid \tau(g) = g\}$) carrying a G -invariant causal structure. In general, a causal structure on a manifold M is the attachment of a cone $C_x \subset T_x M$ to every point $x \in M$. We always assume our cones to be closed, convex and regular, i.e. proper ($C_x \cap -C_x = \{0\}$) and generating ($C_x - C_x = T_x M$). With $\mathfrak{g}$ the Lie algebra of G , and the decomposition $\mathfrak{g} = \mathfrak{h} + \mathfrak{q}$ corresponding to τ , we can identify the tangent space at the identity coset $T_e H G/H$ of the homogenous space G/H with $\mathfrak{q}$. The G -invariant causal structures are now in one-to-one correspondence with H -invariant cones C in $\mathfrak{q}$. Restricting our attention to semisimple Lie algebras $\mathfrak{g}$ or to groups G and irreducible symmetric spaces, the irreducible symmetric Lie algebras $(\mathfrak{g}, \mathfrak{h}, \tau)$ corresponding to a causal symmetric space are classified ([Ól 90, HÓ 96, p. 89]) and can be divided into three classes. For that, let θ be a Cartan involution commuting with τ , and $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ the corresponding Cartan decomposition; then the subclass of compactly causal spaces is characterized by the existence of a cone with $C \cap \mathfrak{k} \neq \emptyset$ and $C \cap \mathfrak{p} = \{0\}$. c -duality, $\mathfrak{g}^c := \mathfrak{h} + i\mathfrak{q}$, defines a correspondence between compactly causal Lie algebras $(\mathfrak{g}, \mathfrak{h}, \tau)$ and noncompactly causal Lie algebras $(\mathfrak{g}^c, \mathfrak{h}, \tau)$. As a third subclass we can single out the intersection of these two classes, the spaces with a self-dual algebra, called causal symmetric spaces of Cayley type.

Definition 1.1 ([ÓØ 97]). Let M be a causal G -space. A *causal compactification* of M is a pair (N, Φ) that satisfies the following conditions:

- (1) N is a compact causal G -space.
- (2) The map $\Phi : M \rightarrow N$ is causal, i.e. if $m \mapsto C_m$ is the causal structure on M and $n \mapsto D_n$ that on N , then $(d\Phi)_m(C_m) \subset D_{\Phi(m)}$ for every $m \in M$.

Received by the editors October 7, 1996.

2000 *Mathematics Subject Classification.* Primary 53C35; Secondary 32M15.

- (3) Φ is injective and G -equivariant, i.e. $\Phi(g \cdot m) = g \cdot \Phi(m)$, for every $g \in G$ and every $m \in M$.
- (4) $\Phi(M)$ is open and dense in N .

Our aim in this article is to construct causal compactifications for compactly causal spaces. The compact space will always be the Bergman-Šilov boundary $\check{S}_1$ of a bounded symmetric domain of tube type. In fact, this boundary is a homogeneous space with an invariant causal structure ([Ka 89]). More precisely, if we choose (G_1, H_1, η) to be of Cayley type — this implies that G_1/K_1 is Hermitian symmetric of tube type ([HÓ 96, Remark 2.6.9]) — there is a parabolic subgroup P' , with $H_1 \subset P'$, such that $\check{S}_1 \simeq G_1/P'$ and the canonical projection $\Phi: G_1/H_1 \rightarrow G_1/P'$ is causal (and of course G_1 -equivariant). The basic strategy is now to find an involution σ of G_1 , commuting with θ and η , such that $(G, H, \tau) := ((G_1^\sigma)_0, H_1 \cap (G_1^\sigma)_0, \eta|(G_1^\sigma)_0)$ is a compactly causal subspace of (G_1, H_1, η) , i.e. its causal structure is defined by restricting the compactly causal structure of G_1/H_1 . If $G \cap P' = H$, we get by use of Φ an embedding $G/H \hookrightarrow G_1/P'$, and it remains to show that $\Phi(G/H)$ is (open) dense. This can be done using Matsuki's classification of $(G_1^\sigma, P_{\min})$ -double cosets, $P_{\min}$ a minimal parabolic subgroup of G_1 . Indeed, there are finitely many open double cosets $Gw_iP_{\min}$, the w_i being elements of an appropriate Weyl group, and it is possible to prove that GP' contains all of them, for $P_{\min} \subset P'$ suitably chosen.

Of course, in general the fixed point group G_1^σ is only reductive, and we therefore define

Definition 1.2. Let (G, H, τ) be a causal symmetric space with G reductive. Then (G, H, τ) is *weak compactly causal* if there exists an H -invariant, closed, convex and regular cone $C \subset \mathfrak{q}$ such that $C \cap \mathfrak{k} \neq \emptyset$ and $C \cap \mathfrak{p} = \{0\}$.

For semisimple G this implies the space is compactly causal by definition ([HÓ 96, p. 76]).

In Theorem 5.1 and Theorem 5.9 we give conditions which ensure that the sketched strategy works. Using these theorems, it is possible to construct causal compactifications for most of the compactly causal spaces — eventually with an additional central extension as explained above — listed in [Ól 90, HÓ 96, p. 89]. We summarize our results on causal compactifications of weak compactly causal spaces in the following three tables.

TABLE I. Theorem 5.1 construction

$(\mathfrak{g}, \mathfrak{h})$	$(\mathfrak{g}_1, \mathfrak{k}_1, \mathfrak{h}_1)$
$(\mathfrak{su}(p, q) \oplus \mathfrak{u}(p, q), \mathfrak{su}(p, q))$	$(\mathfrak{su}(n, n), \mathfrak{su}(n) \oplus \mathfrak{u}(n), \mathfrak{sl}(n, \mathbb{C}) \oplus \mathbb{R})$
$(\mathfrak{u}(2p, 2q), \mathfrak{sp}(p, q))$	$(\mathfrak{so}^*(4n), \mathfrak{u}(2n), \mathfrak{u}^*(2n))$
$(\mathfrak{u}(p, q), \mathfrak{so}(p, q))$	$(\mathfrak{sp}(n, \mathbb{R}), \mathfrak{u}(n), \mathfrak{gl}(n, \mathbb{R}))$
$(\mathfrak{so}(2, n-1) \oplus \mathfrak{so}(2), \mathfrak{so}(1, n-1))$	$(\mathfrak{so}(2, n+1), \mathfrak{so}(2) \oplus \mathfrak{so}(n+1), \mathfrak{so}(1, 1) \oplus \mathfrak{so}(1, n))$
$(\mathfrak{e}_{6(-14)} \oplus \mathfrak{so}(2), \mathfrak{f}_{4(-20)})$	$(\mathfrak{e}_{7(-25)}, \mathfrak{e}_6 \oplus \mathbb{R}, \mathfrak{e}_{6(-26)} \oplus \mathbb{R})$
$(\mathfrak{so}^*(2n) \oplus \mathfrak{so}^*(2n), \mathfrak{so}^*(2n))$	$(\mathfrak{so}^*(4n), \mathfrak{u}(2n), \mathfrak{u}^*(2n))$
$(\mathfrak{sp}(n, \mathbb{R}) \oplus \mathfrak{sp}(n, \mathbb{R}), \mathfrak{sp}(n, \mathbb{R}))$	$(\mathfrak{sp}(2n, \mathbb{R}), \mathfrak{u}(2n), \mathfrak{gl}(2n))$
$(\mathfrak{so}(2, n), \mathfrak{so}(1, n))$	$(\mathfrak{so}(2, n+1), \mathfrak{so}(2) \oplus \mathfrak{so}(n+1), \mathfrak{so}(1, 1) \oplus \mathfrak{so}(1, n))$

TABLE II. Theorem 5.9 construction

$(\mathfrak{g}, \mathfrak{h})$	$(\mathfrak{g}_1, \mathfrak{k}_1, \mathfrak{h}_1)$
$(\mathfrak{so}^*(2n), \mathfrak{so}(n, \mathbb{C}))$	$(\mathfrak{su}(n, n), \mathfrak{s}(\mathfrak{u}(n) \oplus \mathfrak{u}(n)), \mathfrak{so}(n, n))$
$(\mathfrak{sp}(2n, \mathbb{R}), \mathfrak{sp}(n, \mathbb{C}))$	$(\mathfrak{su}(2n, 2n), \mathfrak{s}(\mathfrak{u}(2n) \oplus \mathfrak{u}(2n)), \mathfrak{sp}(n, n))$
$(\mathfrak{so}(2, q) \oplus \mathfrak{so}(p+1), \mathfrak{so}(1, q) \oplus \mathfrak{o}(p))$	$(\mathfrak{so}(2, n+1), \mathfrak{so}(2) \oplus \mathfrak{so}(n+1), \mathfrak{so}(1, 1) \oplus \mathfrak{so}(1, n))$ ¹
$(\mathfrak{g}, \mathfrak{h})$ of Cayley type	$(\mathfrak{g} \oplus \mathfrak{g}, \mathfrak{k} \oplus \mathfrak{k}, \mathfrak{h} \oplus \mathfrak{h})$

Results by Makarevič ([Ma 73]) together with some simple dimension estimates show that the remaining compactly causal spaces have no causal compactification as symmetric orbits by imbedding them in a Bergman-Šilov boundary, cf. Proposition 6.1.

TABLE III. No compactification by Proposition 6.1

$(\mathfrak{g}, \mathfrak{h})$	parameters
$(\mathfrak{so}(2, n), \mathfrak{so}(p, 1) \oplus \mathfrak{so}(1, q))$	$\min(p, q) \geq 2, (p, q) \neq (2, 2), (3, 3)$
$(\mathfrak{e}_{6(-14)}, \mathfrak{sp}(2, 2))$	
$(\mathfrak{e}_{7(-25)}, \mathfrak{su}^*(8))$	
$(\mathfrak{so}(2, n) \oplus \mathfrak{so}(2, n), \mathfrak{so}(2, n))$	$n = 5 \text{ or } n \geq 7$
$(\mathfrak{e}_{6(-14)} \oplus \mathfrak{e}_{6(-14)}, \mathfrak{e}_{6(-14)})$	
$(\mathfrak{e}_{7(-25)} \oplus \mathfrak{e}_{7(-25)}, \mathfrak{e}_{7(-25)})$	

We remark that in the same manner as we use Matsuki's double coset decomposition to prove $\Phi(G/H)$ is dense, density of the image and the decomposition into orbits for other open symmetric orbits in R -spaces $G/H \rightarrow G_1/P'$, cf. [Ma 73] for a list, can be investigated.

This article is the first step towards a detailed study of the Hardy spaces on compactly causal spaces as introduced by Hilgert, Ólafsson and Ørsted ([HÓØ]). Roughly speaking, these are made up of holomorphic functions on certain open domains in $G_{\mathbb{C}}/H_{\mathbb{C}}$ determined by their boundary values on G/H . Similarly, we have the “classical” Hardy spaces on G_1/K_1 — or the biholomorphically equivalent tube domain — formed by certain holomorphic functions on G_1/K_1 determined by their boundary values on the Bergman-Šilov boundary G_1/P' . Causality of Φ assures that it can be prolonged, at least locally, to a map $\tilde{\Phi}$ from the mentioned open domain in $G_{\mathbb{C}}/H_{\mathbb{C}}$ into G_1/K_1 . Thus $\tilde{\Phi}^*$ maps functions on G_1/K_1 to functions on the domain of definition for the Hardy space, and this can be used to construct an imbedding of the “classical” Hardy space into the new one. This program is carried out only for Cayley spaces ([ÓØ 97]) and some of the group cases $(\mathfrak{g}, \mathfrak{h}) \simeq (\tilde{\mathfrak{g}} \oplus \tilde{\mathfrak{g}}, \tilde{\mathfrak{g}})$ ([KØ96, KØ97]) of our list above, and, as a main result, gives explicit formulas for the Cauchy-Szegő kernel. The general case is considered in [FÓ].

Results similar to ours were also obtained by Bertram ([Be]) using Jordan algebra methods. The method used by us has the advantage that it allows one to find the G -orbits in the compactification by looking not only at the open cosets but using the full double coset decomposition of Matsuki. For example, for spaces of Cayley type one recovers the orbit decomposition given by Kaneyuki ([Ka 87]).

I am grateful to H. Holdgrün and G. Ólafsson for their continuing interest, and for discussions concerning the content of the present paper. I would also like to thank the referee for several suggestions that improved the exposition considerably.

¹Here $p \geq 2$, whereas for the other cases p and q are only nonnegative with $p + q = n$.

2. BASIC STRUCTURE THEORY

Let $(\mathfrak{g}_1, \mathfrak{k}_1, \theta)$ be an (irreducible) orthogonal symmetric Lie algebra of noncompact Hermitian type. Denote the complexification of an algebra or group by a subscript $\mathbb{C}$, for example $\mathfrak{g}_{1\mathbb{C}} = \mathfrak{g}_1 \otimes \mathbb{C}$. Let $G_{1\mathbb{C}}$ be the simply connected group with Lie algebra $\mathfrak{g}_{1\mathbb{C}}$, and let (G_1, K_1, θ) be associated to $(\mathfrak{g}_1, \mathfrak{k}_1, \theta)$ with $G_1 \subset G_{1\mathbb{C}}$. For simplicity we will always denote the Cartan involution θ for G_1 or $\mathfrak{g}_1$ and its extension to the respective complexifications by the same letter. (The same notational convention will be used without further comment for all other morphisms.) Let $\mathfrak{g}_1 = \mathfrak{k}_1 + \mathfrak{p}_1$ be the Cartan decomposition for θ .

Let $\mathfrak{t} \subset \mathfrak{k}_1$ be a Cartan subalgebra and $\Delta(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{\mathbb{C}})$ the corresponding root system. Choose a positive system $\Delta^+(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{\mathbb{C}})$ such that the *noncompact roots* $\Delta^+(\mathfrak{p}_{1\mathbb{C}}, \mathfrak{t}_{\mathbb{C}})$ dominate the *compact roots* $\Delta^+(\mathfrak{k}_{1\mathbb{C}}, \mathfrak{t}_{\mathbb{C}})$. Recall that two roots α and β are *strongly orthogonal* if $\alpha \neq \pm\beta$ and $\alpha \pm \beta \notin \Delta(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{\mathbb{C}})$. Let $\{\gamma_1, \dots, \gamma_r\} \subset \Delta^+(\mathfrak{p}_{1\mathbb{C}}, \mathfrak{t}_{\mathbb{C}})$ be a maximal system of strongly orthogonal roots. Choose $E_{\pm j} \in \mathfrak{g}_{1\mathbb{C}, \pm\gamma_j}$, where $\mathfrak{g}_{1\mathbb{C}, \gamma}$ denotes the root space for γ , such that

$$X_j := E_j + E_{-j} \in \mathfrak{k}_1, \quad Y_j := iE_j - iE_{-j} \in \mathfrak{p}_1,$$

and

$$[E_j, E_{-j}] = H_j.$$

Here $H_j \in \mathfrak{t}$ is defined by

$$\langle H_j, H \rangle = 2 \frac{\gamma_j(H)}{\langle \gamma_j, \gamma_j \rangle}$$

for all $H \in \mathfrak{t}$ ([He 78, p. 387]). (As usual, define the pairing for roots α, β by

$$\langle \alpha, \beta \rangle = \langle H_\alpha, H_\beta \rangle = B(H_\alpha, H_\beta),$$

where B is the Killing form of $\mathfrak{g}_1$ and $H_\alpha \in \mathfrak{t}_{\mathbb{C}}$ the vector dual to α with respect to B .) It is known that $\mathfrak{a} = \sum \mathbb{R}X_j$ is maximal abelian in $\mathfrak{p}_1$.

Define the (full) *Cayley transform* by

$$c := \prod_{j=1}^r c_j, \quad \text{where } c_j = \exp \frac{\pi}{4} i Y_j.$$

By the strong orthogonality of the γ_j 's, an $\mathfrak{sl}(2)$ -calculation shows $\text{Ad}(c) i H_j = X_j$ and $\text{Ad}(c) \mathfrak{t}^- = \mathfrak{a}$, where $\mathfrak{t}^- := \sum \mathbb{R} i H_j \subset \mathfrak{t}$. By abuse of notation, let γ_j also denote the restriction of γ_j to $\mathfrak{t}^-$. The restricted root system is then given by the theorem of Moore.

Theorem 2.1 (Moore, [He 78, p. 528]). *For the restricted root system $\Delta(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{\mathbb{C}}^-)$ one of the following cases occurs:*

- (1) $\Delta(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{\mathbb{C}}^-) = \{\pm\gamma_j \mid 1 \leq j \leq r\} \cup \{\pm\frac{1}{2}\gamma_i \pm \frac{1}{2}\gamma_j \mid 1 \leq i < j \leq r\}$; or
- (2) $\Delta(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{\mathbb{C}}^-) = \{\pm\frac{1}{2}\gamma_j \mid 1 \leq j \leq r\} \cup \{\pm\gamma_j \mid 1 \leq j \leq r\} \cup \{\pm\frac{1}{2}\gamma_i \pm \frac{1}{2}\gamma_j \mid 1 \leq i < j \leq r\}$.

Using the Cayley transform, the structure of the root system $\Sigma(\mathfrak{a}) := \Delta(\mathfrak{g}_1, \mathfrak{a})$ can be immediately derived from this theorem.

For example, assume for the moment (G_1, K_1, θ) or $(\mathfrak{g}_1, \mathfrak{k}_1, \theta)$ to be Hermitian symmetric of tube type, i.e. G_1/K_1 is biholomorphically equivalent to a bounded symmetric domain of tube type. For this case the restricted root system is of

type $\mathfrak{c}_r$ ([KW 65, Prop. 3.12]) such that in the theorem above the root system is given by the first case. Therefore we have

$$\Sigma(\mathfrak{a}) = \{\pm\gamma_j^c \mid 1 \leq j \leq r\} \cup \{\pm\frac{1}{2}\gamma_i^c \pm \frac{1}{2}\gamma_j^c \mid 1 \leq i < j \leq r\},$$

where $\gamma_j^c := \gamma_j \circ \text{Ad}(c^{-1})$ is the Cayley transformed root. Furthermore, $\gamma_i^c(X_j) = 2\delta_{ij}$, such that the co-roots to the γ_j^c 's are up to normalization the X_j 's. Since the Weyl group $W(\mathfrak{a})$, which we consider to operate on the roots as well as on the X_j 's, is determined by the root system, we get also

Corollary 2.2. *For $(\mathfrak{g}_1, \mathfrak{k}_1, \theta)$ Hermitian symmetric of tube type we have $W(\mathfrak{a}) \simeq (\mathbb{Z}/2\mathbb{Z})^r \rtimes S_r$, where the permutations in S_r operate by permuting indices and $(\mathbb{Z}/2\mathbb{Z})^r \simeq \{(\epsilon_1, \dots, \epsilon_r) \mid \epsilon_j \in \{\pm 1\}\}$ by sign changes on $(X_1, \dots, X_r)$.*

With $\mathfrak{p}_1^+ := \sum_{\gamma \in \Delta^+(\mathfrak{p}_{1\mathbb{C}}, \mathfrak{t}_{\mathbb{C}})} \mathfrak{g}_{1\mathbb{C}, \gamma}$, $\mathfrak{p}_1^- := \sum_{\gamma \in \Delta^+(\mathfrak{p}_{1\mathbb{C}}, \mathfrak{t}_{\mathbb{C}})} \mathfrak{g}_{1\mathbb{C}, -\gamma}$, and

$$\mathfrak{g}_{1\mathbb{C}} = \mathfrak{p}_1^+ + \mathfrak{k}_{1\mathbb{C}} + \mathfrak{p}_1^-$$

the usual decomposition for a Hermitian symmetric algebra ([He 78, Ch. VIII]), $P_1^\pm$ and $K_{1\mathbb{C}}$ the corresponding analytic subgroups, let $x_0 := eK_{1\mathbb{C}}P_1^- \in G_{1\mathbb{C}}/K_{1\mathbb{C}}P_1^- =: M^*$ be the coset of the identity in the Borel realization of the Hermitian symmetric space G_1/K_1 . The *Bergman-Šilov boundary* $\check{S}_1$ of G_1/K_1 is the closed boundary orbit $G_1(cx_0)$ ([KW 65]). With an $\mathfrak{sl}(2)$ -calculation,

$$c_j = \exp(-E_j) \exp(\log \sqrt{2} H_j) \exp(E_{-j}).$$

Thus in the Harish-Chandra realization of G_1/H_1 in $\mathfrak{p}_1^+$, the Bergman-Šilov boundary is given as the G_1 -orbit through $-\sum E_j$. This boundary can also be described as the G_1 -orbit of $c^{-1}x_0$ or $\sum E_j$. Indeed, $c^{-1}x_0 = \theta(c)^{-1}x_0 = \exp(\sum E_j) \cdot x_0$. This is an element of the Bergman-Šilov boundary, since the exponential of the element $Z^0 := -\frac{i}{2} \sum H_j \in \mathfrak{z}(\mathfrak{k}_1)$, defining the complex structure on $\mathfrak{p}_1$, operates as complex rotations ($z \mapsto e^{i\lambda}z$) on $\mathfrak{p}_1^+$.

For $X^0 := \sum_{j=1}^r X_j$ the eigenvalues of $\text{ad } X^0$ are from $\{0, \pm 1, \pm 2\}$. We define $\mathfrak{p}'_{\mathbb{C}}$ to be the sum of the eigenspaces of $\text{ad } X^0$ in $\mathfrak{g}_{1\mathbb{C}}$ with nonpositive eigenvalues, and $\mathfrak{q}_1^+$ ($\mathfrak{q}_1^-$) to be the sum of the eigenspaces in $\mathfrak{g}_1$ with positive (respectively negative) eigenvalues. Let $P'_{\mathbb{C}}$ respectively Q_1^+ (Q_1^-) be the corresponding analytic subgroups in $G_{1\mathbb{C}}$ respectively G_1 .

Theorem 2.3 ([Wo 72, I.6]). *The stabilizer of the boundary point $cK_{1\mathbb{C}}P_1^-$ of G_1/K_1 is given by $P' := P'_{\mathbb{C}} \cap G_1$. Furthermore, if G_1 is simple, then P' is a maximal parabolic subgroup of G_1 .*

The theorem shows that $\check{S}_1 \simeq G_1/P'$, and we will call this homogeneous space the Bergman-Šilov boundary, too.

We also note for later use that for the parabolic subgroup P' we have the Langlands decomposition

$$P' \simeq M' \times A' \times Q_1^-,$$

where $A' := \exp \mathbb{R}X^0$ and $M'A' = Z_{G_1}(X^0)$.

3. CAUSAL STRUCTURES FOR CAYLEY SPACES AND THE BERGMAN-ŠILOV BOUNDARY

We will from now on always assume $(\mathfrak{g}_1, \mathfrak{k}_1, \theta)$ to be of tube type. Then $\text{ad } X^0$ has eigenvalues 0 and ± 2 . The inner automorphism

$$\eta := \text{Ad} \left(\exp \frac{\pi}{2} i X^0 \right)$$

is therefore involutive, and we get $H_1 := G_1^\eta = Z_{G_1}(X^0)$ and $\theta\eta = \eta\theta$. The triple (G_1, H_1, η) defines an (irreducible) symmetric space of Cayley type, and all Cayley spaces (up to covering) are obtained in this way ([HÓ 96, section 2.6]).

Let us recall the definition of the compactly causal structure on G_1/H_1 , cf. [HÓ 96, section 2.6]. Decomposing $Y^0 := \sum_{j=1}^r Y_j$ in (± 2) -eigenvectors $Y_\pm$ of $\text{ad } X^0$ — it is immediate to verify that $Y_\pm = \frac{1}{2} \sum_{j=1}^r (Y_j \mp i H_j)$ — we define $C_\pm := \overline{\text{conv Ad}(H_1)_0 \mathbb{R}^+ Y_\pm}$, which are H_1 -invariant cones. Then

$$C_k := C_+ - C_- \subset \mathfrak{q}_1^+ \oplus \mathfrak{q}_1^- =: \mathfrak{q}_1$$

is a regular H_1 -invariant cone. We have the canonical identification of $\mathfrak{q}_1$, which is the (-1) -eigenspace of η , with the tangent space $T_{eH_1}G_1/H_1$, and, using this identification, C_k defines the *compactly causal structure* of G_1/H_1 .

We also want to endow the Bergman-Šilov boundary with a G_1 -equivariant causal structure (cf. [Ka 89]). With $\check{S}_1 = G_1(c^{-1}x_0)$ we define the Cayley transform $\check{S}_1^c = cG_1(c^{-1}x_0) = (\text{Ad}(c)G_1)x_0$. Note that the left multiplication L_c on M^* defines a diffeomorphism of $\check{S}_1^c$ onto $\check{S}_1$ if G_1/K_1 is of tube type. Indeed, to be of tube type is equivalent to $\text{Ad}(c^2)\mathfrak{k}_1 = \mathfrak{k}_1$ ([KW 65, Prop. 4.4]), which implies $\text{Ad}(c^2)\mathfrak{g}_1 = \mathfrak{g}_1$, and then we have $L_c\check{S}_1^c = (\text{Ad}(c^2)G_1)(cx_0) = G_1(cx_0) = \check{S}_1$.

Let us identify the tangent space $T_{x_0}M^*$ with $\mathfrak{p}_1^+$ by use of the differential of the map $\xi : \mathfrak{p}_1^+ \rightarrow M^*$, $X \mapsto \exp X \cdot x_0$. Thereby the subspace $T_{x_0}\check{S}_1^c$ corresponds to $\mathfrak{s}_1^+ := \mathfrak{p}_1^+ \cap \text{Ad}(c)\mathfrak{g}_1$, and this space can be given the structure of a Euclidean or formally real Jordan algebra ([KW 65, section 6]). The cone defining the causal structure of $\check{S}_1^c$ is then given by the identity component of the set of all squares in $\mathfrak{s}_1^+$ ([Ka 89]), or equivalently the closure of the K_1^c -orbit of $E := -i\xi^{-1}(cx_0) = i \sum E_j$, with $K_1^c := K_{1\mathbb{C}} \cap \text{Ad}(c)G_1$. We define the causal structure of $\check{S}_1$ now by claiming that L_c is a causal diffeomorphism.

Let us describe this causal structure more explicitly. First of all, by use of the differential of $\xi^c : \text{Ad}(c)\mathfrak{p}_1^+ \rightarrow M^*$, $\text{Ad}(c)X \mapsto \exp(\text{Ad}(c)X)cx_0$, we identify the spaces $\text{Ad}(c)\mathfrak{p}_1^+$ and $T_{cx_0}M^*$. For spaces of tube type $\text{Ad}(c)(iZ^0) = \frac{1}{2}X^0$, and correspondingly $\text{Ad}(c)\mathfrak{p}_1^+$ is the $(+2)$ -eigenspace $\mathfrak{q}_{1\mathbb{C}}^+$ of $\text{ad } X^0$ in $\mathfrak{g}_{1\mathbb{C}}$. We get

$$\text{Ad}(c)\mathfrak{s}_1^+ = \text{Ad}(c)\mathfrak{p}_1^+ \cap \text{Ad}(c^2)\mathfrak{g}_1 = \text{Ad}(c)\mathfrak{p}_1^+ \cap \mathfrak{g}_1 = \mathfrak{q}_1^+$$

for the tangent space $T_{cx_0}\check{S}_1$ (with our identifications). Denote by $H_{1\mathbb{C}}$ the analytic subgroup corresponding to $\mathfrak{h}_{1\mathbb{C}} := \mathfrak{z}_{\mathfrak{g}_{1\mathbb{C}}}(X^0)$; then

$$H_1 = H_{1\mathbb{C}} \cap G_1 = \text{Ad}(c)K_{1\mathbb{C}} \cap G_1 = \text{Ad}(c)K_1^c.$$

Together with $\text{Ad}(c)E = Y_+$, our above definition now shows that the defining cone in $\mathfrak{q}_1^+$ is the closure of the H_1 -orbit of Y_+ , i.e. $C_+ = \overline{\text{Ad}(H_1)Y_+}$.

Remark. Alternatively, one can proceed as follows, cf. [Ka 89]. Since the cone $C_+ = \overline{\text{conv Ad}(H_1)_0 \mathbb{R}^+ Y_+}$ is regular and H_1 -invariant in $\mathfrak{q}_1^+$, it is easy to see that this cone is invariant under the linear isotropy representation of P' in $T_{eP'}G_1/P'$.

It therefore defines a G_1 -equivariant causal structure on G_1/P' , unique up to sign change.

Since the tangent map of the projection

$$\Phi : G_1/H_1 \rightarrow G_1/P'$$

at the identity coset is given, using the identifications introduced above, by the projection of $\mathfrak{q}_1 = \mathfrak{q}_1^+ \oplus \mathfrak{q}_1^-$ onto $\mathfrak{q}_1^+$, and the fact both causal structures are G_1 -equivariant, this projection of the Cayley space G_1/H_1 on the Bergman-Šilov boundary is causal.

Remark. The projection is also causal if we consider the Cayley space with its *noncompactly causal structure* defined by $C_p := C_+ + C_-$.

Later on we will consider subspaces $G/H \subset G_1/H_1$. The question arises: When is G/H a causal subspace, i.e., when does G/H have a causal structure such that $G/H \hookrightarrow G_1/H_1$ is causal? If $G = G_1^\sigma$ is the fixed point group of an involution σ , the following lemma gives a simple criterion.

Lemma 3.1. *Let σ be an involution commuting with η and θ . Let $C \subset \mathfrak{q}_1$ be a convex generating σ -invariant cone. Let $\mathfrak{q}_{1\sigma}$ be the (-1) -eigenspace of σ and $\mathfrak{q}_1 = \mathfrak{q}_1^\sigma \oplus \mathfrak{q}_1 \cap \mathfrak{q}_{1\sigma}$ the decomposition in (± 1) -eigenspaces of σ . Then $C \cap \mathfrak{q}_1^\sigma = \text{pr}_{\mathfrak{q}_1^\sigma} C$, and this cone is generating in $\mathfrak{q}_1^\sigma$.*

Proof. For $v \in \text{pr}_{\mathfrak{q}_1^\sigma} C$ there exists $w \in \mathfrak{q}_1 \cap \mathfrak{q}_{1\sigma}$ such that $v + w \in C$. By σ -invariance $v - w \in C$, and then $v = \frac{1}{2}(v + w) + \frac{1}{2}(v - w) \in C$, which proves that $\text{pr}_{\mathfrak{q}_1^\sigma} C \subset C \cap \mathfrak{q}_1^\sigma$. The other inclusion being trivial, we finally have

$$\mathfrak{q}_1^\sigma = \text{pr}_{\mathfrak{q}_1^\sigma} (C - C) = \text{pr}_{\mathfrak{q}_1^\sigma} C - \text{pr}_{\mathfrak{q}_1^\sigma} C.$$

□

4. DOUBLE COSET DECOMPOSITIONS

In this section we recall some results of Matsuki on the (G, P') -cosets in G_1 , where G is an open subgroup of the fixed point group of an involution and P' a parabolic subgroup.

If we look at minimal parabolic subgroups, these can be parametrized in the usual manner by assigning a maximal abelian subspace $\mathfrak{a}$ of $\mathfrak{p}_1$ and a system of positive roots $\Sigma^+ \subset \Sigma(\mathfrak{a})$, with $\Sigma(\mathfrak{a})$ the root system of $\mathfrak{g}_1$ with respect to $\mathfrak{a}$. We indicate this by writing $P(\mathfrak{a}, \Sigma^+)$, and $\mathfrak{p}(\mathfrak{a}, \Sigma^+)$ for the algebra. For a subset $\mathfrak{r}$ of $\mathfrak{a}$ we define a subset of (positive) roots by $\Sigma^{(+)}(\mathfrak{r}) := \{\gamma \in \Sigma^{(+)} \mid H_\gamma \in \mathfrak{r}\}$, where $H_\gamma \in \mathfrak{a}$ is the co-root to γ defined by $B(H_\gamma, H) = \gamma(H)$ for all $H \in \mathfrak{a}$. Since all minimal parabolic subgroups are conjugate and self-normalizing, we can identify the factor space G_1/P with the set of all minimal parabolic subalgebras of $\mathfrak{g}_1$.

Let (G_1, G, σ) be a symmetric triple, i.e. $(G_1^\sigma)_0 \subset G \subset G_1^\sigma$, and assume σ to commute with θ . Let $\mathfrak{g}_1 = \mathfrak{g} + \mathfrak{q}_{1\sigma}$ be the decomposition into eigenspaces of σ , and $\mathfrak{a}$ a σ -stable maximal abelian subspace of $\mathfrak{p}_1$. We define two subgroups of the Weyl group $W(\mathfrak{a}) = N_{K_1}(\mathfrak{a})/Z_{K_1}(\mathfrak{a})$ by

$$W_\sigma(\mathfrak{a}) := \{w \in W(\mathfrak{a}) \mid w(\mathfrak{a} \cap \mathfrak{g}) = \mathfrak{a} \cap \mathfrak{g}\}$$

and

$$W(\mathfrak{a}, K_1 \cap G) := N_{K_1 \cap G}(\mathfrak{a})/Z_{K_1 \cap G}(\mathfrak{a}) \subset W_\sigma(\mathfrak{a}).$$

Proposition 4.1 ([Ma 79, Prop. 1]). *A minimal parabolic subalgebra $\mathfrak{p}(\mathfrak{a}, \Sigma^+)$, which is identified with a point of G_1/P , is contained in an open G -orbit if and only if the following two conditions are satisfied:*

- (i) $\mathfrak{a} \cap \mathfrak{q}_{1\sigma}$ is maximal abelian in $\mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma}$, and
- (ii) Σ^+ is $\sigma\theta$ -compatible, i.e. $\gamma \in \Sigma^+ \setminus \Sigma^+(\mathfrak{a} \cap \mathfrak{g}) \Rightarrow \sigma\theta(\gamma) \in \Sigma^+ \setminus \Sigma^+(\mathfrak{a} \cap \mathfrak{g})$.

The number of open orbits is $|W_\sigma(\mathfrak{a})| / |W(\mathfrak{a}, K_1 \cap G)|$.

Remark. Condition (i) determines $\mathfrak{a}$ up to $\text{Ad}(K_1 \cap G)_0$ -conjugacy.

Assuming P' to be an arbitrary parabolic subgroup, we want to determine the double cosets $GgP(\mathfrak{a}, \Sigma^+)$ within GP' . Let therefore $\mathfrak{p}' = \mathfrak{m}' + \mathfrak{a}' + \mathfrak{n}'$ be the Langlands decomposition of $\mathfrak{p}'$ such that $\mathfrak{a}' \subset \mathfrak{a}$. Define

$$\mathfrak{a}'_+ := \{X \in \mathfrak{a}' \mid \gamma(X) > 0 \text{ for all } \gamma \text{ with } \mathfrak{g}_{1,\gamma} \subset \mathfrak{n}'\},$$

where $\mathfrak{g}_{1,\gamma}$ is the root space for $\gamma \in \Sigma(\mathfrak{a})$.

Theorem 4.2 ([Ma 82]). *Every minimal parabolic subalgebra of $\mathfrak{g}_1$ contained in $\mathfrak{p}'$ is $G \cap P'$ -conjugate to a minimal parabolic subalgebra $\mathfrak{p}(\mathfrak{a}_1, \Sigma(\mathfrak{a}_1)^+)$, where $\mathfrak{a}_1$ is a σ -stable maximal abelian subspace of $\mathfrak{p}_1$ such that $\mathfrak{a}_1 \supset \mathfrak{a}'$ and $\Sigma(\mathfrak{a}_1)^+$ satisfies $\langle \Sigma(\mathfrak{a}_1)^+, \mathfrak{a}'_+ \rangle \subset \mathbb{R}^+$.*

5. CAUSAL COMPACTIFICATIONS

We are now ready to state and prove our main results. If not otherwise stated, our notation will be as introduced in the previous sections.

Theorem 5.1. *Let (G_1, H_1, η) be an irreducible causal symmetric space of Cayley type with $G_1 \subset G_{1\mathbb{C}}$, $G_{1\mathbb{C}}$ the simply connected Lie group corresponding to $\mathfrak{g}_{1\mathbb{C}}$. Let σ be an involution commuting with η and θ . Assume one of the following cases holds:*

- (i) $\sum \mathbb{R}X_j$ and $\sum \mathbb{R}Y_j$ are maximal abelian in $\mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma}$, where $\mathfrak{q}_{1\sigma}$ is the (-1) -eigenspace of the involution σ .
- (ii) We have $r = 2r'$, and, with $X'_{j'} := X_{j'} + X_{j'+r'}$, $Y'_{j'} := Y_{j'} + Y_{j'+r'}$, for $j' = 1, \dots, r'$, the spaces $\sum \mathbb{R}X'_{j'}$ and $\sum \mathbb{R}Y'_{j'}$ are maximal abelian in $\mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma}$.

Then $(G, H, \tau) := ((G_1^\sigma)_0, H_1 \cap (G_1^\sigma)_0, \eta|(G_1^\sigma)_0)$ is a weak compactly causal subspace of (G_1, H_1, η) , and $\Phi : G/H \rightarrow G_1/P'$ is a causal compactification.

Proof. For the eigenspaces of $\text{ad } X^0$ we have $\sigma(\mathfrak{q}_1^-) = \mathfrak{q}_1^+$, since $X^0 \in \sum \mathbb{R}X_j$ and $X^0 \in \sum \mathbb{R}X'_{j'}$. Using the Langlands decomposition of P' , this implies on the group level $Q_1^- \cap G = \{e\}$ and $G \cap P' = G \cap H_1 = H$, i.e. Φ is injective. The subspace $T_{eH}G/H \subset T_{eH_1}G_1/H_1$ corresponds with our identifications to $\{X + \sigma X \mid X \in \mathfrak{q}_1^+\}$, and the restriction of the tangent map of Φ is given by $T\Phi(X + \sigma X) = X$. Therefore, by G -equivariance, Φ is a submersion onto an open image.

Let us show the density of the image. From (i) $\mathfrak{a} := \sum \mathbb{R}X_j$ is a maximal abelian σ -invariant subspace of $\mathfrak{p}_1$, and the first condition of Proposition 4.1 is fulfilled. Assume $\Sigma(\mathfrak{a})^+$ to be $\sigma\theta$ -compatible; then all other positive systems are conjugates $w(\Sigma(\mathfrak{a})^+)$, with $w \in W(\mathfrak{a})$. The system $w(\Sigma(\mathfrak{a})^+)$ is again $\sigma\theta$ -compatible if and only if $w \in W_\sigma(\mathfrak{a})$. Indeed, by Theorem 2.1 and the following discussion, we have $\Sigma(\mathfrak{a}) = \{\pm \frac{1}{2}(\gamma_s^c + \gamma_t^c), \pm \frac{1}{2}(\gamma_u^c - \gamma_v^c) \mid 1 \leq s, t, u, v \leq r, u \neq v\}$, and X_j is up to normalization the co-root to γ_j^c . Therefore, identifying $\mathfrak{a}$ and its dual

$\mathfrak{a}^*$, $\mathfrak{a} \cap \mathfrak{g}$ is spanned by $\Sigma(\mathfrak{a} \cap \mathfrak{g})^+$. By Proposition 4.1 the $(G, P(\mathfrak{a}, \Sigma^+))$ -double cosets corresponding to open orbits in $G_1/P(\mathfrak{a}, \Sigma^+)$ are then given by $Gw_v P(\mathfrak{a}, \Sigma^+)$, where Σ^+ is an arbitrary $\sigma\theta$ -compatible positive system and w_v a representative for $v \in W(\mathfrak{a}, K_1 \cap G) \backslash W_\sigma(\mathfrak{a})$.

It is now enough to show that all these open double cosets lie in GP' , and we use Theorem 4.2 to do this. With $\mathfrak{a}' = \mathbb{R}X^0$ we have $\mathfrak{a}'_+ = -\mathbb{R}^+X^0$, as $\mathfrak{q}_1^-$ is the (-2) -eigenspace of $\text{ad } X^0$. Let $\Sigma(\mathfrak{a})^+$ fulfill the condition of Theorem 4.2, i.e. $\langle \Sigma(\mathfrak{a})^+, X^0 \rangle \leq 0$; then $w(\Sigma(\mathfrak{a})^+)$ obeys the same condition if and only if

$$\langle w(\Sigma(\mathfrak{a})^+), \mathfrak{a}'_+ \rangle = -\mathbb{R}^+ \langle \Sigma(\mathfrak{a})^+, w^{-1}(X^0) \rangle \geq 0$$

or equivalently $w(X^0) = X^0$. This shows that the open orbits lying in GP' are parametrized by $W(\mathfrak{a})_{\mathfrak{m}'} \cap W(\mathfrak{a}, K_1 \cap G) \backslash W(\mathfrak{a})_{\mathfrak{m}'}$, where

$$W(\mathfrak{a})_{\mathfrak{m}'} := N_{K_1 \cap M'}(\mathfrak{a}) / Z_{K_1 \cap M'}(\mathfrak{a})$$

is the subgroup of the Weyl group leaving X^0 fixed. With the commutative diagram

$$\begin{array}{ccccc} W(\mathfrak{a})_{\mathfrak{m}'} \cap W_\sigma(\mathfrak{a}) & \rightarrow & W_\sigma(\mathfrak{a}) & \rightarrow & W(\mathfrak{a}, K_1 \cap G) \backslash W_\sigma(\mathfrak{a}) \\ & & \downarrow & & \nearrow \varphi \\ W(\mathfrak{a})_{\mathfrak{m}'} \cap W(\mathfrak{a}, K_1 \cap G) \backslash W(\mathfrak{a})_{\mathfrak{m}'} & & & & \end{array}$$

it remains to show the surjectivity of φ . But this is a consequence of Corollary 2.2. Indeed, since the permutations preserve $X^0 = \sum_{j=1}^r X_j$, for our case (i) it is enough to show that the subgroup isomorphic to $(\mathbb{Z}/2\mathbb{Z})^r \times \{\text{id}\}$ lies within $W(\mathfrak{a}, K_1 \cap G)$. But with $X_j, Y_j \in \mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma}$ and therefore $[X_j, Y_j] = -2iH_j \in \mathfrak{k}_1 \cap \mathfrak{g}$ this is a direct consequence of

$$(1) \quad \text{Ad}(\exp \frac{\pi}{2} iH_j)(E_{\pm k}) = \begin{cases} E_{\pm k} & \text{for } k \neq j, \\ -E_{\pm k} & \text{for } k = j. \end{cases}$$

For case (ii) we similarly have to prove that the “sign changes” in $W_\sigma(\mathfrak{a})$ are elements of $W(\mathfrak{a}, K_1 \cap G)$. These sign changes are generated by the reflections which map $X'_{j'} = X_{j'} + X_{j'+r'}$ to its negative and leave all vectors orthogonal to this one fixed. Here $X'_{j'}, Y'_{j'} \in \mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma}$. Thus we have $[X'_{j'}, Y'_{j'}] \in \mathfrak{k}_1 \cap \mathfrak{g}_1$, and the needed result follows again from (1).

Knowing that Φ is a G -equivariant diffeomorphism onto an open and dense subset, it remains to prove the claims concerning causality. We consider first the case (i). With $\sigma(Y_j) = -Y_j$ or $\sigma(iH_j) = iH_j$ we get $\sigma(Y_+) = -Y_-$ by the definition of $Y_\pm$, cf. section 3. This implies $\sigma(C_\pm) = -C_\mp$ and the σ -invariance of C_k . The cone C_k being regular H_1 -invariant, we see by Lemma 3.1 that $C_k \cap \mathfrak{g} \cap \mathfrak{q}_1$ is an H -invariant regular cone in $\mathfrak{g} \cap \mathfrak{q}_1$. This cone has nontrivial intersection with $\mathfrak{k}_1 \cap \mathfrak{g}$, as $Y_+ - Y_- = 2Z^0 \in \mathfrak{k}_1 \cap \mathfrak{g}$, i.e. G/H is weak compactly causal. Case (ii) is treated in the same way, with the obvious modifications. Finally, since we proved $\Phi : G_1/H_1 \rightarrow G_1/P'$ to be causal, it is clear that the restriction of this map to the causal subspace G/H is causal, too. $\square$

Remark. 1. Note that $C_p = C_+ + C_-$ will be mapped by σ to $-C_p$, and therefore G/H is not a noncompactly causal subspace of the Cayley space G_1/H_1 . In turn this shows that $G_a/H_1 \cap G_a$, with $G_a := (G_1^{\sigma\theta})_0$ the associated group to G , is a Cayley subspace.

2. The maximality condition for case (ii) is equivalent to the fact that G_a has real rank r' .

In general, for $G_{1\mathbb{C}}$ not simply connected, some of the inclusions $(G_1)_0 \subset \{g \in G_{1\mathbb{C}} \mid \bar{g} = g\} =: G_1$, where $\bar{g}$ denotes conjugation with respect to the real form $\mathfrak{g}_1$, and $(G_1^\eta)_0 \subset Z_{G_1}(X^0) \subset G_1^\eta = H_1$ may be proper. For the first case, since we are finally interested in quotients like G_1/H_1 and G_1/P' , it does not matter if we work with the group of all “real” matrices G_1 or its identity component. For the second case we have to replace $H_{(1)}$ by $Z_{G_{(1)}}(X^0)$ everywhere in our statements, and all arguments will go through without any change. For the classical groups both complications will only occur for $G_1 = SO(2, n)$.

Example 5.2 (Causal compactification of $SO(2, n)/SO(1, n)$). We start with

$$G_1 = \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in SL(n+3, \mathbb{R}) \mid \begin{array}{l} {}^tAA - {}^tCC = I, \quad {}^tAB = {}^tCD \\ {}^tDD - {}^tBB = I \end{array} \right\},$$

i.e. $G_1 = SO(2, n+1)$, and $\theta(g) = {}^tg^{-1}$ the usual Cartan involution. We assume $n \geq 2$, since $\mathfrak{so}(2, 1) \simeq \mathfrak{su}(1, 1)$ and $\mathfrak{so}(2, 2) \simeq \mathfrak{su}(1, 1) \times \mathfrak{su}(1, 1)$, which thus can be treated differently. The real rank is then two, and $G_{1\mathbb{C}} = SO(n+3, \mathbb{C})$ is not simply connected. We choose

$$E_1 = \frac{1}{2} \begin{pmatrix} 0 & \dots & -i & -1 \\ \vdots & & -1 & i \\ -i & -1 & & \vdots \\ -1 & i & \dots & 0 \end{pmatrix}, \quad E_{-1} = \frac{1}{2} \begin{pmatrix} 0 & \dots & i & -1 \\ \vdots & & -1 & -i \\ i & -1 & & \vdots \\ -1 & -i & \dots & 0 \end{pmatrix}$$

together with

$$E_2 = \frac{1}{2} \begin{pmatrix} 0 & \dots & i & -1 \\ \vdots & & 1 & i \\ i & 1 & & \vdots \\ -1 & i & \dots & 0 \end{pmatrix}, \quad E_{-2} = \frac{1}{2} \begin{pmatrix} 0 & \dots & -i & -1 \\ \vdots & & 1 & -i \\ -i & 1 & & \vdots \\ -1 & -i & \dots & 0 \end{pmatrix}.$$

The involution $\eta = \text{Ad}(\exp i\frac{\pi}{2}X^0)$ has the fixed point group

$$H_1 = \left\{ \begin{pmatrix} a_1 & 0 & 0 & b_2 \\ 0 & a_4 & B_3 & 0 \\ 0 & C_2 & D_1 & 0 \\ c_3 & 0 & 0 & d_4 \end{pmatrix} \in SL(n+3, \mathbb{R}) \mid \dots \right\} \simeq S(O(1, 1) \times O(1, n)),$$

and (G_1, H_1, η) is of Cayley type. It is straightforward to check that $Z_{G_1}(X^0)$ consists of those elements in H_1 for which $a_1d_4 - b_2c_3 = 1$, whereas for an arbitrary element $a_1d_4 - b_2c_3 \in \{\pm 1\}$, i.e. $H_1/Z_{G_1}(X^0) \simeq \mathbb{Z}/2\mathbb{Z}$.

We choose σ to be the inner automorphism

$$\sigma = \text{Ad} \begin{pmatrix} 1_2 & & \\ & -1_n & \\ & & 1 \end{pmatrix};$$

then its fixed point group is

$$G_1^\sigma = \left\{ \begin{pmatrix} A & B' & 0 \\ C' & D'_1 & 0 \\ 0 & 0 & d_4 \end{pmatrix} \in SL(n+3, \mathbb{R}) \mid \dots \right\} \simeq S(O(2, n) \times O(1)),$$

and one verifies condition (ii) of Theorem 5.1 for this involution. We have

$$G = \left\{ \begin{pmatrix} A & B' & 0 \\ C' & D_1' & 0 \\ 0 & 0 & 1 \end{pmatrix} \middle| \begin{pmatrix} A & B' \\ C' & D_1' \end{pmatrix} \in SO_0(2, n) \right\},$$

$$H = \left\{ \begin{pmatrix} a_1 & 0 & 0 & 0 \\ 0 & a_4 & B_3 & 0 \\ 0 & C_2 & D_1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \in G \middle| \begin{array}{l} a_1 a_4 \geq 1 \\ \det D_1 \geq 1 \end{array} \right\},$$

and $Z_G(X^0) = H \cap Z_{G_1}(X^0)$ is the subgroup of matrices with $a_1 = 1$, i.e. $Z_G(X^0) \simeq SO_0(1, n)$ in the obvious manner. Therefore $(SO_0(2, n), SO_0(1, n), \tau)$ is a compactly causal subspace of $(G_1, Z_{G_1}(X^0), \eta)$ with a causal compactification $\Phi : SO_0(2, n)/SO_0(1, n) \rightarrow SO(2, n+1)/P'$ as in Theorem 5.1. Another treatment of this example can be found in [Mo 97].

For every Cayley space (G_1, H_1, η) as in the theorem, there is a whole family of involutions σ for which condition (i) can be easily verified.

Corollary 5.3. *Let (G_1, H_1, η) be an irreducible causal symmetric space of Cayley type. Then, for $\epsilon_j \in \{\pm 1\}$, the inner involution $\sigma = \text{Ad}(\exp \frac{\pi}{2} i \sum \epsilon_j H_j)$ commutes with η and θ , (G, H, τ) is a weak compactly causal subspace of (G_1, H_1, η) , and $\Phi : G/H \rightarrow G_1/P'$ is a causal compactification.*

Proof. An $\mathfrak{sl}(2)$ -calculation shows $\sigma^2 = \text{Ad}(\exp(\pi i \sum \epsilon_j H_j)) = \text{id}$. Since $\theta(iH_j) = -iH_j$, and by another $\mathfrak{sl}(2)$ -calculation $\eta(H_j) = -H_j$, the three involutions η , θ and σ commute pairwise. Finally, from equation (1) — where we can replace H_j also by $\epsilon_j H_j$ — it is immediate that $X_j, Y_j \in \mathfrak{q}_{1\sigma}$. $\square$

Lemma 5.4. *Let $(\mathfrak{g}, \mathfrak{k}, \theta)$ be irreducible Hermitian of noncompact type, ρ an involution commuting with θ . Then $\dim \mathfrak{z}(\mathfrak{g}^\rho) \leq 1$.*

Proof. The Riemannian dual $\mathfrak{g}^r$ of $(\mathfrak{g}, \mathfrak{g}^\rho, \rho)$ is simple with $\mathfrak{g}_{\mathbb{C}}^r = \mathfrak{g}_{\mathbb{C}}$ simple, too. Therefore $\mathfrak{g}^r$ is irreducible orthogonal symmetric of type III ([He 78, p. 379]) and its maximal compact subalgebra $\mathfrak{g}^\rho \cap \mathfrak{k} \oplus i(\mathfrak{g}^\rho \cap \mathfrak{p})$ has a center $\mathfrak{c}$ of dimension at most one. θ is an involution of this subalgebra, which implies $\mathfrak{c} = \mathfrak{g}^\rho \cap \mathfrak{k} \cap \mathfrak{c} \oplus i(\mathfrak{g}^\rho \cap \mathfrak{p}) \cap \mathfrak{c}$ and $\mathfrak{z}(\mathfrak{g}^\rho) = \mathfrak{g}^\rho \cap \mathfrak{k} \cap \mathfrak{c} + i(i(\mathfrak{g}^\rho \cap \mathfrak{p}) \cap \mathfrak{c})$. $\square$

As σ in the corollary is given in a very explicit form, it is possible to derive some additional information on the structure of $(\mathfrak{g}, \mathfrak{h}, \tau)$ for this case.

Proposition 5.5. *Let $(\mathfrak{g}_1, \mathfrak{h}_1, \tau)$ be as in Corollary 5.3. Then:*

- (1) $\mathfrak{z}(\mathfrak{g}) = \mathbb{R}(\sum \epsilon_j i H_j) \subset \mathfrak{q}_1 \cap \mathfrak{k}_1$ and $\mathfrak{z}(\mathfrak{h}_1) = \mathbb{R}X^0 \subset \mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma}$.
- (2) $(\mathfrak{g}, \theta, \tau) \simeq (\mathfrak{k}_1^c, \sigma, \eta)$, where $\mathfrak{k}_1^c := \mathfrak{k}_1 \cap \mathfrak{g} \oplus i(\mathfrak{k}_1 \cap \mathfrak{q}_{1\sigma})$ is the c -dual of $\mathfrak{k}_1 = \mathfrak{k}_1 \cap \mathfrak{g} \oplus \mathfrak{k}_1 \cap \mathfrak{q}_{1\sigma}$.
- (3) $([\mathfrak{g}, \mathfrak{g}], \mathfrak{h}, \tau)$ is an irreducible symmetric space.

Proof. We have $\eta(H_j) = -H_j$ by an $\mathfrak{sl}(2)$ -calculation, i.e. $H_j \in \mathfrak{q}_1$. The first statement then follows by applying Lemma 5.4 to the involutions σ and η .

For the second claim we first prove that $(\mathfrak{g}, \theta) \simeq (\mathfrak{k}_1^c, \sigma)$. Defining

$$A := \frac{\pi}{2} i \left(\sum_{\{j|\epsilon_j=1\}} H_j - \sum_{\{j|\epsilon_j=-1\}} Y_j \right),$$

by $\mathfrak{sl}(2)$ -calculations we get $\text{Ad}(\exp A)(H_j) = \epsilon_j H_j$ and $\text{Ad}(\exp A)^2 = \text{id}$. Since $\theta = \text{Ad}(\exp \frac{\pi}{2} i \sum H_j)$, we see from the definition of σ that $\text{Ad}(\exp A) \circ \theta = \sigma \circ \text{Ad}(\exp A)$. Therefore $\text{Ad}(\exp A) : \mathfrak{g}_{\mathbb{C}} = (\mathfrak{g}_1^{\tau})_{\mathbb{C}} \rightarrow \mathfrak{k}_{1\mathbb{C}}$ is an isomorphism. Further, with $A \in \mathfrak{k}_1 + i\mathfrak{p}_1$, the subalgebra $\mathfrak{k}_1 + i\mathfrak{p}_1$ is trivially $\text{Ad}(\exp A)$ -invariant, and by this $\mathfrak{k} + i\mathfrak{p} = \mathfrak{g}_{\mathbb{C}} \cap (\mathfrak{k}_1 + i\mathfrak{p}_1)$ is mapped onto $\mathfrak{k}_{1\mathbb{C}} \cap (\mathfrak{k}_1 + i\mathfrak{p}_1) = \mathfrak{k}_1$. $\text{Ad}(\exp A)$ also maps the (-1) -eigenspace of θ onto the (-1) -eigenspace of σ , i.e. $\text{Ad}(\exp A)(i\mathfrak{p}_1) = \mathfrak{k}_1 \cap \mathfrak{q}_{1\sigma}$. As the map is complex linear, we finally get an isomorphism $\text{Ad}(\exp A) : \mathfrak{g} = \mathfrak{k} + \mathfrak{p} \rightarrow \mathfrak{k} + i(\mathfrak{k}_1 \cap \mathfrak{q}_{1\sigma}) = \mathfrak{k}_1^c$.

To complete the proof we have to show that $\eta \circ \text{Ad}(\exp A) = \text{Ad}(\exp A) \circ \eta$. But $H_j \in \mathfrak{q}_1$; therefore $Y_j = \frac{1}{2}[iH_j, X_j] \in \mathfrak{q}_1$ and $\eta(A) = -A$, which is enough, as $\text{Ad}(\exp A)$ is involutive.

For the last statement, with $\mathfrak{g}_{\mathbb{C}} \simeq (\mathfrak{k}_1^c)_{\mathbb{C}} = \mathfrak{k}_{1\mathbb{C}}$, the classification of the Hermitian symmetric algebras $(\mathfrak{g}_1, \mathfrak{k}_1, \theta)$ ([He 78, p. 518]) shows that $\mathfrak{g} = \mathfrak{z}(\mathfrak{g}) + [\mathfrak{g}, \mathfrak{g}]$, with $[\mathfrak{g}, \mathfrak{g}]$ simple (and $\mathfrak{z}(\mathfrak{g})$ one dimensional) except for $\mathfrak{g}_1 = \mathfrak{su}(n, n)$. For this Lie algebra in its standard realization we can choose

$$\eta\left(\begin{pmatrix} A & B \\ C & D \end{pmatrix}\right) = \begin{pmatrix} D & C \\ B & A \end{pmatrix},$$

cf. section 7.1, and from this it is also clear that $([\mathfrak{g}, \mathfrak{g}], \mathfrak{h}, \tau) \simeq (\tilde{\mathfrak{g}} \oplus \tilde{\mathfrak{g}}, \Delta(\tilde{\mathfrak{g}} \oplus \tilde{\mathfrak{g}}), \tilde{\tau})$ with $\tilde{\mathfrak{g}}$ simple and $\tilde{\tau} : \tilde{\mathfrak{g}} \oplus \tilde{\mathfrak{g}} \rightarrow \tilde{\mathfrak{g}} \oplus \tilde{\mathfrak{g}}, (X, Y) \mapsto (Y, X)$. $\square$

Example 5.6. Choosing all $\epsilon_i = +1$ (or -1), we get $\sigma = \theta$, and our “compactification” is the causal diffeomorphism $K_1/K_1 \cap H_1 \rightarrow G_1/P'$.

Example 5.7 (Causal compactification of $U(p, q)/O(p, q)$). With

$$G_1 = \left\{ \begin{pmatrix} A & B \\ \overline{B} & A \end{pmatrix} \in SU(n, n) \mid \begin{array}{l} A, B \in M(n \times n, \mathbb{C}), \\ {}^t \overline{A} A - {}^t B \overline{B} = 1, {}^t A \overline{B} = {}^t \overline{B} A \end{array} \right\} \simeq Sp(n, \mathbb{R}),$$

the Cartan involution $\theta(g) = {}^t \overline{g}^{-1}$, and the Cartan algebra $\mathfrak{t}$ given by the set of diagonal matrices in $\mathfrak{g}_1$, we choose $E_j = E_{j, j+n}$ and $E_{-j} = E_{j+n, j}$ for $j = 1, \dots, n$, where $E_{k, l}$ is the matrix with entry 1 in the k -th row and l -th column and otherwise zero. Then $\eta = \text{Ad}(\exp \frac{\pi}{2} i X^0)$ is given by complex conjugation, and for the subgroup of real matrices H_1 we have an isomorphism

$$H_1 \rightarrow GL(n, \mathbb{R}), \quad \begin{pmatrix} A & B \\ B & A \end{pmatrix} \mapsto A + B.$$

Choosing $\epsilon_j = +1$ for $j = 1, \dots, p$, and $\epsilon_j = -1$ for the remaining j , for the fixed point group of σ we get

$$G = \left\{ \begin{pmatrix} A_1 & 0 & 0 & B_2 \\ 0 & A_4 & B_3 & 0 \\ 0 & \overline{B}_2 & \overline{A}_1 & 0 \\ \overline{B}_3 & 0 & 0 & \overline{A}_4 \end{pmatrix} \mid \begin{array}{l} A_1 \in M(p \times p, \mathbb{C}), A_4 \in M(q \times q, \mathbb{C}), \\ B_2 \in M(p \times q, \mathbb{C}), B_3 \in M(q \times p, \mathbb{C}), \\ {}^t \overline{A} A - {}^t B \overline{B} = I, {}^t A \overline{B} = {}^t \overline{B} A \end{array} \right\},$$

where $p + q = n$. It is straightforward to check that

$$\rho : G \rightarrow U(p, q), \quad \begin{pmatrix} A & B \\ \overline{B} & A \end{pmatrix} \mapsto \begin{pmatrix} A_1 & B_2 \\ \overline{B}_3 & \overline{A}_4 \end{pmatrix},$$

is a group isomorphism, and that the subgroup of real matrices H is mapped onto $O(p, q)$.

We give the causal compactification in the Harish-Chandra realization of the Bergman-Šilov boundary. Remember that — with the usual choice of positive

roots compatible with our choice of the $E_{\pm j}$'s — the decomposition in P_1^+ , $K_{1\mathbb{C}}$ and P_1^- -components for G_1 is given by

$$\begin{pmatrix} A & B \\ \overline{B} & \overline{A} \end{pmatrix} = \begin{pmatrix} I & B\overline{A}^{-1} \\ 0 & I \end{pmatrix} \begin{pmatrix} A - B\overline{A}^{-1}\overline{B} & 0 \\ 0 & \overline{A} \end{pmatrix} \begin{pmatrix} I & 0 \\ \overline{A}^{-1}B & I \end{pmatrix},$$

and the operation of G_1 on P_1^+ by

$$\begin{pmatrix} A & B \\ \overline{B} & \overline{A} \end{pmatrix} \begin{pmatrix} I & Z \\ 0 & I \end{pmatrix} K_{1\mathbb{C}} P_1^- = \begin{pmatrix} I & (AZ + B)(\overline{B}Z + \overline{A})^{-1} \\ 0 & I \end{pmatrix} K_{1\mathbb{C}} P_1^-.$$

By these equations, and inverting ρ , we now easily get the causal compactification

$$U(p, q)/O(p, q) \rightarrow \check{S}_1,$$

$$\begin{pmatrix} A_1 & B_2 \\ \overline{B}_3 & \overline{A}_4 \end{pmatrix} O(p, q) \mapsto \begin{pmatrix} -A_1 & B_2 \\ B_3 & -A_4 \end{pmatrix} \begin{pmatrix} \overline{A}_1 & -\overline{B}_2 \\ -\overline{B}_3 & \overline{A}_4 \end{pmatrix}^{-1},$$

where the Bergman-Šilov boundary is given by $\check{S}_1 = \{Z \in M(n \times n, \mathbb{C}) \mid \overline{Z}Z = I, {}^t Z = Z\}$ ([Wo 72, He 78, p. 527]).

In fact, in proving Theorem 5.1 we used only the compactly causal structure of the Cayley space G_1/H_1 . In some cases it is therefore possible to construct causal compactifications while assuming G_1/H_1 to be only compactly causal.

Definition 5.8. Let $(\mathfrak{g}_1, \mathfrak{k}_1, \theta)$ be an irreducible orthogonal Lie algebra of non-compact Hermitian type. Let $\{\gamma_1, \dots, \gamma_r\}$ be a maximal system of strongly orthogonal noncompact roots, and $E_{\pm j} \in \mathfrak{g}_{1\mathbb{C}, \pm \gamma_j}$, $j = 1, \dots, r$, vectors with $E_j + E_{-j}$, $iE_j - iE_{-j} \in \mathfrak{g}_1$. Let τ be an involutive automorphism of $\mathfrak{g}_1$ commuting with θ . The set $\{E_{\pm j} \mid 1 \leq j \leq r\}$ is called a *system of first kind* with respect to τ if $\tau(E_{\pm j}) = -E_{\mp j}$ for all j . The set is called a *system of second kind* if $r = 2r'$, and $\tau(E_{\pm j}) = -E_{\mp(j+r')}$ or $\tau(E_{\pm(j+r')}) = -E_{\mp j}$, for $j = 1, \dots, r'$.

Theorem 5.9. Let (G_1, H_1, τ) , $H_1 := G_1^\tau$, be an irreducible compactly causal symmetric space with $G_1 \subset G_{1\mathbb{C}}$, $G_{1\mathbb{C}}$ the simply connected Lie group corresponding to $\mathfrak{g}_{1\mathbb{C}}$, and (G_1, K_1, θ) Hermitian symmetric of tube type. Define $\sigma := \eta \circ \tau$, and assume that one of the following cases holds:

- (i) The $\{E_{\pm j} \mid 1 \leq j \leq r\}$ form a system of first kind with respect to τ .
- (ii) The $\{E_{\pm j} \mid 1 \leq j \leq r\}$ form a system of second kind and $(\sum \mathbb{R}X_j) \cap \mathfrak{q}_{1\sigma}$ is maximal abelian in $\mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma}$.

Then $(G, H, \tau) := ((G_1^\sigma)_0, H_1 \cap (G_1^\sigma)_0, \tau)$ is a weak compactly causal subspace of (G_1, G_1^η, η) , and $\Phi : G/H \rightarrow G_1/P'$ is a causal compactification of it.

Proof. With $\tau(X^0) = \theta(X^0) = -X^0$ we see immediately that $\sigma^2 = \text{id}$ and that this involution commutes with τ and θ . From the Langlands decomposition we have $P' \simeq Z_{G_1}(X^0) \times Q_1^-$, and $\sigma(Q_1^-) = Q_1^+$ by $\sigma(X^0) = \tau(X^0) = -X^0$. On the other hand, by this the centralizer $Z_{G_1}(X^0)$ is σ - and τ -invariant, and therefore $P' \cap G = Z_{G_1}(X^0) \cap G$. In fact, our assumptions assure that (G_1, G_1^η, η) is of Cayley type with $G_1^\eta = Z_{G_1}(X^0)$. As $\sigma = \eta\tau$ and $H_1 = G_1^\tau$, we have $H \subset G_1^\eta$, and that $P' \cap G = H$. Now, by the same arguments as in the proof of Theorem 5.1, the map Φ is an injective submersion.

We show next the density of the image, again by using Matsuki's double coset decomposition. For systems of the first kind we have $\mathfrak{a} = \sum \mathbb{R}X_j \subset \mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma}$,

$W_\sigma(\mathfrak{a}) = W(\mathfrak{a})$, and the open double cosets are $Gw_vP(\mathfrak{a}, \Sigma^+)$, where w_v represents the element $v \in W(\mathfrak{a}, K_1 \cap G) \backslash W(\mathfrak{a})$.

In contrast, for systems of the second kind, by condition (i) of Proposition 4.1 the set $\mathfrak{a} \cap \mathfrak{q}_{1\sigma} = \sum_{j=1}^{r'} \mathbb{R}(X_j + X_{j+r'})$ must be maximal abelian in $\mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma}$. With $\mathfrak{a} \cap \mathfrak{g} = \sum \mathbb{R}(X_j - X_{j+r'})$ and $W(\mathfrak{a}) \simeq (\mathbb{Z}/2\mathbb{Z})^r \times S_r$, cf. Corollary 2.2, we see that $W_\sigma(\mathfrak{a})$ is generated by all permutations of the pairs $(X_1, X_{1+r'}), \dots, (X_{r'}, X_{r'+r'})$, the transpositions $\langle X_j, X_{j+r'} \rangle$ — which together form a semidirect product of permutations $S_{r'} \times (\mathbb{Z}/2\mathbb{Z})^{r'}$ — and the reflections which map $X_j + X_{j+r'}$ onto its negative and leave all vectors orthogonal to this one fixed. According to Theorem 2.1 and the following discussion, we have $\Sigma(\mathfrak{a}) \backslash \Sigma(\mathfrak{a} \cap \mathfrak{g}) = \{\pm \frac{1}{2}(\gamma_s^c + \gamma_t^c), \pm \frac{1}{2}(\gamma_u^c - \gamma_v^c) \mid 1 \leq s, t, u, v \leq r, u \neq v, |u - v| \neq r'\}$ and a $\sigma\theta$ -compatible positive subset in the sense of Proposition 4.1 can be defined using the lexicographic ordering $\gamma_1^c > \gamma_{1+r'}^c > \gamma_2^c > \gamma_{2+r'}^c > \dots > \gamma_{r'+r'}^c$. Of course, also in this case all other $\sigma\theta$ -compatible positive systems are $W_\sigma(\mathfrak{a})$ -conjugate to the given one.

We determine the open double cosets in GP' by Theorem 4.2. As in the proof of Theorem 5.1, a minimal parabolic subalgebra $\mathfrak{p}(\mathfrak{a}, w\Sigma(\mathfrak{a})^+)$, where $\langle \Sigma(\mathfrak{a})^+, \mathfrak{a}'_+ \rangle \subset \mathbb{R}^+$, lies in $\mathfrak{p}'$ if and only if $w \in W(\mathfrak{a})_{\mathfrak{m}'}$. Hence the classes of subalgebras in open orbits are parametrized by $W(\mathfrak{a})_{\mathfrak{m}'} \cap W_\sigma(\mathfrak{a})$. If we restrict our attention to open double cosets and divide out algebras within the same orbit, we get the diagram

$$\begin{array}{ccc} W(\mathfrak{a})_{\mathfrak{m}'} \cap W_\sigma(\mathfrak{a}) & \rightarrow & W(\mathfrak{a}, K_1 \cap G) \backslash W_\sigma(\mathfrak{a}) \\ \downarrow & & \nearrow \varphi \\ W(\mathfrak{a})_{\mathfrak{m}'} \cap W(\mathfrak{a}, K_1 \cap G) \backslash W(\mathfrak{a})_{\mathfrak{m}'} & & \cap W_\sigma(\mathfrak{a}) \end{array}$$

where we have to prove that φ is surjective. For spaces of first kind $Y_j \in \mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma}$, and the arguments given in the proof of Theorem 5.1 apply without any change. In the other case we have to show that the reflections with respect to $X_j + X_{j+r'}$ are elements of $W(\mathfrak{a}, K_1 \cap G)$. But now $X_j + X_{j+r'}$ and $Y_j + Y_{j+r'}$ are in $\mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma}$; therefore

$$[X_j + X_{j+r'}, Y_j + Y_{j+r'}] = -2i(H_j + H_{j+r'}) \in \mathfrak{k}_1 \cap \mathfrak{g}$$

and $\text{Ad}(\exp \frac{\pi}{2}i(H_j + H_{j+r'}))$ is a representative of the reflection we seek.

It remains to prove the causality of G/H and Φ . First of all we see that $\tau(Y_j) = Y_j$, and for systems of the second kind $\tau(Y_j) = Y_{j+r'}$ and $\tau(Y_{j+r'}) = Y_j$. In both cases we have $\tau(Y^0) = Y^0$. Thus $Z^0 = \frac{1}{4}[X^0, Y^0] \in \mathfrak{q}_1$, and this shows that $\tau(Y_\pm) = \tau(\frac{1}{2}Y^0 \mp Z^0) = Y_\mp$. With $\text{Ad}(\exp \frac{\pi}{2}iX^0)(Y^0) = -Y^0$, we then conclude that $\sigma(Y_\pm) = -Y_\mp$, showing the σ -invariance of the cone C_k . For the Cayley space (G_1, G_1^η, η) the compactly causal structure is defined by $C_k \subset \mathfrak{q}_1$. By $\sigma = \eta \circ \tau$ we have $\tau|_{\mathfrak{g}} = \eta|_{\mathfrak{g}}$ and $\tau|_G = \eta|_G$, and, applying Lemma 3.1, the cone $\text{pr}_{\mathfrak{g}}C_k \subset \mathfrak{q}_1 \cap \mathfrak{g}$ makes G/H into a weak compactly causal subspace of G_1/G_1^η . This also ensures that Φ is a causal map. $\square$

Example 5.10. For (G_1, H_1, τ) of Cayley type, $\tau(E_{\pm j}) = E_{\pm j}$. Defining $\tilde{E}_{+j} := iE_{+j}$ and $\tilde{E}_{-j} := -iE_{-j}$ gives a system of the first kind. Applying Theorem 5.9, we recover the “compactification” $\Phi : K_1/H_1 \cap K_1 \rightarrow \check{S}_1$.

Example 5.11 (Causal compactification of spaces of Cayley type). Let $(\mathfrak{g}_1, \mathfrak{h}_1, \tau)$ be compact causal of group type, i.e. $(\mathfrak{g}_1, \mathfrak{h}_1, \tau) = (\tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}}, \Delta(\tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}}), \tau)$, with involution $\tau : \tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}} \rightarrow \tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}}, (X, Y) \mapsto (Y, X)$, and $\Delta(\tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}}) := \{(X, X) \mid X \in \tilde{\mathfrak{g}}\}$. Then (G_1, K_1, θ_1) , $\theta_1 := \theta \times \theta$, is Hermitian symmetric of tube type if and only if $(\tilde{G}, \tilde{K}, \theta)$ is. If we choose the $\{\tilde{E}_{\pm j} \mid j = 1, \dots, r'\}$ in $\tilde{\mathfrak{g}}_{\mathbb{C}}$ with the usual properties, then

$E_{\pm j} := (\tilde{E}_{\pm j}, 0)$ and $E_{\pm(j+r')} := (0, -\tilde{E}_{\mp j})$ form a system of the second kind. Theorem 5.9 then gives $G = \{(g, \tilde{\eta}(g)) \mid g \in \tilde{G}\}$, where $\tilde{\eta} := \text{Ad}(\exp \frac{\pi}{2} i \tilde{X}^0)$, and therefore $(G, H, \tau) \simeq (\tilde{G}, \tilde{G}^{\tilde{\eta}}, \tilde{\eta})$, which is of Cayley type. This gives us a variant of the causal compactification for Cayley spaces already constructed by different methods, cf. [ØØ 97, Ko 93, Ka 87].

Example 5.12 (Causal compactification of $SO^*(2n)/O(n, \mathbb{C})$). For

$$G_1 = \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in SL(2n, \mathbb{C}) \mid \begin{matrix} {}^t A \overline{A} - {}^t C \overline{C} = I, {}^t A \overline{B} = {}^t C \overline{D} \\ {}^t D \overline{D} - {}^t B \overline{B} = I \end{matrix} \right\} = SU(n, n),$$

let τ be complex conjugation and correspondingly $H_1 \simeq O(n, n)$, the subgroup of real matrices. (Note that (G_1, H_1, τ) , as the ρ -image of the semisimple symmetric space $([G, G], H, \tau)$ defined in Example 5.7, is compactly causal.) Then $E_{+j} = iE_{j,j+n}$ and $E_{-j} = -iE_{j+n,j}$ form a system of first kind. The involution σ has as fixed point group

$$G = \left\{ \begin{pmatrix} A & B \\ -\overline{B} & \overline{A} \end{pmatrix} \in SL(2n, \mathbb{C}) \mid \begin{matrix} {}^t \overline{A} A - {}^t \overline{B} B = I \\ {}^t A \overline{B} + {}^t \overline{B} A = 0 \end{matrix} \right\} \simeq SO^*(2n)$$

([He 78, p. 527]), and H is the subgroup of all real matrices with

$$H \rightarrow O(n, \mathbb{C}), \quad \begin{pmatrix} A & B \\ -B & A \end{pmatrix} \mapsto A + iB,$$

an isomorphism. In the Harish-Chandra realization we now get

$$SO^*(2n)/O(n, \mathbb{C}) \rightarrow \check{S}_1$$

$$\begin{pmatrix} A & B \\ -\overline{B} & \overline{A} \end{pmatrix} H \mapsto (-iA + B)(i\overline{B} + \overline{A})^{-1},$$

where $\check{S}_1 = \{Z \in M(n \times n, \mathbb{C}) \mid {}^t \overline{Z} Z = I\}$ is the Bergman-Šilov boundary of $SU(n, n)/S(U(n) \times U(n))$.

6. COMPACTLY CAUSAL SPACES WITH NO COMPACTIFICATION

Using our results of the last section, it is easy to construct causal compactifications for the spaces listed in the introduction. For those with “no compactification” causal compactifications can exist for some small n from the existence of special isomorphisms, cf. [He 78, p. 519f]. We will prove here the nonexistence of further compactifications as symmetric orbits by making use of the classification of open symmetric orbits in symmetric R -spaces ([Ma 73]).

For this we note first that the Bergman-Šilov boundary $\check{S}_1$ of a Hermitian symmetric space of tube type is such a space, since, by Theorem 2.3, it is an R -space, i.e., a quotient G_1/P' of a semisimple symmetric group by a parabolic subgroup, and in addition $(K_1, K_1 \cap P', \eta)$ and $\check{S}_1 = K_1/K_1 \cap P'$ possess K_1 -invariant Riemannian structures ([KW 65, Thm. 4.9]). An open orbit of $G \subset G_1$ is called symmetric if it is a symmetric subspace of G_1/P' . We remark that our causal compactifications already constructed are symmetric orbits.

Proposition 6.1. *For the following compactly causal symmetric Lie algebras $(\mathfrak{g}, \mathfrak{h})$ there exists no compactly causal space (G_1, H_1, τ) such that the subspace (G, H, τ) associated to $(\mathfrak{g}, \mathfrak{h}) \subset (\mathfrak{g}_1, \mathfrak{h}_1)$ is a compactly causal subspace and $\Phi : G/H \rightarrow G_1/P'$, with P' appropriately chosen, a causal compactification as symmetric orbit.*

$\mathfrak{g}$	$\mathfrak{h}$	parameters
$\mathfrak{so}(2, p+q)$	$\mathfrak{so}(p, 1) \oplus \mathfrak{so}(1, q)$	$\min(p, q) \geq 2, (p, q) \neq (2, 2), (3, 3)$
$\mathfrak{e}_{6(-14)}$	$\mathfrak{sp}(2, 2)$	
$\mathfrak{e}_{7(-25)}$	$\mathfrak{su}^*(8)$	
$\mathfrak{so}(2, n) \oplus \mathfrak{so}(2, n)$	$\mathfrak{so}(2, n)$	$n = 5$ or $n \geq 7$
$\mathfrak{e}_{6(-14)} \oplus \mathfrak{e}_{6(-14)}$	$\mathfrak{e}_{6(-14)}$	
$\mathfrak{e}_{7(-25)} \oplus \mathfrak{e}_{7(-25)}$	$\mathfrak{e}_{7(-25)}$	

Proof. Assume to the contrary that $G/H \subset G_1/P'$ is a symmetric orbit. When we decompose $G_1/P' = \tilde{S}_1 \times \dots \times \tilde{S}_n$ into irreducible symmetric R -spaces, $G = \tilde{G}_1 \times \dots \times \tilde{G}_m$ decomposes also into factors, where each factor gives a symmetric orbit in $\tilde{S}_j$ or $\tilde{S}_j \times \tilde{S}_{j+1}$, with $\tilde{S}_j \simeq \tilde{S}_{j+1}$ ([Ma 73]). By looking at the classification² of symmetric orbits $G/H \rightarrow \tilde{S}_1$, we see that no space on our list — with the exception of the already mentioned low dimensional cases — possesses such a compactification. For the last three cases with $G = G_0 \times G_0$, a compactification may be constructed from maps $G_0 \times \{e\} \rightarrow \tilde{S}_1$ and $\{e\} \times G_0 \rightarrow \tilde{S}_2$. However then $\mathfrak{h} \simeq \mathfrak{g}_0$ would not be simple, which is the case only for $\mathfrak{so}(2, 2) \simeq \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{sl}(2, \mathbb{R})$.

It remains to check the case $G \rightarrow \tilde{S}_1 \simeq \tilde{S} \times \tilde{S}$. For spaces of tube type the dimension of the Bergman-Šilov boundary $\tilde{S}_1$ is $\frac{1}{2} \dim(G_1/K_1)$, and we get the necessary condition $\dim \mathfrak{g} - \dim \mathfrak{h} = \frac{1}{2}(\dim \mathfrak{g}_1 - \dim \mathfrak{k}_1)$. Moreover, for G simple the only possible imbedding is the diagonal imbedding $G \rightarrow \Delta(G) \subset G \times G$ or a modification of this map as in Example 5.11 ([Ma 73]). For the first three groups of our list we therefore have the condition $\dim \mathfrak{h} = \dim \mathfrak{k}$, and this equation can only be solved for $\mathfrak{so}(2, p+q)$ with $p = 1$ or $q = 1$, recovering the compactification for the spaces of Cayley type. For the last three groups a compactification would give us an imbedding $\mathfrak{g}_0 \oplus \mathfrak{g}_0 \hookrightarrow \mathfrak{g}_1 \oplus \mathfrak{g}_1$, with $(\mathfrak{g}_1, \mathfrak{k}_1, \theta)$ Hermitian symmetric of tube type, such that $\Delta(\mathfrak{g}_0 \oplus \mathfrak{g}_0)$ goes into the sum $\mathfrak{p}' \oplus \tilde{\mathfrak{p}}'$ of two maximal parabolic subalgebras. With the Langlands decomposition $\mathfrak{p}' = \mathfrak{m}' + \mathbb{R}X^0 + \mathfrak{q}_1^-$ we get from this an imbedding $\mathfrak{g}_0 \hookrightarrow \mathfrak{m}'$ ([Va 84, Cor. 3.14.3]) with θ -invariant image ([Wa 72, Lemma 1.1.5.5]). Therefore the rank and real rank of $\mathfrak{m}'$ are restricted from below by the corresponding quantities of $\mathfrak{g}_0$, excluding $\mathfrak{g}_1 = \mathfrak{so}(2, k)$. Using our equation on the dimensions, we are left with $\mathfrak{g}_0 = \mathfrak{so}(2, n)$ and $\mathfrak{g}_1 = \mathfrak{sp}(k, \mathbb{R})$ or $\mathfrak{g}_1 = \mathfrak{so}^*(4k)$, $k \geq 3$. For the first $\mathfrak{g}_1$, equality of dimensions leads to $(n+1)(n+2) = 2k(k+1)$, whereas from $\mathfrak{g}_0 \hookrightarrow \mathfrak{m}' = \mathfrak{sl}(k, \mathbb{R})$ necessarily $(n+1)(n+2) \leq 2(k^2 - 1)$, a contradiction. For the second $\mathfrak{g}_1$, comparing the dimensions gives $n \geq 2k$; but $\mathfrak{g}_0 \hookrightarrow \mathfrak{m}' = \mathfrak{su}^*(2k)$, mapping maximal compact subalgebras into compact subalgebras, gives $1 + [\frac{n}{2}] < k$, a contradiction, too. $\square$

7. CAUSAL COMPACTIFICATIONS FOR THE CLASSICAL CAUSAL SPACES

To be complete, we give here some details for the compactification of the classical symmetric spaces not already treated in the various examples. For the definition of these spaces as well as for our notation we follow [He 78, Chapter X].

²According to W. Bertram (personal communication), the classification in [Ma 73] is not complete. However, the part of the classification needed by us is not affected by this.

7.1. The compactification of $S(U(p, q) \times U(q, p))/SU(p, q)$. The group

$$G_1 = \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in SL(2n, \mathbb{C}) \left| \begin{array}{l} {}^t A \bar{A} - {}^t C \bar{C} = I, \quad {}^t A \bar{B} = {}^t C \bar{D} \\ {}^t D \bar{D} - {}^t B \bar{B} = I \end{array} \right. \right\} = SU(n, n)$$

verifies the conditions of Corollary 5.3. The real rank of $SU(n, n)$ is n , and we can use the same system of strongly orthogonal roots as for $Sp(n, \mathbb{R})$ in Example 5.7. However, η is no longer given by complex conjugation, but

$$\eta\left(\begin{pmatrix} A & B \\ C & D \end{pmatrix}\right) = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} = \begin{pmatrix} D & C \\ B & A \end{pmatrix}.$$

Then

$$H_1 = \left\{ \begin{pmatrix} A & B \\ B & A \end{pmatrix} \in SL(2n, \mathbb{C}) \left| \begin{array}{l} {}^t A \bar{A} - {}^t B \bar{B} = I, \\ {}^t A \bar{B} = {}^t B \bar{A} \end{array} \right. \right\} \simeq SL(n, \mathbb{C}) \times \mathbb{R}^+ I,$$

where the isomorphism is the same as that for H_1 in Example 5.7.

Again let $\epsilon_j = +1$ for $j = 1, \dots, p$, and $\epsilon_j = -1$ for $j = p+1, \dots, n$. Then for the fixed point group of σ we get

$$\begin{aligned} G &= \left\{ \begin{pmatrix} A_1 & 0 & 0 & B_2 \\ 0 & A_4 & B_3 & 0 \\ 0 & C_2 & D_1 & 0 \\ C_3 & 0 & 0 & D_4 \end{pmatrix} \in SU(n, n) \left| \begin{array}{l} A_1, D_1 \in M(p \times p, \mathbb{C}) \\ A_4, D_4 \in M(q \times q, \mathbb{C}) \\ B_2, C_2 \in M(q \times p, \mathbb{C}) \\ B_3, C_3 \in M(p \times q, \mathbb{C}) \end{array} \right. \right\} \\ &\simeq \left\{ \left(\begin{pmatrix} A_1 & B_2 \\ C_3 & D_4 \end{pmatrix}, \begin{pmatrix} A_4 & B_3 \\ C_2 & D_1 \end{pmatrix} \right) \left| \begin{array}{l} {}^t \bar{A}_k A_k - {}^t \bar{C}_m C_m = I, \quad {}^t \bar{D}_n D_n - {}^t \bar{B}_l B_l = I \\ {}^t A_k \bar{B}_l = {}^t C_m \bar{D}_n, \text{ product of determinants } 1 \\ (k, l, m, n) \in \{(1, 2, 3, 4), (4, 3, 2, 1)\} \end{array} \right. \right\} \\ &= S(U(p, q) \times U(q, p)). \end{aligned}$$

The subgroup $H = G \cap H_1$ maps under the same isomorphism onto

$$\tilde{H} = \left\{ \left(\begin{pmatrix} A_1 & B_2 \\ B_3 & A_4 \end{pmatrix}, \begin{pmatrix} A_4 & B_3 \\ B_2 & A_1 \end{pmatrix} \right) \left| \dots \right. \right\} \simeq SU(p, q),$$

and we get the causal compactification

$$S(U(p, q) \times U(q, p)) / \tilde{H} \rightarrow \check{S}_1,$$

$$\left(\begin{pmatrix} A_1 & B_2 \\ C_3 & D_4 \end{pmatrix}, \begin{pmatrix} A_4 & B_3 \\ C_2 & D_1 \end{pmatrix} \right) \tilde{H} \mapsto \left(\begin{pmatrix} -A_1 & B_2 \\ B_3 & -A_4 \end{pmatrix} \begin{pmatrix} D_1 & -C_2 \\ -C_3 & D_4 \end{pmatrix}^{-1} \right),$$

where the Bergman-Šilov boundary of $SU(n, n)/S(U(n) \times U(n))$ in the Harish-Chandra realization is given by $\check{S}_1 = \{Z \in M(n \times n, \mathbb{C}) \mid {}^t Z \bar{Z} = I\}$ ([Wo 72, He 78]).

7.2. The compactification of $U(2p, 2q)/Sp(p, q)$. We apply Corollary 5.3 to

$$G_1 = \left\{ \begin{pmatrix} A & B \\ -\bar{B} & \bar{A} \end{pmatrix} \in SL(4n, \mathbb{C}) \left| \begin{array}{l} {}^t A \bar{A} - {}^t \bar{B} B = I \\ {}^t A \bar{B} + {}^t \bar{B} A = 0 \end{array} \right. \right\} \simeq SO^*(4n),$$

with Cartan algebra $\mathfrak{t} = \{i \operatorname{diag}(x_1, \dots, x_{2n}, -x_1, \dots, -x_{2n}) \mid x_j \in \mathbb{R}\}$, and

$$E_j = \begin{pmatrix} 0 & E_{j,j+n} - E_{j+n,j} \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad E_{-j} = \begin{pmatrix} 0 & 0 \\ -E_{j,j+n} + E_{j+n,j} & 0 \end{pmatrix},$$

$j = 1, \dots, n$. With the given data we have

$$\eta\left(\begin{pmatrix} A & B \\ -\bar{B} & \bar{A} \end{pmatrix}\right) = \begin{pmatrix} -J_n \bar{A} J_n & -J_n \bar{B} J_n \\ J_n B J_n & -J_n A J_n \end{pmatrix}, \quad \text{where } J_n := \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}.$$

The fixed point group of η is

$$H_1 = \left\{ \left(\begin{array}{cccc} A_1 & A_2 & B_1 & B_2 \\ -\overline{A}_2 & \overline{A}_1 & -\overline{B}_2 & \overline{B}_1 \\ -\overline{B}_1 & -\overline{B}_2 & \overline{A}_1 & \overline{A}_2 \\ B_2 & -B_1 & -A_2 & A_1 \end{array} \right) \mid \begin{array}{l} {}^t A \overline{A} - \overline{B} B = I \\ {}^t A \overline{B} + \overline{B} A = 0 \end{array} \right\},$$

which is isomorphic to $SU^*(2n)$ by mapping the element of the above given form to

$$\begin{pmatrix} A_1 + B_2 & A_2 - B_1 \\ -\overline{A}_2 + \overline{B}_1 & \overline{A}_1 + \overline{B}_2 \end{pmatrix}.$$

With $\epsilon_j = 1$ for $j = 1, \dots, p$, and $\epsilon_j = -1$ for $j = p + 1, \dots, n$, we then have $\sigma = \text{Ad}(\text{diag}(I_{p,q}, I_{p,q}, -I_{p,q}, -I_{p,q}))$, with $I_{p,q} := \text{diag}(1, \dots, 1, -1, \dots, -1)$, and for the set of fixed points

$$G = \left\{ \left(\begin{array}{cccc} A_{11} & 0 & A_{21} & 0 \\ 0 & A_{14} & 0 & A_{24} \\ A_{31} & 0 & A_{41} & 0 \\ 0 & A_{34} & 0 & A_{44} \\ \dots & & & \\ 0 & -\overline{B}_{12} & 0 & -\overline{B}_{22} \\ -\overline{B}_{13} & 0 & -\overline{B}_{23} & 0 \\ 0 & -\overline{B}_{32} & 0 & -\overline{B}_{42} \\ -\overline{B}_{33} & 0 & -\overline{B}_{43} & 0 \end{array} \right) \mid \begin{array}{l} A_{11}, A_{21}, A_{31}, A_{41} \in M(p \times p) \\ A_{14}, A_{24}, A_{34}, A_{44} \in M(q \times q) \\ B_{12}, B_{22}, B_{32}, B_{42} \in M(p \times q) \\ B_{13}, B_{23}, B_{33}, B_{43} \in M(q \times p) \\ {}^t A \overline{A} - \overline{B} B = I, \\ {}^t A \overline{B} + \overline{B} A = 0 \end{array} \right\}$$

$$\simeq \left\{ \left(\begin{array}{cccc} A_{11} & A_{21} & B_{12} & B_{22} \\ A_{31} & A_{41} & B_{32} & B_{42} \\ -\overline{B}_{13} & -\overline{B}_{23} & \overline{A}_{14} & \overline{A}_{24} \\ -\overline{B}_{33} & -\overline{B}_{43} & \overline{A}_{34} & \overline{A}_{44} \end{array} \right) \mid \dots \right\} = U(2p, 2q).$$

The image $\tilde{H}$ of H is isomorphic to $Sp(p, q)$ via conjugation with

$$\begin{pmatrix} i1_p & 0 & 0 & 0 \\ 0 & 0 & 1_p & 0 \\ 0 & 1_q & 0 & 0 \\ 0 & 0 & 0 & i1_q \end{pmatrix},$$

and for the causal compactification we find that

$$SU(2p, 2q)/\tilde{H} \rightarrow \check{S}_1,$$

$$\begin{pmatrix} A_{11} & A_{21} & B_{12} & B_{22} \\ A_{31} & A_{41} & B_{32} & B_{42} \\ -\overline{B}_{13} & -\overline{B}_{23} & \overline{A}_{14} & \overline{A}_{24} \\ -\overline{B}_{33} & -\overline{B}_{43} & \overline{A}_{34} & \overline{A}_{44} \end{pmatrix} \tilde{H}$$

$$\mapsto \begin{pmatrix} A_{21} & B_{12} & -A_{11} & B_{22} \\ B_{13} & A_{24} & B_{23} & -A_{14} \\ A_{41} & B_{32} & -A_{31} & B_{42} \\ B_{33} & A_{44} & B_{43} & -A_{34} \end{pmatrix} \begin{pmatrix} \overline{A}_{11} & -\overline{B}_{22} & \overline{A}_{21} & \overline{B}_{12} \\ -\overline{B}_{23} & \overline{A}_{14} & \overline{B}_{13} & \overline{A}_{24} \\ \overline{A}_{31} & -\overline{B}_{42} & \overline{A}_{41} & \overline{B}_{32} \\ -\overline{B}_{43} & \overline{A}_{34} & \overline{B}_{33} & \overline{A}_{44} \end{pmatrix}^{-1},$$

with Bergman-Šilov boundary $\check{S}_1 = \{Z \in M(2n \times 2n, \mathbb{C}) \mid {}^t Z = -Z, {}^t Z \overline{Z} = 1\}$.

7.3. The compactification of $SO_0(2, n-1) \times SO(2)/(O(1) \times SO_0(1, n-1))$. We start with $G_1 = SO(2, n+1)$ as in Example 5.2, again assume $n \geq 2$, and choose the same $E_{\pm j}$ and η . We have to replace H_1 by $Z_{G_1}(X^0)$, since $G_{1\mathbb{C}}$ is not simply connected. Applying Corollary 5.3, with $\epsilon_1 = 1 = -\epsilon_2$, we have $\sigma = \text{Ad}(I_{n,2})$ and correspondingly

$$G = \left\{ \begin{pmatrix} A & B' & 0 \\ C' & D'_1 & 0 \\ 0 & 0 & D_4 \end{pmatrix} \in SL(n+3, \mathbb{R}) \mid \dots \right\} \simeq SO_0(2, n-1) \times SO(2).$$

The centralizer $Z_G(X^0)$ is easily determined from $Z_{G_1}(X^0)$, as given in Example 5.2, to be

$$Z_G(X^0) = \left\{ \begin{pmatrix} \epsilon & 0 & 0 & 0 & 0 \\ 0 & \epsilon a_4 & B'_3 & 0 & 0 \\ 0 & C'_2 & D'_1 & 0 & 0 \\ 0 & 0 & 0 & \epsilon & 0 \\ 0 & 0 & 0 & 0 & \epsilon \end{pmatrix} \in SL(n+3, \mathbb{R}) \mid \begin{matrix} \epsilon \in \{\pm 1\}, \\ \begin{pmatrix} a_4 & B'_3 \\ C'_2 & D'_1 \end{pmatrix} \in O(1, n) \end{matrix} \right\},$$

and therefore $G/Z_G(X^0) \simeq SO_0(2, n+1) \times SO(2)/(O(1) \times SO_0(1, n-1))$.

7.4. Causal compactification of $Sp(n, \mathbb{R}) \times Sp(n, \mathbb{R})/Sp(n, \mathbb{R})$. We begin with $G_1 \simeq Sp(2n, \mathbb{R})$ and the Cartan involution $\theta(g) = {}^t\bar{g}^{-1}$ as in Example 5.7. But as root vectors we choose now

$$E_j = \frac{1}{2} \begin{pmatrix} 0 & E_{j,j} + E_{j+n,j} + E_{j,j+n} + E_{j+n,j+n} \\ 0 & 0 \end{pmatrix}$$

and

$$E_{j+n} = \frac{1}{2} \begin{pmatrix} 0 & -E_{j,j} + E_{j+n,j} + E_{j,j+n} - E_{j+n,j+n} \\ 0 & 0 \end{pmatrix},$$

for $j = 1, \dots, n$, and $E_{-j} := -\theta(E_j)$ for all j . With

$$\sigma = \text{Ad} \begin{pmatrix} I_{n,n} & 0 \\ 0 & I_{n,n} \end{pmatrix}$$

this involution commutes with θ and

$$\eta = \text{Ad}(\exp(\frac{\pi}{2}iX^0)) = \text{Ad} \begin{pmatrix} 0 & L_n \\ L_n & 0 \end{pmatrix}, \text{ where } L_n := \begin{pmatrix} 0 & I_n \\ I_n & 0 \end{pmatrix},$$

giving the fixed point group

$$G = \left\{ \begin{pmatrix} A_1 & 0 & B_1 & 0 \\ 0 & A_4 & 0 & B_4 \\ \bar{B}_1 & 0 & \bar{A}_1 & 0 \\ 0 & \bar{B}_4 & 0 & \bar{A}_4 \end{pmatrix} \mid \begin{matrix} {}^tA_i\bar{A}_i - {}^t\bar{B}_iB_i = I \\ {}^tA_i\bar{B}_i = {}^t\bar{B}_iA_i \\ i \in \{1, 4\} \end{matrix} \right\} \simeq Sp(n, \mathbb{R}) \times Sp(n, \mathbb{R}).$$

For the associated dual group we have

$$G_a = \left\{ \begin{pmatrix} A_1 & 0 & 0 & B_2 \\ 0 & A_4 & B_3 & 0 \\ 0 & \bar{B}_2 & \bar{A}_1 & 0 \\ \bar{B}_3 & 0 & 0 & \bar{A}_4 \end{pmatrix} \mid \dots \right\} \simeq U(n, n)$$

and, with n the real rank of $U(n, n)$, condition (ii) of Theorem 5.1 can be verified immediately. Finally,

$$H = \left\{ \left(\begin{pmatrix} A_1 & 0 & B_1 & 0 \\ 0 & \bar{A}_1 & 0 & \bar{B}_1 \\ \bar{B}_1 & 0 & \bar{A}_1 & 0 \\ 0 & B_1 & 0 & A_1 \end{pmatrix} \right) \left| \begin{array}{l} {}^t A_1 \bar{A}_1 - {}^t \bar{B}_1 B_1 = I, \\ {}^t A_1 \bar{B}_1 = {}^t \bar{B}_1 A_1 \end{array} \right. \right\} \simeq Sp(n, \mathbb{R}).$$

Putting things together, we get the causal compactification in the Harish-Chandra realization

$$Sp(n, \mathbb{R}) \times Sp(n, \mathbb{R}) / \Delta(Sp \times Sp) \rightarrow \check{S}_1,$$

$$\left(\left(\begin{pmatrix} A_1 & B_1 \\ \bar{B}_1 & \bar{A}_1 \end{pmatrix}, \begin{pmatrix} \bar{A}_4 & \bar{B}_4 \\ B_4 & A_4 \end{pmatrix} \right) \Delta(Sp \times Sp) \mapsto \begin{pmatrix} B_1 & -A_1 \\ -A_4 & B_4 \end{pmatrix} \begin{pmatrix} \bar{A}_1 & -\bar{B}_1 \\ -\bar{B}_4 & \bar{A}_4 \end{pmatrix}^{-1}, \right.$$

where the Bergman-Šilov boundary is $\check{S}_1 = \{Z \in M(2n \times 2n, \mathbb{C}) \mid \bar{Z}Z = I, {}^t Z = Z\}$, and $\Delta(Sp \times Sp) := \{(g, g) \mid g \in Sp(n, \mathbb{R})\}$, the diagonal in $Sp(n, \mathbb{R}) \times Sp(n, \mathbb{R})$.

We note that our (G_1, H_1) is a Cayley subspace of $(SU(2n, 2n), SL(2n, \mathbb{C}) \cdot \mathbb{R}^+ I)$. If we construct the causal compactification of $S(U(n, n) \times U(n, n))/SU(n, n)$ as in section 7.1, starting from $G_1 = SU(2n, 2n)$, then our G/H is a causal subspace of $S(U(n, n) \times U(n, n))/SU(n, n)$, and the causal compactification of G/H constructed above is that of $S(U(n, n) \times U(n, n))/SU(n, n)$ restricted to G/H .

7.5. The compactification of $SO^*(2n) \times SO^*(2n)/SO^*(2n)$. For

$$G_1 = \left\{ \begin{pmatrix} A & B \\ -\bar{B} & \bar{A} \end{pmatrix} \in SL(4n, \mathbb{C}) \left| \begin{array}{l} {}^t A \bar{A} - {}^t \bar{B} B = I \\ {}^t A \bar{B} + {}^t \bar{B} A = 0 \end{array} \right. \right\} \simeq SO^*(4n),$$

we choose $E_j = E_{j, j+3n} - E_{j+n, j+2n}$ and $E_{-j} = E_{j+3n, j} - E_{j+2n, j+n}$, for $j = 1, \dots, n$. With θ and σ as in section 7.4 we get

$$G = \left\{ \left(\begin{pmatrix} A_1 & 0 & B_1 & 0 \\ 0 & A_4 & 0 & B_4 \\ -\bar{B}_1 & 0 & \bar{A}_1 & 0 \\ 0 & -\bar{B}_4 & 0 & \bar{A}_4 \end{pmatrix} \right) \left| \begin{array}{l} {}^t A_i \bar{A}_i - {}^t \bar{B}_i B_i = I \\ {}^t A_i \bar{B}_i + {}^t \bar{B}_i A_i = 0 \\ i \in \{1, 4\} \end{array} \right. \right\} \\ \simeq SO^*(2n) \times SO^*(2n),$$

and our root vectors give X_j and Y_j , which satisfy condition (i) of Theorem 5.1. The involution

$$\eta = \text{Ad}(\exp i \frac{\pi}{2} X^0) = \text{Ad} \begin{pmatrix} 0 & J_n \\ -J_n & 0 \end{pmatrix}$$

makes (G_1, H_1, η) into a Cayley space, and it commutes with σ . Also,

$$H = \left\{ \left(\begin{pmatrix} A_1 & 0 & B_1 & 0 \\ 0 & \bar{A}_1 & 0 & \bar{B}_1 \\ -\bar{B}_1 & 0 & \bar{A}_1 & 0 \\ 0 & -B_1 & 0 & A_1 \end{pmatrix} \right) \left| \begin{array}{l} {}^t A_1 \bar{A}_1 - {}^t \bar{B}_1 B_1 = I, \\ {}^t A_1 \bar{B}_1 + {}^t \bar{B}_1 A_1 = 0 \end{array} \right. \right\} \simeq SO^*(2n),$$

so that for the causal compactification we now get, similarly to the last section,

$$SO^*(2n) \times SO^*(2n) / \Delta(SO^* \times SO^*) \rightarrow \check{S}_1,$$

$$\left(\left(\begin{pmatrix} A_1 & B_1 \\ -\bar{B}_1 & \bar{A}_1 \end{pmatrix}, \begin{pmatrix} \bar{A}_4 & \bar{B}_4 \\ -B_4 & A_4 \end{pmatrix} \right) \Delta(SO^* \times SO^*) \mapsto \begin{pmatrix} B_1 & -A_1 \\ A_4 & B_4 \end{pmatrix} \begin{pmatrix} \bar{A}_1 & \bar{B}_1 \\ -\bar{B}_4 & \bar{A}_4 \end{pmatrix}^{-1}, \right.$$

with $\check{S}_1$ as in section 7.2 and $\Delta(SO^* \times SO^*)$ again the diagonal subgroup.

7.6. The compactification of $Sp(2n, \mathbb{R})/Sp(n, \mathbb{C})$. The group $G_1 = SU(2n, 2n)$ together with

$$H_1 = \left\{ \left(\begin{pmatrix} A_{11} & A_{21} & B_{12} & B_{22} \\ -\bar{A}_{21} & \bar{A}_{11} & -\bar{B}_{12} & \bar{B}_{22} \\ -\bar{B}_{13} & -\bar{B}_{23} & \bar{A}_{14} & \bar{A}_{24} \\ B_{23} & -B_{13} & -A_{24} & A_{14} \end{pmatrix} \in SL(4n, \mathbb{C}) \mid \dots \right\} \simeq Sp(n, n),$$

and the involution

$$\tau(g) = \text{Ad} \begin{pmatrix} J_p & 0 \\ 0 & J_p \end{pmatrix} (\bar{g}),$$

make up a compactly causal space (G_1, H_1, τ) . Indeed, G_1/H_1 is isomorphic to the semisimple symmetric subspace contained in the weak compactly causal space G/H constructed in section 7.2. The real rank is $2n$, and $E_j := iE_{j,j+3n}$, $E_{-j} := -iE_{j+3n,j}$, $E_{j+n} := iE_{j+2n,j+n}$ together with $E_{-(j+n)} := -iE_{j+n,j+2n}$, for $j = 1, \dots, n$, form a system of the second kind. Defining σ as in Theorem 5.9, we find immediately that

$$G = \left\{ \begin{pmatrix} A & B \\ \bar{B} & \bar{A} \end{pmatrix} \in SU(2n, 2n) \right\} \simeq Sp(2n, \mathbb{R})$$

and

$$H = \left\{ \left(\begin{pmatrix} A_{11} & A_{21} & B_{12} & B_{22} \\ -\bar{A}_{21} & \bar{A}_{11} & -\bar{B}_{22} & \bar{B}_{12} \\ \bar{B}_{12} & \bar{B}_{22} & \bar{A}_{11} & \bar{A}_{21} \\ -B_{22} & B_{12} & -A_{21} & A_{11} \end{pmatrix} \in SL(4n, \mathbb{C}) \mid \dots \right\} \simeq Sp(n, \mathbb{C}),$$

the last isomorphism being given by mapping an element of the given form onto

$$\begin{pmatrix} A_{11} + iB_{22} & A_{21} - iB_{12} \\ -\bar{A}_{21} + i\bar{B}_{12} & \bar{A}_{11} + i\bar{B}_{22} \end{pmatrix}.$$

Since the associated group $G_a := G^{\sigma\theta} \simeq SO^*(4n)$ has real rank n , the cited theorem gives us a causal compactification of (G, H, τ) .

7.7. The compactification of $SO_0(2, q) \times SO(p+1)/(SO_0(1, q) \times O(p))$. With $G_1 = SO(2, n+1)$ we choose $\tau := \text{Ad}(\text{diag}(1, -1, \dots, -1, 1, \dots, 1))$, where -1 ranges from the second to the $(p+2)$ -th position. Then

$$H_1 = \left\{ \left(\begin{pmatrix} a_1 & 0 & 0 & B_2 \\ 0 & a_4 & B_3 & 0 \\ 0 & C_2 & D_1 & 0 \\ C_3 & 0 & 0 & D_4 \end{pmatrix} \in SL(n+3, \mathbb{R}) \mid \dots \right\} \simeq S(O(1, p) \times O(1, q+1))$$

and therefore G_1/H_1 is compactly causal ([HÓ 96, p. 89]). With the symmetric matrices

$$\tilde{E}_1 := \frac{1}{2} \begin{pmatrix} 0 & \dots & -i & -1 \\ \vdots & & -1 & i \\ -i & -1 & & \\ -1 & i & & \vdots \\ & & \dots & 0 \end{pmatrix} \text{ and } \tilde{E}_2 := \frac{1}{2} \begin{pmatrix} 0 & \dots & i & -1 \\ \vdots & & 1 & i \\ i & 1 & & \\ -1 & i & & \vdots \\ & & \dots & 0 \end{pmatrix},$$

where the first nonzero entries $-i$ and i occur in the first row in the $(p+1)$ -th column, and $\tilde{E}_{-j} := -\theta(\tilde{E}_j)$, we have $\tau(\tilde{E}_{\pm j}) = \tilde{E}_{\mp j}$. Similarly to Example 5.10,

we define $E_{\pm j} := \pm i \tilde{E}_{\pm j}$ so that we get a system of the first kind. Applying Theorem 5.9 gives $\sigma = \text{Ad}(\text{diag}(1, 1, -1, \dots, -1, 1, \dots, 1))$, with -1 now from the third up to the $(p+3)$ -th position, and

$$G = \left\{ \left(\begin{pmatrix} A & 0 & B' \\ 0 & D'_1 & 0 \\ C' & 0 & D'_4 \end{pmatrix} \in SO(2, n+1) \mid \begin{pmatrix} A & B' \\ C' & D'_4 \end{pmatrix} \in SO_0(2, q) \right. \right. \\ \left. \left. D'_1 \in SO(p+1) \right\} \right. \\ \simeq SO_0(2, q) \times SO(p+1).$$

Intersecting with H_1 gives us

$$H = \left\{ \left(\begin{pmatrix} a_1 & 0 & 0 & 0 & B'_2 \\ 0 & a_4 & 0 & 0 & 0 \\ 0 & 0 & D_1 & 0 & 0 \\ 0 & 0 & 0 & d_4 & 0 \\ C'_3 & 0 & 0 & 0 & D'_4 \end{pmatrix} \in SO(2, n+1) \mid \begin{pmatrix} a_4, d_4 \in \{\pm 1\} \\ \begin{pmatrix} a_1 & B'_2 \\ C'_2 & D'_4 \end{pmatrix} \in SO_0(1, q) \\ a_1 a_4 \geq 1 \\ d_4 \det D_1 = 1 \end{pmatrix} \right. \right. \\ \left. \left. \simeq SO_0(1, q) \times O(1) \times O(p), \right. \right.$$

and the centralizer $Z_G(X^0)$ is formed by the elements with $a_1 d_4 = 1$, i.e. $Z_G(X^0) \simeq SO_0(1, q) \times O(p)$. Therefore we have a causal compactification for $G/Z_G(X^0) \simeq SO_0(2, q) \times SO(p+1)/(SO_0(1, q) \times O(p))$.

REFERENCES

- [Be] W. Bertram, *Algebraic structures of Makarevič spaces I*, Transformation Groups **3** (1988), 3–32. MR **99c**:32047
- [FÓ] F. Betten and G. Ólafsson, *Causal compactification and Hardy spaces for spaces of Hermitian type*, Pacific J. Math. **200** (2001), 273–312. CMP 2002:05
- [He 78] S. Helgason, *Differential Geometry, Lie Groups and Symmetric Spaces*, Academic Press, New York 1978 MR **80k**:53081
- [HÓ 96] J. Hilgert and G. Ólafsson, *Causal symmetric spaces, The Geometry and Harmonic Analysis*, Perspectives in Math. 18, Academic Press, New York 1996 MR **97m**:43006
- [HÓØ] J. Hilgert, G. Ólafsson, and B. Ørsted, *Hardy spaces on affine symmetric spaces*, J. reine angew. Math. **415** (1991), 189–218 MR **92h**:22030
- [Ka 87] S. Kaneyuki, *On orbit structure of compactification of para-hermitian symmetric spaces*, Japan J. Math. **13** (1987), 333–370 MR **88m**:53094
- [Ka 89] S. Kaneyuki, *On the causal structures of the Šilov boundary of symmetric bounded domains*, in “Prospects in Complex Geometry”, Springer LN 1468, New York 1989, pp. 127–159 MR **92h**:32049
- [Ko 93] K. Koufany, *Semi-groupe de Lie associé à une algèbre de Jordan euclidienne*, thesis, Université de Nancy I, 1993
- [KØ96] K. Koufany and B. Ørsted, *Function spaces on the Ol’shanskii semigroup and the Gelf’and-Gindikin program*, Ann. Inst. Fourier **46** (1996), 689–722 MR **97k**:22021
- [KØ97] K. Koufany and B. Ørsted, *Hardy spaces on two sheeted covering semigroups*, J. Lie Theory **7** (1997), 245–267 MR **98k**:22060
- [KW 65] A. Korányi and J. A. Wolf, *Realization of Hermitian symmetric spaces as generalized half planes*, Ann. of Math. **81** (1965), 265–288 MR **30**:4980
- [Ma 73] B. O. Makarevič, *Open symmetric orbits of reductive groups in symmetric R-spaces*, Math. USSR Sbornik **20** (1973), 406–418 MR **50**:1170
- [Ma 79] T. Matsuki, *The orbits of affine symmetric spaces under the action of minimal parabolic subgroups*, J. Math. Soc. Japan **31** (1979), 331–357 MR **81a**:53049
- [Ma 82] T. Matsuki, *Orbits on affine symmetric spaces under the action of parabolic subgroups*, Hiroshima Math. J. **12** (1982), 307–320 MR **83k**:53072
- [Mo 97] V. F. Molchanov, *Holomorphic discrete series for hyperboloids of Hermitian type*, J. Funct. Anal. **147** (1997), 26–50 MR **99b**:22026
- [ÓI 90] G. Ólafsson, *Causal symmetric spaces*, Mathematica Gottingensis 15, 1990

- [ÓØ 97] G. Ólafsson and B. Ørsted, *Causal compactification and Hardy spaces*, Trans. AMS **351** (1999), 3771–3792 MR **99m**:22005
- [Va 84] V. S. Varadarajan, *Lie Groups, Lie Algebras and Their Representations*, 2nd edition, Springer, New York 1984 MR **85e**:22001
- [Wa 72] G. Warner, *Harmonic Analysis on Semi-Simple Lie Groups I*, Springer Grundlehren der mathematischen Wissenschaften 188, Berlin 1972 MR **58**:16979
- [Wo 72] J. A. Wolf, *Fine Structure of Hermitian Symmetric Spaces*, in “Symmetric Spaces”, pp. 271–357, Dekker, New York 1972 MR **53**:8516

MATHEMATISCHES INSTITUT, UNIVERSITÄT GÖTTINGEN, BUNSENSTRASSE 3–5, D–37073 GÖTTINGEN, GERMANY