

ANALYSING FINITE LOCALLY s -ARC TRANSITIVE GRAPHS

MICHAEL GIUDICI, CAI HENG LI, AND CHERYL E. PRAEGER

ABSTRACT. We present a new approach to analysing finite graphs which admit a vertex intransitive group of automorphisms G and are either locally (G, s) -arc transitive for $s \geq 2$ or G -locally primitive. Such graphs are bipartite with the two parts of the bipartition being the orbits of G . Given a normal subgroup N which is intransitive on both parts of the bipartition, we show that taking quotients with respect to the orbits of N preserves both local primitivity and local s -arc transitivity and leads us to study graphs where G acts faithfully on both orbits and quasiprimively on at least one. We determine the possible quasiprimitive types for G in these two cases and give new constructions of examples for each possible type. The analysis raises several open problems which are discussed in the final section.

1. INTRODUCTION

Let Γ be a connected graph with vertex set $V\Gamma$, edge set $E\Gamma$ and adjacency denoted by \sim . An s -arc in Γ is an $(s + 1)$ -tuple (v_0, v_1, \dots, v_s) of vertices in Γ such that $v_i \sim v_{i-1}$ and $v_{j-1} \not\sim v_{j+1}$ for each $i = 1, \dots, s$ and $j = 1, \dots, s - 1$. Given $G \leq \text{Aut}(\Gamma)$ we call Γ *locally (G, s) -arc transitive* if Γ contains an s -arc and given any two s -arcs α and β starting at the same vertex v , there exists an element $g \in G_v$ mapping α to β . We say Γ is *locally s -arc transitive* if it is locally (G, s) -arc transitive for some $G \leq \text{Aut}(\Gamma)$. If all vertices in Γ have valency at least two, local (G, s) -arc transitivity implies local $(G, s - 1)$ -arc transitivity. We concentrate on the case where $s \geq 2$ as local 1-arc transitivity is equivalent to edge transitivity, a far more general class of graphs.

Locally s -arc transitive graphs are intimately linked to many areas of mathematics. A locally s -arc transitive graph of girth g with at least one vertex of valency greater than two satisfies $s \leq (g + 2)/2$. (The *girth* of Γ is the smallest integer g such that Γ contains a cycle of length g .) When equality holds, various interrelated objects arise such as generalised $(s - 1)$ -gons, groups with a (B, N) -pair of rank two, Moufang polygons and Tutte's m -cages. These connections are explained in more detail in [33]. Locally s -arc transitive graphs also arose in the work of Delgado and Stellmacher in [6] on weak (B, N) -pairs of rank two. Other interesting examples arise naturally from incidence graphs of various structures, for example the points and lines of an affine space (Example 4.2) or the vertices and maximal cliques of a Hamming graph (Example 4.3).

Received by the editors November 22, 2002.

2000 *Mathematics Subject Classification*. Primary 05C25, 20B25.

This paper forms part of an Australian Research Council large grant project which supported the first author. The second author was supported by an ARC Fellowship.

A group G acting on a set Ω is said to be *primitive* if the only partitions of Ω that G preserves are the trivial ones, that is, the whole set or the partition into sets of size one. We call a graph G -*locally primitive* if for each vertex v , the stabiliser G_v acts primitively on $\Gamma(v)$, where $\Gamma(v)$ is the set of vertices adjacent to v . We say Γ is locally primitive if it is G -locally primitive for some automorphism group G . Now suppose that each vertex of a graph Γ has valency at least two. Then Γ is locally $(G, 2)$ -arc transitive if and only if for every vertex v , G_v acts 2-transitively on the set $\Gamma(v)$ (see Lemma 3.2). Thus locally $(G, 2)$ -arc transitive graphs are G -locally primitive.

Let Γ be a locally (G, s) -arc transitive graph. If G is transitive on vertices, then G acts transitively on the set of s -arcs of Γ , that is, Γ is (G, s) -arc transitive. Interest in such graphs goes back to Tutte's seminal work ([29], [30]) showing that (G, s) -arc transitive graphs of valency three satisfy $s \leq 5$. Later, Weiss [32] proved that if the valency is at least three, then $s \leq 7$. Unlike Tutte's ingenious elementary methods, Weiss's proof uses the classification of finite 2-transitive groups and thereby the finite simple groups classification. Note that cycles are s -arc transitive for all s . The second author [19] has determined all (G, s) -arc transitive graphs with $s \geq 4$ for which G acts primitively or biprimitively on vertices. This result and others mentioned below also depend on the finite simple group classification.

If G does not act transitively on vertices and all vertices have valency at least two, then as G is edge transitive, Γ is bipartite and G has two orbits on vertices: the two parts of the bipartition. Weiss (see [31]), has proved that $s \leq 7$ in the valency three case while Stellmacher announced in Siena in 1996 that $s \leq 9$ for locally (G, s) -arc transitive graphs with all vertices of valency at least three (see [27]). The maximum $s = 9$ is attained by the classical generalised octagons. We see in Corollary 3.13 that if Γ has a vertex of valency two and a vertex of valency at least three, then $s \leq 13$. If all vertices have valency two, then the graph is a cycle. Most work on locally (G, s) -arc transitive graphs has centered around the local structure and more specifically the possible point stabilisers of two adjacent vertices. The theory of amalgams plays a crucial role in such a strategy and Goldschmidt's [14] paper determines all possible vertex stabilisers in the valency three case. Stellmacher and van Bon are currently attempting to determine all possible vertex stabilisers for locally $(G, 4)$ -arc transitive graphs where all vertices have valency at least three.

Studying certain quotients of graphs has been very successful in investigating various families of graphs, for example distance transitive graphs [26] and s -arc transitive graphs [23]. We now outline this method for the latter case. Let Γ be a graph with a group of automorphisms G . If G has a normal subgroup N which acts intransitively on vertices, then we define the *quotient graph* Γ_N to have vertex set the N -orbits on $V\Gamma$, and two N -orbits B_1 and B_2 are adjacent in Γ_N if and only if there exist $v \in B_1$ and $w \in B_2$ such that v and w are adjacent in Γ . The original graph Γ is said to be a *cover* of Γ_N if $|\Gamma(v) \cap B_2| = 1$ for each edge $\{B_1, B_2\}$ in Γ_N and $v \in B_1$. If Γ is a nonbipartite (G, s) -arc transitive graph, then Γ is a cover of Γ_N and Γ_N is a $(G/N, s)$ -arc transitive graph. This reduces the problem to finding all examples where G acts quasiprimatively on vertices and then studying their covers. (A transitive permutation group G on a set Ω is said to be *quasiprimitive* if every nontrivial normal subgroup of G acts transitively on Ω .) There are eight types of quasiprimitive groups (these are described in Section 2), and it was shown

in [23] that only four of them, HA, TW, AS and PA, occur for nonbipartite (G, s) -arc transitive graphs. All graphs arising in the HA case were determined in [16], while Baddeley [1] provided a nice characterisation for the graphs of type TW. A similar reduction for the vertex transitive, bipartite case was done in [22].

The purpose of this paper is to initiate a similar program for locally (G, s) -arc transitive graphs for which G acts intransitively on vertices. We see that the appropriate quotients to take are those with respect to the orbits of a certain normal subgroup which is intransitive on both the G -orbits. The O’Nan–Scott Theorem for quasiprimitive groups is then used to study the “basic” graphs, those for which there are no suitable normal quotients. Wherever possible we will prove our results in the more general case of G -vertex intransitive, G -locally primitive graphs. Note that if Γ is a bipartite (G, s) -arc transitive graph, then Γ is locally (G^+, s) -arc transitive for the index two subgroup G^+ of G which fixes both parts of the bipartition. Our results applied to G^+ provide a slightly different analysis for bipartite (G, s) -arc transitive graphs to the one in [22].

Our first result is a reduction theorem, but before stating it we need the following definition. A graph Γ is called a *star* if there exists a vertex v adjacent to every other vertex of Γ and all vertices apart from v have valency one, that is, $\Gamma = K_{1,n}$ for some positive integer n . We call the vertex v the *central vertex* of Γ .

Theorem 1.1. *Let Γ be a finite connected G -locally primitive graph such that G has two orbits Δ_1 and Δ_2 on the set of vertices of Γ with each $|\Delta_i| > 1$. Let N be a normal subgroup of G , the maximal subject to being intransitive on both orbits Δ_1 and Δ_2 . Then Γ_N is a (G/N) -locally primitive graph, Γ is a cover of Γ_N and Γ_N satisfies at least one of the following:*

- (1) $\Gamma_N \cong K_{n,n'}$.
- (2) G/N acts faithfully and quasiprimively on both of its orbits on $V\Gamma_N$.
- (3) G/N acts faithfully on both of its orbits on $V\Gamma_N$ but only acts quasiprimively on one of them. Furthermore, there exists $M \triangleleft G$ containing N , transitive on one G -orbit but intransitive on the other, and Γ_M is a star.

Moreover, if Γ is locally (G, s) -arc transitive, then Γ_N is locally $(G/N, s)$ -arc transitive. In fact, every star occurs as a normal quotient of some locally 3-arc transitive graph.

This suggests that to investigate locally (G, s) -arc transitive graphs we should concentrate on three problems: determining such graphs where G acts quasiprimively on both of its orbits, determining such graphs where G acts quasiprimively on only one of its orbits, and investigating locally s -arc transitive covers of these two types of graphs and of complete bipartite graphs. A similar result for the case where Γ is locally s -arc transitive for $s \geq 4$ and valency at least three was obtained in [17], but the situation is different there, as the stars and complete bipartite graphs do not occur. See also Lemma 5.6.

Now $\text{Aut}(\Gamma)$ may or may not act transitively on the vertices of Γ . Hence Γ is not necessarily regular and G may act differently on each of its orbits. We say that G is of *type* $\{X, Y\}$ if G acts quasiprimively of type X on one G -orbit and quasiprimively of type Y on the other G -orbit, where the types X and Y refer to quasiprimitive types as described in Section 2. Often when $X = Y$, we will relax notation and say that G is of type X . The following theorem determines the

possibilities for the type of G and shows that the two actions of G are usually of the same type.

Theorem 1.2. *Let Γ be a finite G -locally primitive connected graph such that G has two orbits on vertices and G acts faithfully and quasiprimively on both orbits with type $\{X, Y\}$. Then either $X = Y$, or $\{X, Y\} = \{\text{SD}, \text{PA}\}$ or $\{\text{CD}, \text{PA}\}$, and examples exist in each case. Furthermore, if Γ is locally (G, s) -arc transitive with $s \geq 2$, then either $X = Y \in \{\text{HA}, \text{TW}, \text{AS}, \text{PA}\}$ or $\{X, Y\} = \{\text{SD}, \text{PA}\}$, and again examples exist in each case.*

Examples of locally (G, s) -arc transitive graphs where G is quasiprimitive of type HA, TW, AS and PA will be provided in Lemma 3.3 while Example 4.1 gives examples where G is of type $\{\text{SD}, \text{PA}\}$. Examples of G -locally primitive graphs of the remaining possible types can be found in Example 6.6. Note that Theorem 1.2 corresponds closely to the result in [23] that for a (G, s) -arc transitive, nonbipartite graph where G acts on vertices quasiprimively, G has quasiprimitive type HA, TW, AS or PA [23]. If G is of quasiprimitive type HA on both of its orbits, Γ was shown to be vertex transitive in [15, Lemma 3.2] and this situation is discussed in [16]. When G acts quasiprimively on only one orbit we have the following theorem.

Theorem 1.3. *Let Γ be a finite locally (G, s) -arc transitive connected graph with $s \geq 2$ such that G acts faithfully on both of its orbits Δ_1 and Δ_2 but only acts quasiprimively on Δ_1 . Then the quasiprimitive action of G on Δ_1 is of type HA, HS, AS, PA or TW, and examples exist for each of these types.*

We provide examples for each of the five possibilities in Section 4. For the more general class of G -locally primitive graphs, the quasiprimitive action on Δ_1 may be of the three remaining types as seen in Example 7.2.

In Section 3, we give some general constructions of edge transitive graphs and in particular we characterise locally s -arc transitive graphs having a vertex of valency two and show that for such graphs $s \leq 13$ (see Theorem 3.10 and Corollary 3.13). We provide examples in Section 4 verifying our claims about the existence of locally (G, s) -arc transitive graphs with G quasiprimitive of various types. Section 5 is devoted to the proof of Theorem 1.1, Section 6 to the proof of Theorem 1.2 and Section 7 for Theorem 1.3. Finally in Section 8 we discuss some open problems and future directions arising from this work.

2. QUASIPRIMITIVE GROUPS

Recall that a transitive permutation group G on a set Ω is called quasiprimitive if every nontrivial normal subgroup of G acts transitively on Ω . This is a generalisation of primitivity as every normal subgroup of a primitive group is transitive, but there exist quasiprimitive groups which are not primitive. The third author [23] generalised the O’Nan–Scott Theorem for primitive groups to quasiprimitive groups and showed that a finite quasiprimitive group is one of eight types. We now provide a description of these types along the lines of that in [24]. Throughout, T will be a finite nonabelian simple group.

The first three types of quasiprimitive groups are subgroups of the *holomorph* $\text{Hol}(N)$ of a certain group N . This is defined as the semidirect product $\text{Hol}(N) = N \cdot \text{Aut}(N)$ acting on $\Omega = N$ where, for all $n \in N$ and $\sigma \in \text{Aut}(N)$,

$$n\sigma : x \mapsto x^\sigma n^\sigma \text{ for all } x \in \Omega.$$

Thus the group N is normal in $\text{Hol}(N)$ and acts regularly by right multiplication.

The first quasiprimitive type is HA, and all groups of this type are in fact primitive. This class consists of all quasiprimitive groups with an elementary abelian minimal normal subgroup. Such groups are subgroups of $\text{AGL}(d, p)$, for some prime p and positive integer d , acting on the points of the affine space $\text{AG}(d, p)$. The group $N \cong C_p^d$ of all translations is the unique minimal normal subgroup of G and $G = N : G_0$ for some irreducible subgroup $G_0 \leq \text{GL}(d, p)$. Note that $\text{AGL}(d, p) = \text{Hol}(N)$.

The next type of quasiprimitive group is HS. Groups of this type are subgroups of the holomorph $\text{Hol}(T) = T \cdot \text{Aut}(T)$ containing $T \cdot \text{Inn}(T)$ and acting on $\Omega = T$ as above. Such groups have two minimal normal subgroups, both of which are isomorphic to T and act regularly on Ω . One minimal normal subgroup is T in its action on itself by right multiplication and the other is T acting on itself by left multiplication. Thus the socle, $\text{soc}(G)$, of G is $T \times T = T \cdot \text{Inn}(T)$ and acts on Ω by

$$(t_1, t_2) : s \mapsto t_2^{-1} s t_1 \text{ for all } s \in T.$$

All groups of this type are primitive.

The third type of quasiprimitive group HC also consists only of primitive groups. These groups are similar to those of type HS, but here $G \leq \text{Hol}(T^k)$ for $k \geq 2$. The action on $\Omega = T^k$ is again the holomorph action and G has two minimal normal subgroups, both regular and isomorphic to T^k . Now $\text{Inn}(T^k) \leq G_1 \leq \text{Aut}(T^k)$ and we require that G_1 acts transitively on the k simple direct factors of T^k . The groups of types HS and HC are the only quasiprimitive groups with two minimal normal subgroups. All the rest have only one.

We now define a twisted wreath product by following the description in [28, p. 269]. Let T be a finite nonabelian simple group, P a group with a proper subgroup Q and let $\phi : Q \rightarrow T$ be a homomorphism. We define the *complete base group* \mathcal{B} to be the set of maps $f : P \rightarrow T$ under pointwise multiplication and so $\mathcal{B} \cong T^{|P|}$. The group P acts on \mathcal{B} by

$$f^p(x) = f(px) \text{ for all } x, p \in P, f \in \mathcal{B}.$$

We define \mathcal{X} to be the semidirect product $\mathcal{B} : P$ with respect to this action. Define the ϕ -base group

$$B_\phi = \{f : P \rightarrow T \mid f(pq) = f(p)^{\phi(q)} \text{ for all } p \in P, q \in Q\}.$$

This group is isomorphic to T^k , where $k = |P : Q|$. Also B_ϕ is normalised by P , and so B_ϕ and P generate the subgroup $X_\phi = B_\phi : P$ of \mathcal{X} which we call the *twisted wreath product* $T \text{ twr}_\phi P$ of T by P with respect to ϕ . The action of X_ϕ on its base group is quasiprimitive if and only if $\phi^{-1}(\text{Inn}(T))$ is a core free subgroup of P (see [23]). Such groups are said to be of quasiprimitive type TW and are the only quasiprimitive groups with a unique minimal normal subgroup isomorphic to T^k for $k \geq 2$ which acts regularly. The action is primitive if and only if ϕ does not extend to a larger subgroup of P and $\text{Inn}(T) \leq \phi(Q)$ (see [21]).

The next type of quasiprimitive group is AS, and these are all the almost simple groups with transitive socle T , that is $T \leq G \leq \text{Aut}(T)$. We are not given any further information about the action in this case, and in particular it is possible for T to act regularly on Ω . The group G is primitive if and only if a point stabiliser is a maximal subgroup of G not containing T .

Next we describe quasiprimitive groups of type SD. Let $N = T^k$ for $k \geq 2$. A *full diagonal subgroup* of N is a subgroup isomorphic to T whose projection onto every coordinate is also isomorphic to T . We let N act on the set Ω of right cosets of the full diagonal subgroup $N_\alpha = \{(t, \dots, t) : t \in T\}$. Then the set $\{[t_1, \dots, t_{k-1}, 1] : t_i \in T\}$ forms a set of coset representatives for N_α in N and so we may identify Ω with T^{k-1} . The action of N is then given by

$$(s_1, \dots, s_k) : [t_1, \dots, t_{k-1}, 1] \mapsto [s_k^{-1}t_1s_1, \dots, s_k^{-1}t_{k-1}s_{k-1}, 1].$$

Let W be the normaliser of N in $\text{Sym}(\Omega)$. Then for each $h \in \text{Aut}(T)$, the permutation

$$[t_1, \dots, t_{k-1}, 1] \mapsto [t_1^h, \dots, t_{k-1}^h, 1]$$

induced by h is in W . Note that the permutations induced by $\text{Inn}(T)$ are precisely those induced by N_α . Also, for each $\sigma \in S_k$, if we let $t_k = 1$, the permutation

$$[t_1, \dots, t_{k-1}, 1] \mapsto [t_{1\sigma^{-1}}, \dots, t_{k\sigma^{-1}}] = [t_{k\sigma^{-1}}^{-1}t_{1\sigma^{-1}}, \dots, t_{k\sigma^{-1}}^{-1}t_{(k-1)\sigma^{-1}}, 1]$$

is in W . In fact, $W = \langle N, \text{Aut}(T), S_k \rangle \cong T^k \cdot (\text{Out}(T) \times S_k)$, where the extension of T^k by $\text{Out}(T) \times S_k$ is not necessarily a split extension. A quasiprimitive group of type SD is a group G such that $N \triangleleft G \leq W$ and G acts transitively by conjugation on the set of simple direct factors of N , that is, N is the unique minimal normal subgroup of G . Such groups are primitive if and only if G acts primitively on the k simple direct factors of N . We note that, in the $k = 2$ case, if $N \triangleleft G < W$ and G does not act transitively by conjugation on the two simple direct factors of N , then N is the product of two minimal normal subgroups of G and G is primitive of type HS.

Quasiprimitive groups of type CD are built from quasiprimitive groups of type SD. Let H be a quasiprimitive group of type SD on the set Δ with a unique minimal normal subgroup T^l . For a positive integer k divisible by l , if G satisfies $N = T^k \leq G \leq H$ wr $S_{k/l}$, then G acts on the set $\Omega = \Delta^{k/l}$ with the product action of the wreath product, namely, for $(\delta_1, \dots, \delta_{k/l}) \in \Omega$, $h = (h_1, \dots, h_{k/l}) \in H^{k/l}$ and $\sigma \in S_{k/l}$,

$$(\delta_1, \dots, \delta_{k/l})^h = (\delta_1^{h_1}, \dots, \delta_{k/l}^{h_{k/l}})$$

and

$$(\delta_1, \dots, \delta_{k/l})^\sigma = (\delta_{1\sigma^{-1}}, \dots, \delta_{(k/l)\sigma^{-1}}).$$

This action is quasiprimitive if and only if G acts transitively by conjugation on the set of simple direct factors of N . Then N is the unique minimal normal subgroup of G . The action is primitive if H is primitive. We note here that groups of type HC can be built up in the same way from groups of type HS.

The last type of quasiprimitive group is PA. A group G of this type preserves some partition \mathcal{P} (possibly trivial in the sense of having parts of size 1) of Ω upon which G acts faithfully preserving a product structure Δ^k . Furthermore, $N = T^k \leq G \leq H$ wr S_k , where H acts quasiprimitively on Δ of type AS with nonregular socle T and G acts transitively by conjugation on the set of simple direct factors of N . Let $\delta \in \Delta$ and $B = (\delta, \dots, \delta) \in \mathcal{P}$. Then $N_B = T_\delta^k$ and for $\omega \in B$, the point stabiliser N_ω is a subdirect subgroup of N_B , that is, N_ω projects onto T_δ in every coordinate. The action of G is primitive if and only if \mathcal{P} is trivial and the action of H on Δ is primitive.

3. GENERAL GRAPH CONSTRUCTIONS

In this section we provide three methods for constructing G -edge transitive graphs. The constructions in Sections 3.1 and 3.3 enable us to build a locally (G, s) -arc transitive graph from a given (G, s) -arc transitive graph. Section 3.2 shows how to construct arbitrary locally (G, s) -arc transitive, G -vertex intransitive graphs as coset graphs. The construction in Section 3.3 characterises locally (G, s) -arc transitive graphs with a vertex of valency two.

We use the following conventions for graphs. A *digraph* Γ consists of a vertex set $V\Gamma$ and a subset $A\Gamma$ of $(V\Gamma \times V\Gamma) \setminus \{(v, v) : v \in V\Gamma\}$ called *arcs*. If $A\Gamma$ is self-paired in the sense that $(x, y) \in A\Gamma$ if and only if $(y, x) \in A\Gamma$, then we say that Γ is *self-paired*. In this case we usually replace every pair of arcs between two vertices with a single undirected edge to form an undirected graph. The edge set is denoted $E\Gamma$. If $G \leq \text{Aut}(\Gamma)$ is edge transitive on Γ with two vertex orbits Δ_1 and Δ_2 , then all vertices in the same orbit Δ_i have the same valency k_i say, and we say that Γ is *biregular* of valency $\{k_1, k_2\}$. If $k_1 = k_2 = k$, then Γ is regular of valency k .

Let G be a transitive permutation group on the set V and let \mathcal{O} be a nontrivial orbital for G , that is, an orbit of G on $(V \times V) \setminus \{(v, v) : v \in V\}$. We define the *orbital digraph* Γ for G with respect to \mathcal{O} to be the directed graph with vertex set V and arc set \mathcal{O} . This graph is G -vertex transitive and G -arc transitive. Conversely, every G -vertex transitive and G -arc transitive graph can be constructed in this way. If \mathcal{O} is self-paired, then the corresponding undirected graph is called an *orbital graph*.

We note the following elementary lemmas. The proof of the first can be found, for example, in [2].

Lemma 3.1. *Let Γ be a connected locally (G, s) -arc transitive graph such that $s \geq 1$ and all vertices have valency at least two. Then G acts transitively on the set of edges of Γ . Furthermore, if G acts intransitively on $V\Gamma$, then Γ is a bipartite graph and the two parts of the bipartition are G -orbits.*

Lemma 3.2. *Let Γ be a graph such that all vertices have valency at least two. Then Γ is locally $(G, 2)$ -arc transitive if and only if for every vertex v , G_v acts 2-transitively on $\Gamma(v)$.*

Proof. Let Γ be locally $(G, 2)$ -arc transitive with all vertices having valency at least two and let $v \in V\Gamma$. By Lemma 3.1, G acts transitively on the set of edges of Γ and so G_v acts transitively on $\Gamma(v)$. Let $u \in \Gamma(v)$. Then for every $w \in \Gamma(v) \setminus \{u\}$ there is a 2-arc (u, v, w) . As Γ is locally $(G, 2)$ -arc transitive, it follows that G_{uv} acts transitively on $\Gamma(v) \setminus \{u\}$ and so G_v acts 2-transitively on $\Gamma(v)$.

Conversely, suppose that for every vertex v , G_v acts 2-transitively on $\Gamma(v)$. Let (v, u_1, w_1) and (v, u_2, w_2) be two 2-arcs in Γ . Then as G_v acts transitively on $\Gamma(v)$, there exists $g \in G_v$ such that $(v, u_1, w_1)^g = (v, u_2, w'_1)$ for some $w'_1 \in \Gamma(u_2) \setminus \{v\}$. Also, as G_{u_2} acts 2-transitively on $\Gamma(u_2)$, there exists $h \in G_{u_2 v}$ such that $(w'_1)^h = w_2$. Hence $(v, u_1, w_1)^{gh} = (v, u_2, w_2)$ and so Γ is locally $(G, 2)$ -arc transitive. \square

3.1. Double covers. Let Γ be a directed or undirected vertex transitive graph with vertex set $V\Gamma$ and arc set $A\Gamma$. The *standard double cover* of Γ is the undirected graph $\bar{\Gamma}$ with a vertex set $V\Gamma \times \{1, 2\}$, and two vertices $(x, 1)$ and $(y, 2)$ are adjacent if and only if $(x, y) \in A\Gamma$. The new graph is bipartite with bipartite halves $V\Gamma \times \{i\}$ for each $i = 1, 2$.

Now if $G \leq \text{Aut}(\Gamma)$, then G also acts as a group of automorphisms of $\overline{\Gamma}$ with the action $g : (x, i) \mapsto (x^g, i)$. If G is vertex transitive on Γ , then G has two orbits on the set of vertices of $\overline{\Gamma}$ and the action of G on each orbit is permutationally isomorphic to the action of G on $V\Gamma$. Furthermore, $G_v = G_{(v,i)}$ for each $i = 1, 2$. Then if Γ is undirected, the action of G_v on $\Gamma(v)$ is the same as the action of $G_{(v,i)}$ on $\overline{\Gamma}((v, i))$. Thus in this case, if Γ is G -locally primitive, $\overline{\Gamma}$ is also G -locally primitive.

Suppose again that Γ is undirected. Then $(x, 1) \sim (y, 2)$ if and only if $(y, 1) \sim (x, 2)$. If Γ is also connected, then for each $x, y \in V\Gamma$ there exists a path P in Γ between x and y . This path lifts to a path in $\overline{\Gamma}$ between $(x, 1)$ and $(y, 1)$ if P has even length, and to one between $(x, 1)$ and $(y, 2)$ if P has odd length. There is a path between $(y, 1)$ and $(y, 2)$ if and only if y is in an odd cycle in Γ . Thus for an undirected connected graph Γ , $\overline{\Gamma}$ is connected if and only if Γ contains an odd cycle, that is, if and only if Γ is not bipartite. If $((v, i), (v_1, j), \dots, (v_s, i))$ is an s -arc in $\overline{\Gamma}$, then (v, v_1, \dots, v_s) is an s -arc in Γ . Thus if Γ is (G, s) -arc transitive, then $\overline{\Gamma}$ is locally (G, s) -arc transitive. Since there are nonbipartite, $(G, 2)$ -arc transitive graphs with G quasiprimitive of each of the types HA, TW, AS and PA (see [24]), we can construct locally $(G, 2)$ -arc transitive graphs with G of types HA, TW, AS and PA on both of its orbits by taking their standard double covers. Note that these graphs are $\text{Aut}(\Gamma)$ -vertex transitive, as the map $\tau : (x, i) \mapsto (x, 3 - i)$ is an automorphism interchanging the two halves of the bipartition. We summarise this discussion in the following lemma.

Lemma 3.3. *Let Γ be an undirected graph. Then*

- (1) $\overline{\Gamma}$ is connected if and only if Γ is connected and not bipartite.
- (2) If Γ is G -locally primitive, then $\overline{\Gamma}$ is G -locally primitive.
- (3) If Γ is (G, s) -arc transitive, then $\overline{\Gamma}$ is locally (G, s) -arc transitive. In particular, there exist quasiprimitive locally $(G, 2)$ -arc transitive graphs of types HA, TW, AS and PA.

Note that when Γ is directed, paths and s -arcs in Γ no longer lift to paths and s -arcs in $\overline{\Gamma}$.

Now let Γ be a G -edge transitive, bipartite graph such that each vertex has valency at least two, and suppose that G has two orbits Δ_1 and Δ_2 on vertices and all vertex stabilisers are conjugate. Then the two actions of G are permutationally isomorphic and so there exists a bijection $\phi : \Delta_1 \rightarrow \Delta_2$ such that $\phi(v^g) = \phi(v)^g$ for all $v \in \Delta_1$ and $g \in G$. Note that v and $\phi(v)$ have the same vertex stabilisers. By identifying the vertices v and $\phi(v)$, we can label the vertex set of Γ with $\Delta_1 \times \{1, 2\}$. Define the directed graph Σ with vertex set Δ_1 and arc set such that (v, w) is an arc if and only if $(v, 1) \sim (w, 2)$ in Γ . Note that $(v, 1)$ is not adjacent to $(v, 2)$ in Γ as $G_{(v,1)}$ acts transitively on edges emerging from $(v, 1)$. Then Σ is a G -vertex transitive and G -arc transitive directed graph and so is an orbital digraph for G on Δ_1 . Furthermore, Γ is the standard double cover of the directed graph Σ . The bijection $\varphi : V\Gamma \rightarrow V\Gamma$ which interchanges $(v, 1)$ and $(v, 2)$ for all $v \in \Delta_1$ is an automorphism for Γ if and only if for every arc (v, w) in Σ , (w, v) is also an arc. Thus we have the following.

Lemma 3.4. *Let Γ be a G -edge transitive graph of valency at least two with two vertex orbits Δ_1 and Δ_2 such that all vertex stabilisers are conjugate. Then Γ is the standard double cover of an orbital digraph Σ for G on Δ_1 , and the bijection*

which interchanges $(v, 1)$ with $(v, 2)$ is an automorphism for Γ if and only if Σ is self paired.

Another condition for such a graph Γ to be $\text{Aut}(\Gamma)$ -vertex transitive is given in [10, Lemma 2.5]. There are examples in [10] of locally (G, s) -arc transitive graphs for which all vertex stabilisers are conjugate in G , yet $\text{Aut}(\Gamma)$ is not vertex transitive.

3.2. Coset graphs. Let Γ be a G -edge transitive graph with vertex set $V\Gamma$, where $G \leq \text{Aut}(\Gamma)$. Assume that G is not transitive on $V\Gamma$. Then by Lemma 3.1, Γ is a bipartite graph with parts Δ_1 and Δ_2 such that G is transitive on both Δ_1 and Δ_2 . Since G is transitive on Δ_1 , for $w \in \Delta_1$ we may write Δ_1 as the set $[G : G_w]$ of right cosets of G_w in G so that G acts transitively on Δ_1 by right multiplication, that is,

$$\begin{aligned} \Delta_1 &= \{G_v x : x \in G\}, \text{ and} \\ z &: G_v x \rightarrow G_v xz, \text{ for each } z \in G. \end{aligned}$$

Similarly, for $w \in \Delta_2$, we may write Δ_2 as the set $[G : G_w]$ of right cosets of G_w in G so that G acts transitively on Δ_2 by right multiplication, that is,

$$\begin{aligned} \Delta_2 &= \{G_w x : x \in G\} \text{ and} \\ z &: G_w x \rightarrow G_w xz, \text{ for each } z \in G. \end{aligned}$$

Thus the vertices of Γ may be identified with right cosets of G_v and G_w in G . Now assume that v is adjacent to w . Then $G_v \cap G_w$ is the stabiliser of the edge $\{v, w\}$ in G . It follows since $G \leq \text{Aut}(\Gamma)$ that $G_v \cap G_w$ is core free in G . Since G_v is transitive on $\Gamma(v)$ (by right multiplication), the neighbours of v are exactly the images of w under G_v , that is $\{G_w z : z \in G_v\}$. Similarly, the neighbours of w may be identified with $\{G_v z : z \in G_w\}$. The adjacency relation of Γ is given by

$$G_v x \sim G_w y \iff xy^{-1} \in G_v G_w, \text{ or equivalently } yx^{-1} \in G_w G_v.$$

Hence the vertex set and the adjacency relation of the graph Γ can be explicitly described in terms of cosets of the point stabilisers. Thus we have the following result.

Lemma 3.5. *Let Γ be a bipartite G -edge transitive graph with parts Δ_1 and Δ_2 , where $G \leq \text{Aut}(\Gamma)$ and G is intransitive on $V\Gamma$. Let $v \in \Delta_1$ and $w \in \Delta_2$ be adjacent. Then we may identify $\Delta_1 = [G : G_v]$ and $\Delta_2 = [G : G_w]$ such that*

- (1) $G_v x$ is adjacent to $G_w y$ if and only if $xy^{-1} \in G_v G_w$, or equivalently, $yx^{-1} \in G_w G_v$;
- (2) $\Gamma(v) = \{G_w z : z \in G_v\} = G_w G_v$ and $\Gamma(w) = \{G_v z : z \in G_w\} = G_v G_w$;
- (3) the valencies are $|\Gamma(v)| = |G_v : G_v \cap G_w|$ and $|\Gamma(w)| = |G_w : G_v \cap G_w|$.

Conversely, we may construct edge transitive graphs from any given abstract group.

Definition 3.6. Let G be a group, and let $L, R < G$ be such that $L \cap R$ is core free in G . Let $\Delta_1 = \{Lx : x \in G\}$ and $\Delta_2 = \{Ry : y \in G\}$. Define the bipartite graph

$$\Gamma = \text{Cos}(G, L, R)$$

to be the graph with vertex set $V = \Delta_1 \dot{\cup} \Delta_2$ such that

$$Lx \sim Ry \iff xy^{-1} \in LR, \text{ or equivalently } yx^{-1} \in RL.$$

We refer to $(L, R, L \cap R)$ as the associated *amalgam*.

We note that $xy^{-1} \in LR$ if and only if $Lx \cap Ry \neq \emptyset$, so the above definition of coset graphs coincides with Tits's construction of flag transitive incidence structures; see [7]. It is also related to Du and Xu's [9] construction of bi-coset graphs. The next simple lemma gives some properties of the graphs $Cos(G, L, R)$.

Lemma 3.7. *For a group G and subgroups $L, R < G$ such that $L \cap R$ is core free in G , the graph $\Gamma = Cos(G, L, R)$ satisfies the following properties:*

- (1) Γ is connected if and only if $\langle L, R \rangle = G$;
- (2) $G \leq \text{Aut}(\Gamma)$, and Γ is G -edge transitive and G -vertex intransitive;
- (3) G acts faithfully on both Δ_1 and Δ_2 if and only if both L and R are core free;
- (4) Γ is G -locally primitive if and only if $L \cap R$ is a maximal subgroup of both L and R ;
- (5) Γ is locally $(G, 2)$ -arc transitive if and only if L acts 2-transitively on $[L : L \cap R]$ and R acts 2-transitively on $[R : L \cap R]$.

Conversely, if Γ is a G -edge transitive but not G -vertex transitive graph, and v and w are adjacent vertices, then $\Gamma \cong Cos(G, G_v, G_w)$.

Proof. (1) Assume that $G = \langle L, R \rangle$. Then for any $w \in G$, $w = x_1y_1 \dots x_ky_k$ for some $x_i \in L$ and some $y_i \in R$ such that $x_i \neq 1$ if $i \neq 1$, and $y_i \neq 1$ if $i \neq k$. There is a path in Γ from L to Lw :

$$L, Ry_k, Lx_ky_k, \dots, Ry_1 \dots x_ky_k, Lx_1y_1 \dots x_ky_k = Lw.$$

There is also a path in Γ from L to Rw :

$$L, R, Ly_k, Rx_ky_k, \dots, Ly_1 \dots x_ky_k, Rx_1y_1 \dots x_ky_k = Rw.$$

Thus Γ is connected. Conversely, if Γ is connected, it is easily shown that $G = \langle L, R \rangle$.

(2) Let $\Delta_1 = \{Lx : x \in G\}$ and $\Delta_2 = \{Ry : y \in G\}$. For $z \in G$, let \hat{z} be the permutation on $V = \Delta_1 \cup \Delta_2$ induced by z defined as

$$\hat{z}: Lx \rightarrow Lxz, Ry \rightarrow Ryz.$$

Then

$$\begin{aligned} Lx \sim Ry &\iff xy^{-1} \in LR \\ &\iff xz(yz)^{-1} = xzz^{-1}y^{-1} \in LR \\ &\iff Lxz \sim Ryz \\ &\iff (Lx)^{\hat{z}} \sim (Ry)^{\hat{z}}. \end{aligned}$$

Thus \hat{z} is an automorphism of Γ , and so there is a homomorphism $\phi : G \rightarrow \text{Aut}(\Gamma)$ given by $\phi(z) = \hat{z}$. The kernel of this homomorphism is trivial since $L \cap R$ is core free and hence $G \leq \text{Aut}(\Gamma)$. Clearly, no element of G maps $L \in \Delta_1$ to $R \in \Delta_2$, and hence G is intransitive on V .

Write v and w to be the vertices corresponding to L and R , respectively. By definition, the set of neighbours of v equals $\{Rx : x \in L\}$. For any $z \in L$, $L^{\hat{z}} = Lz = L$ and $R^{\hat{z}} = Rz \in \{Ry : y \in L\} = \Gamma(v)$. When z runs through L , Rz runs through $\{Ry : y \in L\}$. Therefore, L is transitive on $\Gamma(v)$. Similarly, R is transitive on $\Gamma(w) = \{Lx : x \in G\}$ and so Γ is G -edge transitive.

- (3) This follows as $G_v = L$ and $G_w = R$.
- (4) This follows as $G_{vw} = G_w \cap G_v = L \cap R$.
- (5) This follows from Lemma 3.2.

Finally, let Γ be any G -edge transitive graph that is not G -vertex transitive and let v and w be adjacent vertices. The fact that $\Gamma \cong \text{Cos}(G, G_v, G_w)$ follows from Lemma 3.5 and Definition 3.6. \square

Given a group G with subgroup L and element g we can also form the coset digraph $\Sigma = \text{Cos}(G, L, LgL)$ which has vertex set $[G : L]$ and $(Lx, Ly) \in A\Sigma$ if and only if $yx^{-1} \in LgL$. This directed graph is connected if and only if $\langle L, g \rangle = G$ and is self paired if and only if $g^{-1} \in LgL$. Furthermore, G acts as an arc transitive group of automorphisms by right multiplication on the cosets. This construction is due to Sabidussi [25] and is related to our coset construction in the following way.

Lemma 3.8. *Let G be a group with subgroup L and element g such that $L \cap L^g$ is core free. Then $\Gamma = \text{Cos}(G, L, L^g)$ is the standard double cover of*

$$\Sigma = \text{Cos}(G, L, LgL).$$

Proof. Let $\bar{\Sigma}$ be the standard double cover of Σ and define the map

$$\begin{aligned} \phi : \quad V\bar{\Sigma} &\rightarrow V\Gamma, \\ (Lx, 1) &\mapsto Lx, \\ (Lx, 2) &\mapsto L^g g^{-1}x. \end{aligned}$$

Let $g_1, g_2 \in G$. Then

$$\begin{aligned} L^g g^{-1}g_1 = L^g g^{-1}g_2 &\iff g^{-1}g_1g_2^{-1}g \in L^g \\ &\iff g_1g_2^{-1} \in L \\ &\iff Lg_1 = Lg_2. \end{aligned}$$

Furthermore, if \mathcal{T} is a transversal for L in G , then $g^{-1}\mathcal{T}$ is a transversal for L^g in G . Thus ϕ is a well-defined bijection. Suppose that $(Lx, 1) \sim (Ly, 2)$ in $\bar{\Sigma}$. Then (Lx, Ly) is an arc in Σ and so $yx^{-1} \in LgL$. Hence $xy^{-1} \in Lg^{-1}L$ and so $xy^{-1}g \in Lg^{-1}Lg = LL^g$. Thus $\phi((Lx, 1)) = Lx$ is adjacent to $\phi((Ly, 2)) = L^g g^{-1}y$ in Γ . Conversely, suppose that $(Lx, L^g y) \in E\Gamma$. Then $xy^{-1} \in LL^g$ and so $yx^{-1} \in L^g L$. Now $\phi((Lx, 1)) = Lx$ and $\phi((L^g y, 2)) = L^g y$. Furthermore, $gyx^{-1} \in gL^g L = LgL$ and so $(Lx, 1) \sim (L^g y, 2)$ in $\bar{\Sigma}$. Thus ϕ is a graph isomorphism. \square

A version of Lemma 3.8 was proved in [10, Lemma 2.6] in the case where $g^2 \in L$.

3.3. Graphs with a vertex of valency one or two. If Γ is a locally (G, s) -arc transitive graph with all vertices of valency at least two, then Γ is $(G, s - 1)$ -arc transitive. Furthermore, if G acts intransitively on vertices, then Γ is bipartite with the two parts of the bipartition being G -orbits. However, if Γ has a vertex of valency one, then we cannot make these two deductions. We have the following lemma.

Lemma 3.9. *Let Γ be a connected locally (G, s) -arc transitive graph with $s \geq 1$ which contains a vertex of valency one. Then Γ is a tree.*

Proof. Let v be a vertex of valency one in Γ and suppose that Γ contains a cycle. Then as Γ is connected, there exists a shortest path $\{w_0 = v, w_1, \dots, w_t\}$ such that w_t is contained in a cycle C . Since v has valency one, we have $t \geq 1$ and none of the vertices w_0, w_1, \dots, w_{t-1} lies on a cycle. There is an s -arc that starts with the arc (w_{t-1}, w_t) and then proceeds to loop around C finishing at a vertex $u \in C$. Let β be the reverse of this s -arc running from u to w_{t-1} . Let α be the s -arc that agrees with β in its first s vertices but ends in a vertex of C adjacent to w_t . As

$w_{t-1} \in \beta$ and is not contained in a cycle while all vertices in α belong to C , there is no element of G_u mapping α to β . This contradicts Γ being locally (G, s) -arc transitive and so Γ is a tree. \square

Let Γ be a (G, s) -arc transitive graph with valency $k \geq 2$. We follow [2] and form a new graph Γ^* by placing a vertex at the midpoint of each edge of Γ . Then $G \leq \text{Aut}(\Gamma^*)$ and has two orbits on vertices: the set Δ_1 of vertices of Γ and the set Δ_2 of midpoints of edges of Γ . All vertices in Δ_1 have valency k while those in Δ_2 have valency two. If Γ is a cycle, then the new graph Γ^* is also a cycle. We will show in Theorem 3.10 that Γ^* is locally $(G, 2s - 1)$ -arc transitive.

In fact every locally $(G, 2s - 1)$ -arc transitive graph with a vertex of valency two can be constructed in this way. First we need a definition. Given a connected graph Γ , the *distance two graph* of Γ is the graph with vertex set $V\Gamma$ such that two vertices are adjacent if and only if they are at distance two in Γ . Note that if Γ is bipartite, then the distance two graph of Γ has two connected components. Also Γ is the connected component of the distance two graph of Γ^* with vertex set Δ_1 .

Theorem 3.10. *Let $s \geq 1$.*

- (a) *Let Γ be a connected locally $(G, 2s - 1)$ -arc transitive graph of valency $\{2, k\}$ with $k \geq 3$ and let $\tilde{\Gamma}$ be a connected component of the distance two graph of Γ containing a vertex of valency k . Then $V\tilde{\Gamma}$ is the set of all vertices of Γ of valency k and $\tilde{\Gamma}$ is (G, s) -arc transitive of valency k .*
- (b) *Let Σ be a connected (G, s) -arc transitive graph of valency k . Then Σ^* is a connected locally $(G, 2s - 1)$ -arc transitive graph. Moreover, if $\Gamma = \Sigma^*$, then the graph $\tilde{\Gamma}$ in part (a) is equal to Σ .*

Proof. (a) Let Γ be a connected locally $(G, 2s - 1)$ -arc transitive graph such that G has two orbits Δ_1 and Δ_2 on vertices. Suppose that all vertices in Δ_2 have valency two while the vertices in Δ_1 have valency $k \geq 3$. Let $\tilde{\Gamma}$ be a connected component of the distance two graph of Γ containing a vertex of valency k . Then as Γ is connected, $V\tilde{\Gamma} = \Delta_1$. Let $v \in \Delta_1$ and $w \in \Gamma(v)$. Then there exists a unique element $v(w) \in \Gamma(w) \setminus \{v\}$ and so the map $w \mapsto v(w)$ is a 1-1 correspondence between $\Gamma(v)$ and $\Gamma_2(v)$. Thus $\tilde{\Gamma}$ has valency k . Now $G \leq \text{Aut}(\tilde{\Gamma})$ and acts transitively on vertices. Let (v_0, v_1, \dots, v_s) be an s -arc in $\tilde{\Gamma}$. Then for each $i = 0, \dots, s - 1$ there exists a unique w_i such that $\Gamma(w_i) = \{v_i, v_{i+1}\}$ and so $(v_0, w_0, v_1, \dots, v_{s-1}, w_{s-1})$ is a $(2s - 1)$ -arc in Γ . Since Γ is locally $(G, 2s - 1)$ -arc transitive, it follows that G_{v_0} is transitive on the set of all s -arcs in $\tilde{\Gamma}$ beginning with v_0 . As $\tilde{\Gamma}$ is G -vertex transitive, this implies that $\tilde{\Gamma}$ is (G, s) -arc transitive.

(b) Let Σ be a connected (G, s) -arc transitive graph of valency k and form the new graph Σ^* by placing a new vertex at the midpoint of every edge. We have seen that $G \leq \text{Aut}(\Sigma^*)$ and has two orbits Δ_1 and Δ_2 on vertices, with Δ_1 being the set of all vertices of Σ and Δ_2 the set of midpoints. Let $v_0 \in \Delta_1$ and consider the $(2s - 1)$ -arc $(v_0, p_0, v_1, p_1, \dots, v_{s-1}, p_{s-1})$ in Σ^* , where each p_i is the midpoint of the edge $\{v_i, v_{i+1}\}$. Then (v_0, v_1, \dots, v_s) is an s -arc in Σ . As G_{v_0} acts transitively on the set of s -arcs in Σ starting at v_0 , it also acts transitively on the set of $(2s - 1)$ -arcs in Σ^* emanating from v_0 . Thus to show that Σ^* is locally $(G, 2s - 1)$ -arc transitive it remains to show that for all $p_0 \in \Delta_2$, G_{p_0} acts transitively on the set of $(2s - 1)$ -arcs starting at p_0 . Given two such $(2s - 1)$ -arcs $\alpha = (p_0, v_1, p_1, \dots, p_{s-1}, v_s)$ and $\alpha' = (p_0, v'_1, p'_1, \dots, p'_{s-1}, v'_s)$ in Σ^* , there exist

s -arcs $\beta = (v_0, v_1, \dots, v_s)$ and $\beta' = (v'_0, \dots, v'_s)$ in Σ such that p_i is the midpoint of $\{v_i, v_{i+1}\}$ and p'_i is the midpoint of $\{v'_i, v'_{i+1}\}$ for each $i \geq 0$. As Σ is (G, s) -arc transitive, there exists $g \in G$ such that $\beta^g = \beta'$ and hence $\alpha^g = \alpha'$. However, p_0 is the midpoint of both $\{v_0, v_1\}$ and $\{v'_0, v'_1\}$, and so $\{v_0, v_1\} = \{v'_0, v'_1\}$. Hence $g \in G_{\{v_0, v_1\}} = G_{p_0}$. Thus G_{p_0} acts transitively on the set of $(2s - 1)$ -arcs starting at p_0 , so Σ^* is locally $(G, 2s - 1)$ -arc transitive. The last assertion of (b) follows from the discussion immediately preceding the statement of the theorem. \square

Corollary 3.11. *The connected locally $(G, 2s - 1)$ -arc transitive graphs of valency $\{2, k\}$, where $k \geq 3$, are in 1-1 correspondence with the connected (G, s) -arc transitive graphs of valency k .*

We now aim to give an upper bound for s when Γ has a vertex of valency two and a vertex of valency at least three.

Theorem 3.12. *Let Γ be a connected locally $(G, 2s)$ -arc transitive graph of valency $\{2, k\}$, where $k \geq 3$ and $s \geq 1$. Then Γ is locally $(G, 2s + 1)$ -arc transitive.*

Proof. Let Δ_1 and Δ_2 denote the sets of vertices of valency k and 2, respectively. As all vertices in Γ have valency at least two, Γ is locally $(G, 2s - 1)$ -arc transitive. Let $\tilde{\Gamma}$ be the connected component of the distance two graph of Γ with vertex set Δ_1 . Then by Theorem 3.10(a), $\tilde{\Gamma}$ is (G, s) -arc transitive. Let $(u_0, v_1, u_1, v_2, \dots, v_s, u_s)$ be a $2s$ -arc in Γ where each $v_i \in \Delta_1$ and $u_i \in \Delta_2$. Now as Γ is locally $(G, 2s)$ -arc transitive, $G_{u_0 v_1 u_1 \dots v_s}$ acts transitively on $\Gamma(v_s) \setminus \{u_{s-1}\}$. Then as each vertex of $\Gamma(v_s) \setminus \{u_{s-1}\}$ is adjacent to a unique vertex of $\Gamma_2(v_s) \setminus \{v_{s-1}\}$, it follows that $G_{u_0 v_1 u_1 \dots v_s}$ acts transitively on $\Gamma_2(v_s) \setminus \{v_{s-1}\}$. Now as u_0 has valency two, there exists a unique $v_0 \in \Gamma(u_0) \setminus \{v_1\}$. Thus $G_{u_0 v_1 u_1 \dots v_s} \leq G_{v_0}$. Furthermore, (v_0, v_1, \dots, v_s) is an s -arc in $\tilde{\Gamma}$ and $G_{v_0 v_1 \dots v_s}$ acts transitively on $\tilde{\Gamma}(v_s) \setminus \{v_{s-1}\} = \Gamma_2(v_s) \setminus \{v_{s-1}\}$. Hence G_{v_0} acts transitively on the set of $(s + 1)$ -arcs in $\tilde{\Gamma}$ starting at v_0 and so $\tilde{\Gamma}$ is $(G, s + 1)$ -arc transitive. As $\Gamma = (\tilde{\Gamma})^*$ it follows from Theorem 3.10(b) that Γ is locally $(2s + 1)$ -arc transitive. \square

Corollary 3.13. *Let Γ be a locally (G, s) -arc transitive graph of valency $\{2, k\}$ with $k \geq 3$. Then $s \leq 13$, and this bound can be attained.*

Proof. By Theorem 3.12 we may assume that s is odd. Then by Corollary 3.11, there exists a $(G, \frac{s+1}{2})$ -arc transitive graph of valency k . Thus by the result of Weiss [32], $\frac{s+1}{2} \leq 7$ and so $s \leq 13$. Moreover, examples of $(G, 7)$ -arc transitive graphs of valency at least three are well known (see [19]), and so by Corollary 3.11 locally $(G, 13)$ -arc transitive graphs exist. \square

4. EXAMPLES

We now prove our assertions about the existence of locally (G, s) -arc transitive graphs with $s \geq 2$ for certain quasiprimitive types. We have already seen in Lemma 3.3 that there are examples of locally (G, s) -arc transitive graphs with $s \geq 2$ such that G is quasiprimitive on both orbits of type HA, TW, AS or PA. Many more examples where G is of type AS on both orbits can be found in [6, page 98]. From Theorem 1.2 the only remaining possible type when G is quasiprimitive on both orbits is $\{SD, PA\}$, and the following example shows that such graphs do occur.

Example 4.1. $\{\text{SD, PA}\}$: a family of locally 3-arc transitive graphs of valency $\{q, q+1\}$ with amalgam

$$(\text{PSL}(2, q) \times C_2, (C_2^{2^f} : C_{q-1}) : C_2, (C_2^f : C_{q-1}) \times C_2).$$

Let $T = \text{PSL}(2, q)$ for $q = 2^f \geq 4$. Now T has a 2-transitive representation of degree $q+1$ and if H is the stabiliser of a point in this representation, then $H \cong C_2^f : C_{q-1}$. The subgroup H has a unique minimal normal subgroup $L \cong C_2^f$. Let $G = T \text{ wr } S_2$ and consider the graph $\Gamma = \text{Cos}(G, G_v, G_w)$, where

$$G_v = \{(t, t) : t \in T\} \times S_2$$

and

$$G_w = \{(l_1 h, l_2 h) : l_i \in L, h \in H\} : S_2.$$

As G_v is a maximal subgroup of G , Γ is connected. The action of G on $\Delta_1 = [G : G_v]$ is primitive of type SD while the action of G on $\Delta_2 = [G : G_w]$ is quasiprimitive of type PA. Now $G_v \cap G_w = \{(h, h) : h \in H\} \times S_2$. The action of G_v on $[G_v : G_v \cap G_w]$ is then equivalent to the 2-transitive representation of T of degree $q+1$. The set $\{(l, 1) : l \in L\}$ is a set of coset representatives for $G_v \cap G_w$ in G_v . For each $l \in L$, set $[l, 1] = (G_v \cap G_w)(l, 1)$. Now if $h \in H$, then $(h, h) : [l, 1] \mapsto [l^h, 1]$. Then as H induces the group C_{q-1} as a group of automorphisms of L , it follows that G_{vw} acts transitively on $\{[l, 1] : l \in L \setminus \{1\}\}$. Hence the permutation group induced by G_w on the set of cosets $[G_w : G_v \cap G_w]$ is 2-transitive of degree q . Thus Γ is locally $(G, 2)$ -arc transitive with G quasiprimitive of type $\{\text{SD, PA}\}$, and Γ is biregular of valency $\{q+1, q\}$.

Let v be the vertex in Δ_1 corresponding to the coset G_v and let w be the vertex of Δ_2 corresponding to the coset G_w . Let $w' \in \Gamma(v) \setminus \{w\}$. Then

$$G_{w'vw} \cong \{(h, h) : h \in C_{q-1}\} \times S_2$$

which still acts transitively on $\Gamma(w) \setminus \{v\}$. Also, given $u \in \Gamma(w) \setminus \{v\}$,

$$G_{uvw} = \{(l, l) : l \in L\} \times S_2$$

is still transitive on $\Gamma(v) \setminus \{w\}$. Hence Γ is locally $(G, 3)$ -arc transitive.

We now look at graphs where G acts quasiprimitively on Δ_1 but not quasiprimitively on Δ_2 . First we provide a family of examples for which G acts primitively of type HA on Δ_1 .

Example 4.2. HA stars: a family of locally 3-arc transitive graphs of valency $\{q, \frac{q^d-1}{q-1}\}$ with amalgam

$$(\text{GL}(d, q), [q] : ([q]^{d-1} : (C_q \times \text{GL}(d-1, q))), [q]^{d-1} : (C_q \times \text{GL}(d-1, q))).$$

Define Γ to be the bipartite graph whose vertices are all the points and lines of the affine space $\text{AG}(d, q)$ and with adjacency given by incidence. There are q^d points and $q^{d-1}(q^d-1)/(q-1)$ lines. Each point is adjacent to $(q^d-1)/(q-1)$ lines while each line is adjacent to q points. Let $\text{AGL}(d, q) \leq G \leq \text{AFL}(d, q)$. Then G is a group of automorphisms of Γ with two orbits on the vertex set: the set Δ_1 of points and the set Δ_2 of lines. Given a point p , G_p acts 2-transitively on the set of lines through p while for any line l , G_l acts 2-transitively on the points of l . Thus Γ is locally $(G, 2)$ -arc transitive and is biregular of valency $\{q, \frac{q^d-1}{q-1}\}$. In fact, it is not hard to show that Γ is locally $(G, 3)$ -arc transitive.

Now let N be the group of translations of G . Then N acts transitively on the set of points of $\text{AG}(d, q)$ but has $(q^d - 1)/(q - 1)$ orbits on the set of lines. Each of these orbits is a set of parallel lines which partitions the set of points. Thus each point is incident to a unique line in each N -orbit and of course each line in an N -orbit is incident with q points. Hence Γ_N is the star $K_{1, (q^d - 1)/(q - 1)}$. As N always has at least three orbits on Δ_2 , it follows from Lemma 5.6, which we will prove in the next section, that Γ is never locally $(G, 4)$ -arc transitive.

If we take k -dimensional affine subspaces, where $1 < k < d - 1$, instead of affine lines we get a G -locally primitive graph which is not locally $(G, 2)$ -arc transitive. Its quotient graph modulo N is again a star and its central vertex has valency

$$\frac{(q^d - 1)(q^{d-1} - 1) \dots (q^{d-k+1} - 1)}{(q^k - 1)(q^{k-1} - 1) \dots (q - 1)}.$$

The following family shows that examples also exist where G is primitive of type PA on Δ_1 .

Example 4.3. PA stars: a family of locally 3-arc transitive graphs of valency $\{k, n\}$ with amalgam

$$(S_{n-1} \text{ wr } S_k, S_n \times (S_{n-1} \text{ wr } S_{k-1}), S_{n-1}^k : S_{k-1}).$$

The Hamming graph $H(k, n)$ has vertices given by all the k -tuples with entries from a set Ω of size n . Let $k \geq 2, n \geq 5$, and let Δ_1 be the set of vertices of $H(k, n)$. Also let Δ_2 be the set of maximal cliques of $H(k, n)$. Define the graph Γ to have vertex set $\Delta_1 \cup \Delta_2$ and adjacency given by incidence. Then if $v = (\omega_1, \dots, \omega_k) \in \Delta_1$, the maximal cliques in Δ_2 adjacent to v are the k cliques $w = \{(\alpha, \omega_2, \dots, \omega_k) : \alpha \in \Omega\}, \{(\omega_1, \alpha, \omega_3, \dots, \omega_k) : \alpha \in \Omega\}, \dots, \{(\omega_1, \dots, \omega_{k-1}, \alpha) : \alpha \in \Omega\}$.

Let $G = S_n \text{ wr } S_k$. Then the action of G on Δ_1 is primitive of type PA. Also $G_v \cong S_{n-1} \text{ wr } S_k$ and $G_v^{\Gamma(v)} \cong S_k$ acting on k points. Now $G_w \cong S_n \times (S_{n-1} \text{ wr } S_{k-1})$ and $G_w^{\Gamma(w)} \cong S_n$ acting on n points. Thus Γ is locally $(G, 2)$ -arc transitive and is biregular of valency $\{k, n\}$. Let $u = \{(\omega_1, \alpha, \omega_3, \dots, \omega_k) : \alpha \in \Omega\}$. Then $u \in \Gamma(v)$ and $G_{uvw} \cong (S_{n-1})^2 \times (S_{n-1} \text{ wr } S_{k-2})$ which acts transitively on $\Gamma(w) \setminus \{v\}$. Also let $x = (\beta, \omega_2, \dots, \omega_k)$ for some $\beta \in \Omega \setminus \{\omega_1\}$. Then $x \in \Gamma(w)$ and $G_{xwv} \cong S_{n-2} \times (S_{n-1} \text{ wr } S_{k-1})$ which acts transitively on $\Gamma(v) \setminus \{w\}$. Thus Γ is locally $(G, 3)$ -arc transitive. Let $N = S_n^k \triangleleft G$. Then N is transitive on Δ_1 but has k orbits on Δ_2 , each orbit being the set of maximal cliques associated with a given coordinate. Thus $\Gamma_N = K_{1,k}$. Then by Lemma 5.6, Γ is not locally $(G, 4)$ -arc transitive when $k \geq 3$.

When $k = 2$ the vertices of Δ_1 have valency two. Also, given a vertex $\{(\alpha, \omega_2) : \alpha \in \Omega\}$ in Δ_2 the vertices at distance two are all vertices of the form $\{(\omega_1, \alpha) : \alpha \in \Omega\}$ for some $\omega_1 \in \Omega$. Then the connected component $\tilde{\Gamma}$ of the distance two graph of Γ with $V\tilde{\Gamma} = \Delta_2$ is $K_{n,n}$ and is $(G, 3)$ -arc transitive. Thus by Theorem 3.10, Γ is locally $(G, 5)$ -arc transitive when $k = 2$.

Note that for each $k \geq 2$ our last example provides us with a locally $(G, 3)$ -arc transitive graph Γ such that G has a normal subgroup N with k orbits on Δ_2 and $\Gamma_N = K_{1,k}$. Hence we have proved the following.

Lemma 4.4. Every star graph $K_{1,n}$ for $n \geq 2$ is the normal quotient of some locally 3-arc transitive graph and $K_{1,2}$ is the normal quotient of a locally 5-arc transitive graph.

Examples where G is quasiprimitive of type HS also occur.

Example 4.5. HS stars: a family of locally 3-arc transitive graphs of valency $\{q, q+1\}$ with amalgam

$$(\mathrm{PSL}(2, q), C_2^{2f} : C_{q-1}, C_2^f : C_{q-1}).$$

Let Γ be the graph from Example 4.1 and let $G = T \times T$ where $T = \mathrm{PSL}(2, q)$. Then Γ is locally $(G, 3)$ -arc transitive such that G acts primitively of type HS on Δ_1 but has two normal subgroups $T \times 1$ and $1 \times T$ which are both intransitive on Δ_2 .

All examples where G acts quasiprimitively of type HS on Δ_1 , yet has a normal subgroup which is intransitive on Δ_2 , will be determined in [12]. All such graphs are also of $\{\mathrm{SD}, \mathrm{PA}\}$ type. There are also examples for which G is almost simple.

Example 4.6. AS star: a locally 3-arc transitive graph of valency $\{3, 8\}$ with amalgam

$$((C_7 : C_3) \times S_3, \mathrm{PSL}(2, 7) \times C_2, (C_7 : C_3) \times C_2).$$

Let $G = \mathrm{P}\Gamma\mathrm{L}(3, 4)$ which is an almost simple group with socle $T = \mathrm{PSL}(3, 4)$. Let ϕ be a field automorphism of T and consider $C_T(\phi) \cong \mathrm{PSL}(3, 2) \cong \mathrm{PSL}(2, 7)$. Let $R = N_G(C_T(\phi))$. Then $R = \langle C_T(\phi), \phi \rangle \cong \mathrm{PSL}(2, 7) \times C_2 \leq \mathrm{P}\Sigma\mathrm{L}(3, 4)$. There are three conjugacy classes of subgroups of T isomorphic to $C_T(\phi)$, and they are fused together by G . In fact $G/T \cong S_3$ permutes the three conjugacy classes naturally. Let δ be the element of $\mathrm{PGL}(3, 4)$ corresponding to the matrix

$$\begin{pmatrix} \xi & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

where $\xi \in \mathrm{GF}(4)$ is of order three. Then $A = \langle \delta, \phi \rangle \cong S_3$ and $G = T : A$. Now let L_0 be the intersection of $C_T(\phi)$ and its two images under A . Then $L_0 = C_7 : C_3$ and its normaliser in G is the maximal subgroup $L = (C_7 : C_3) \times S_3$. (See [5].) Now $L \cap R = (C_7 : C_3) \times C_2$. The action of L on the set of cosets $[L : L \cap R]$ is equivalent to the action of S_3 on three points while the action of R on the set of cosets $[R : L \cap R]$ is equivalent to the action of $\mathrm{PSL}(2, 7)$ on 8 points. Both actions are 2-transitive and so $\Gamma = \mathrm{Cos}(G, L, R)$ is a locally $(G, 2)$ -arc transitive graph of valency $\{3, 8\}$. One can deduce that Γ is in fact locally $(G, 3)$ -arc transitive. Furthermore, since L is a maximal subgroup of G , Γ is connected.

Further examples of locally $(G, 3)$ -arc transitive graphs where G is quasiprimitive of type AS on Δ_1 but has a normal subgroup which is intransitive on Δ_2 can be found in [13]. Finally we provide an example where G is quasiprimitive of type TW on Δ_1 , so all types listed in Theorem 1.3 occur.

Example 4.7. TW star: a locally 2-arc transitive graph of valency $\{6, 16\}$ with amalgam $(A_6, C_2^4 : A_5, A_5)$. Let $T = A_5$, $P = A_6$, $Q = A_5 < P$, the stabiliser of the point 6 in the natural action and $\phi : Q \mapsto \mathrm{Inn}(T)$. Then $G = T \mathrm{twr}_\phi P$ has a normal subgroup $N \cong T^6$ and the action of G on the set of cosets of $L = P$ is primitive of type TW. Let $\{z_1, \dots, z_5, z_6 = 1\}$ be a transversal for Q in P and for each $i = 1, \dots, 6$, using the notation introduced in Section 2, let $T_i = \{f \in N : f(z_j) = 1 \text{ for all } j \neq i\}$. Then each T_i is isomorphic to T and P acts on the set $\{T_1, \dots, T_6\}$ such that Q is the stabiliser of T_6 . The subgroup Q also normalises

$T_1 \times \cdots \times T_5$. For $i = 1, \dots, 5$, let Q_i be the stabiliser in Q of T_i and let V_i be the Klein 4-group of Q_i . Note that $V_i \triangleleft Q_i$. Let $R_i = \{f \in T_i : f(z_i) \in V_i\}$. Then for each $q \in Q$ we have $R_i^q = R_{iq}$. Thus $M = R_1 \times \cdots \times R_5 \cong C_2^{10}$ is normalised by Q . Now A_5 does not have any irreducible representations of degree 10 over $\text{GF}(2)$ so Q normalises some proper subgroup of M . Calculations show that there exists R_0 in M of order 2^4 which is normalised by Q and Q acts irreducibly on R_0 . Then let $R = R_0 : Q \cong \text{ASL}(2, 4)$. Now $L \cap R = Q$ and the action of L on the set of cosets $[L : L \cap R]$ is equivalent to the action of A_6 on six points. Furthermore, the action of R on the set of cosets $[R : L \cap R]$ is equivalent to the 2-transitive action of $\text{ASL}(2, 4)$ on the 2^4 points of the affine plane $\text{AG}(2, 4)$. Thus the graph $\Gamma = \text{Cos}(G, L, R)$ is locally $(G, 2)$ -arc transitive and is biregular of valency $\{6, 16\}$. Now G acts primitively on $\Delta_1 = [G : L]$ but N has six orbits on $\Delta_2 = [G : R]$. Hence the quotient graph Γ_N is the star $K_{1,6}$. The graph Γ is not locally $(G, 3)$ -arc transitive as given, $v = L \in \Delta_1$, $w = R \in \Delta_2$ and $w' \in \Gamma(v) \setminus \{w\}$, we have $G_{w'vw} = A_4$ and there are 15 vertices in $\Gamma(w) \setminus \{v\}$.

5. NORMAL QUOTIENTS

In this section we investigate the normal quotients of a G -edge transitive bipartite graph Γ and in particular we prove Theorem 1.1. We begin by showing that taking quotients with respect to the orbits of a normal subgroup which is intransitive on both G -orbits, preserves local primitivity and local s -arc transitivity.

Lemma 5.1. *Let Γ be a connected G -locally primitive bipartite graph with G -orbits Δ_1 and Δ_2 on $V\Gamma$ and each $|\Delta_i| > 1$. Suppose that there exists $N \triangleleft G$ such that N is intransitive on Δ_1 and on Δ_2 . Then*

- (1) Γ is a cover of Γ_N .
- (2) N acts semiregularly on $V\Gamma$ and $G^{V\Gamma_N} \cong G/N$.
- (3) Γ_N is (G/N) -locally primitive. Furthermore, if Γ is locally (G, s) -arc transitive, then Γ_N is locally $(G/N, s)$ -arc transitive.

Proof. Let $v \in \Delta_1$ and let $B = v^N$. Choose $u \in \Gamma(v) \subseteq \Delta_2$ and set $C = u^N$. Then C is a block of imprimitivity for the action of G on Δ_2 , and hence $C \cap \Gamma(v)$ is a block of imprimitivity for the action of G_v on $\Gamma(v)$. As $N_v \triangleleft G_v$ and G_v acts primitively on $\Gamma(v)$, it follows that either $\Gamma(v) \subseteq C$ or $|\Gamma(v) \cap C| = 1$.

Suppose that $\Gamma(v) \subseteq C$. Then for every vertex $w \in \Delta_1$, the set $\Gamma(w)$ is contained in some N -orbit. Thus if B' is an N -orbit on Δ_1 containing a vertex adjacent to a vertex in C , then $\Gamma(B') \subseteq C$. It follows from the connectedness of Γ that $V\Gamma = C \cup \Gamma(C)$, contradicting the intransitivity of N on Δ_2 . Thus $|\Gamma(v) \cap C| = 1$. Similarly $|\Gamma(u) \cap B| = 1$ and so Γ is a cover of Γ_N .

Let K be the kernel of the action of G on the set of N -orbits on $V\Gamma$ and let $v \in V\Gamma$. Now K_v fixes each N -orbit setwise and since distinct vertices of $\Gamma(v)$ lie in distinct N -orbits, we have that K_v acts trivially on $\Gamma(v)$. Since Γ is connected it follows that K_v fixes all the vertices of Γ and hence $K_v = 1$. Since this is true for all v , K acts semiregularly on $V\Gamma$ and hence so does N . Furthermore, as $N \leq K$ and acts transitively on the orbits of K , we see that $K = N$. Thus $G^{V\Gamma_N} \cong G/N$ and so $G/N \leq \text{Aut}(\Gamma_N)$.

For a vertex v in the N -orbit B , the group NG_v fixes B , contains G_v and is transitive on B . Hence $G_B = NG_v$. Then as N is the kernel of the action of G on $V\Gamma_N$ and as each block in $\Gamma_N(B)$ contains exactly one vertex of $\Gamma(v)$, we have

that $G_B^{\Gamma_N(B)}$ is permutationally isomorphic to $G_v^{\Gamma(v)}$ and so is primitive. Thus Γ_N is (G/N) -locally primitive.

Let (B, B_1, \dots, B_s) and (B, C_1, \dots, C_s) be s -arcs in Γ_N . Choose $v \in B$. Then there exist unique $v_i \in B_i$ and $u_i \in C_i$ such that (v, v_1, \dots, v_s) and (v, u_1, \dots, u_s) are s -arcs in Γ . If Γ is locally (G, s) -arc transitive, then there exists $g \in G_v$ mapping one to the other. As the orbits of N form a system of imprimitivity for G , it follows that $g \in G_B$ and maps the s -arc (B, B_1, \dots, B_s) to (B, C_1, \dots, C_s) . Thus Γ_N is locally $(G/N, s)$ -arc transitive. \square

Note that if N is maximal in G subject to being intransitive on both orbits Δ_1 and Δ_2 , then each normal subgroup of G containing N is transitive on at least one G -orbit.

We now consider the question of faithfulness.

Lemma 5.2. *Let Γ be a connected G -locally primitive bipartite graph with G -orbits Δ_1 and Δ_2 on $V\Gamma$ of sizes n and n' , respectively. Then either $\Gamma \cong K_{n,n'}$ or G is faithful on both Δ_1 and Δ_2 .*

Proof. If either of n or n' is 1, then $\Gamma = K_{n,n'}$ so assume that both n and n' are at least 2. Let K_1 and K_2 be the kernels of G on Δ_1 and Δ_2 , respectively. Since G acts faithfully on $V\Gamma$, we know that $K_1 \cap K_2 = 1$.

Suppose that $K_1 \neq 1$ and note that K_1 acts faithfully on Δ_2 . Let B be a nontrivial orbit of K_1 on Δ_2 and u a vertex in B . Let v be a vertex in Δ_1 adjacent to u . Since K_1 fixes v , it follows that v is adjacent to every vertex in B . Furthermore, as $K_1 \triangleleft G_v$, the orbits of K_1 on $\Gamma(v)$ are blocks of imprimitivity for the action of G_v . Then as Γ is G -locally primitive, $\Gamma(v) = B$. As this holds for every v adjacent to a vertex of B , it follows from the connectedness of Γ that $\Gamma \cong K_{n,n'}$. The same holds if $K_2 \neq 1$. \square

Note that if either n or n' is at least three and neither is equal to one, then $K_{n,n'}$ is locally 3-arc transitive but not locally 4-arc transitive as there are 4-arcs $(v_0, v_1, v_2, v_3, v_4)$ for which $v_4 = v_0$ and some for which $v_4 \neq v_0$. The graph $K_{2,2}$ is a cycle of length four and is locally (G, s) -arc transitive for all $s \geq 1$ with $G = S_2 \times S_2$. On the other hand, the graphs $K_{1,n}$ for $n \geq 2$ are only locally 2-arc transitive as they do not contain any 3-arcs.

We prove the following lemma regarding edge transitive graphs with a vertex of valency one.

Lemma 5.3. *Let Γ be a connected G -edge transitive but not G -vertex transitive graph such that $|\Gamma(u)| = 1$ for some vertex u . Then Γ is a star $K_{1,k}$, and if G acts faithfully on both G -orbits on vertices, then $k = 1$, $\Gamma = K_2$ and $G = 1$.*

Proof. Let Δ_1 and Δ_2 be the G -orbits on $V\Gamma$. Without loss of generality we may suppose that $u \in \Delta_1$. Since u is joined by an edge to only one vertex of Δ_2 and Γ is connected, Γ is a star $K_{1,k}$. If G is faithful on Δ_2 , then $G = 1$ and hence $|\Delta_1| = 1$. Thus $\Gamma = K_{1,1} = K_2$. \square

Now assuming that G acts faithfully on both of its orbits we investigate the case where every nontrivial normal subgroup of G is transitive on at least one G -orbit.

Lemma 5.4. *Let Γ be a finite connected graph such that G has two orbits Δ_1 and Δ_2 on vertices and G acts faithfully on both orbits. Suppose that every nontrivial*

normal subgroup N of G is transitive on at least one of the Δ_i . Then G acts quasiprimively on at least one of its orbits.

Proof. Suppose that G is not quasiprimitive on either of the Δ_i . Then for each $i \in \{1, 2\}$ there exists $N_i \triangleleft G$ such that N_i is intransitive on Δ_i and transitive on Δ_{3-i} . Now $N_1 \cap N_2 \triangleleft G$ and so if nontrivial would be transitive on at least one Δ_i . But then N_1 and N_2 would both be transitive on the same set, a contradiction. Thus $N_1 \cap N_2 = 1$ and hence $N_1 \times N_2 \triangleleft G$.

As each N_i is transitive on Δ_{3-i} , it follows that $C_{\text{Sym}(\Delta_{3-i})}(N_i)$ is semiregular. Thus each N_i is semiregular on Δ_i and transitive on Δ_{3-i} . Therefore $|N_1|$ divides $|\Delta_1|$ and $|\Delta_2|$ divides $|N_1|$, so $|\Delta_2|$ divides $|\Delta_1|$. Similarly $|\Delta_1|$ divides $|N_2|$ and $|N_2|$ divides $|\Delta_2|$ implying that $|\Delta_1|$ divides $|\Delta_2|$. Thus $|\Delta_1| = |\Delta_2|$. Furthermore, $|N_1| = |\Delta_1| = |N_2|$, contradicting N_1 being intransitive on Δ_1 . Thus G must be quasiprimitive on at least one of the Δ_i . \square

Note that this is the only lemma necessary for the proof of Theorem 1.1 whose proof does not extend to the case of infinite locally s -arc transitive graphs.

We now analyse the structure of Γ in the case where G has a normal subgroup which is transitive on one G -orbit yet intransitive on the other. A proof of Lemma 5.5 will complete the proof of Theorem 1.1.

Lemma 5.5. *Let Γ be a connected G -locally primitive bipartite graph with G -orbits Δ_1 and Δ_2 on $V\Gamma$. Suppose there exists $N \triangleleft G$ such that N is transitive on Δ_1 but intransitive on Δ_2 . Then Γ_N is a star whose central vertex has valency the number of orbits of N on Δ_2 . Furthermore, for each vertex $v \in \Delta_1$ and N -orbit B in Δ_2 , $|B \cap \Gamma(v)| = 1$ and the vertex stabiliser N_v acts trivially on $\Gamma(v)$.*

Proof. Choose $u \in \Delta_2$ and $v \in \Delta_1$ such that $v \sim u$. Let $B = u^N$. For each $w \in B$, we have that $w = u^g$ for some $g \in N$ and $v^g \in \Gamma(u^g) = \Gamma(w)$, that is, each vertex of B is adjacent to some vertex in Δ_1 . Conversely, as N acts transitively on Δ_1 each vertex of Δ_1 is adjacent to some vertex in B . Thus Γ_N is a star whose central vertex has valency the number of orbits of N on Δ_2 .

Now the set $\Gamma(v) \cap B$ is an orbit of N_v on $\Gamma(v)$ and is hence a block for G_v . If $\Gamma(v) \subseteq B$, then $\Gamma(v') \subseteq B$ for all $v' \in \Delta_1$ since B and Δ_1 are N -orbits, and this contradicts the connectivity of Γ . Hence $|B \cap \Gamma(v)| = 1$ and so N_v acts trivially on $\Gamma(v)$. \square

Finally, in this section we state a lemma from [17] and give a short proof.

Lemma 5.6. *Let Γ be a locally (G, s) -arc transitive graph such that all vertices have valency at least two and G has a normal subgroup N which is transitive on Δ_1 but has at least three orbits on Δ_2 . Then $s \leq 3$.*

Proof. Let B_1, B_2 and B_3 be three orbits of N on Δ_2 . Choose $v_0 \in B_1$ and let $v_1 \in \Gamma(v_0)$. Then by Lemma 5.5, v_1 is adjacent to a unique vertex v_2 in B_2 . Let $v_3 \in \Gamma(v_2) \setminus \{v_2\}$. Then (v_0, v_1, v_2, v_3) is a 3-arc in Γ . Now by Lemma 5.5, there exist $u, w \in \Gamma(v_3)$ such that $u \in B_1$ and $w \in B_3$. Then (v_0, v_1, v_2, v_3, u) and (v_0, v_1, v_2, v_3, w) are 4-arcs in Γ for which there does not exist an element $g \in G_{v_0}$ mapping one to the other, since such a g would fix B_1 and so cannot map u to w . Hence Γ is not locally $(G, 4)$ -arc transitive. \square

6. QUASIPRIMITIVE ON BOTH ORBITS

In this section we analyse the case where G acts faithfully and quasiprimively on both of its orbits. In particular we prove Theorem 1.2.

We say that Γ is G -locally quasiprimitive if $G_v^{\Gamma(v)}$ is a quasiprimitive permutation group for every vertex v . We begin with two very useful lemmas.

Lemma 6.1. *Let Γ be a connected G -locally quasiprimitive graph. Suppose that Γ is bipartite and the orbits of G are the bipartite halves Δ_1 and Δ_2 . Suppose also that G acts faithfully and quasiprimively on Δ_1 and on Δ_2 . If $N \triangleleft G$, then N^{Δ_1} is regular if and only if N^{Δ_2} is regular.*

Proof. If there exists a vertex with valency one, then by Lemma 5.3, $\Gamma = K_2$ and the result is trivially true. So we assume $|\Gamma(v)| \geq 2$ for all $v \in V\Gamma$. Suppose that N^{Δ_1} is regular and N^{Δ_2} is not regular. Then for all $v \in \Delta_1$, $N_v = 1$, while there exists $u \in \Delta_2$ such that $N_u \neq 1$. This implies that N_u acts nontrivially on $\Gamma(u)$ and so

$$1 \neq N_u^{\Gamma(u)} \triangleleft G_u^{\Gamma(u)}.$$

As $G_u^{\Gamma(u)}$ is a quasiprimitive permutation group, N_u acts transitively on $\Gamma(u)$. Since G is quasiprimitive and faithful on Δ_2 , N is transitive on Δ_2 and so for all $w \in \Delta_2$, N_w acts transitively on $\Gamma(w)$. This implies that N is transitive on the set of edges of Γ . Thus N_v acts transitively on $\Gamma(v)$ which contradicts $N_v = 1$ as v has at least two neighbours. \square

Lemma 6.2. *Let Γ and G be as in Lemma 6.1 and let $N \triangleleft G$. If N is not regular on Δ_1 , then $N_v^{\Gamma(v)}$ is transitive for all $v \in V\Gamma$.*

Proof. If Γ has a vertex of valency one, then by Lemma 5.3, $\Gamma = K_2$ and $G = 1$ so no such N exists. Thus all vertices have valency at least two. By Lemma 6.1, N is not regular on Δ_2 either, and so for all $v \in V\Gamma$, $N_v \neq 1$. Suppose that there exists $v \in V\Gamma$ such that $N_v^{\Gamma(v)} = 1$. Then as Γ is connected and G acts faithfully on $V\Gamma$, there exists a path (v_0, v_1, \dots, v_r) in Γ with $v_0 = v$ such that N_v fixes v_0, \dots, v_{r-1} but not v_r . Now $N_v \leq N_{v_{r-1}}$ and N_v moves $v_r \in \Gamma(v_{r-1})$ so

$$1 \neq N_{v_{r-1}}^{\Gamma(v_{r-1})} \triangleleft G_{v_{r-1}}^{\Gamma(v_{r-1})}.$$

Now $G_{v_{r-1}}^{\Gamma(v_{r-1})}$ is quasiprimitive, so $N_{v_{r-1}}^{\Gamma(v_{r-1})}$ is transitive. The vertex v_{r-1} is in Δ_1 or in Δ_2 and so as N is transitive on each of the sets Δ_1 and Δ_2 , we have that N is transitive on the edges of Γ . This contradicts $N_v^{\Gamma(v)} = 1$ as v has at least two neighbours. Thus $N_v^{\Gamma(v)} \neq 1$ and arguing as above, $N_v^{\Gamma(v)}$ is transitive. \square

We are now in a position to determine the possible quasiprimitive types for G .

Proposition 6.3. *Let Γ be a finite connected G -locally primitive graph with G -orbits Δ_1 and Δ_2 on vertices. Suppose that G acts faithfully and quasiprimively on Δ_1 and on Δ_2 of type $\{X, Y\}$. Then either $X = Y$, or $\{X, Y\} = \{\text{SD}, \text{PA}\}$ or $\{\text{CD}, \text{PA}\}$.*

Proof. Note that $G^{\Delta_1} \cong G^{\Delta_2} \cong G$. So if G is of type HA, AS, HS or HC on one of the G -orbits, then it must have the same type on the other G -orbit, as such types are determined by the abstract structure and number of minimal normal subgroups of G .

If G is of type TW on one G -orbit, then G has a unique minimal normal subgroup N isomorphic to T^k for some finite nonabelian simple group T and N is regular on that orbit. Lemma 6.1 implies that N also acts regularly on the other G -orbit and so G must be of type TW on both orbits. Thus we may assume that $\{X, Y\} \subseteq \{\text{SD}, \text{CD}, \text{PA}\}$, and so the only thing that remains to be proved is that $\{X, Y\} \neq \{\text{SD}, \text{CD}\}$.

Suppose to the contrary that G is of type $\{\text{SD}, \text{CD}\}$ and without loss of generality assume that G is quasiprimitive of type SD on Δ_1 and quasiprimitive of type CD on Δ_2 . Let N be the unique minimal normal subgroup of G . Then $N \cong T^k$ for some finite nonabelian simple group T . Since G acts faithfully on Δ_1 , we may assume that $N < G < N \cdot (\text{Out}(T) \times S_k) < \text{Aut}(T) \text{ wr } S_k$ and for some $v \in \Delta_1$, we have $N_v = \{(t, \dots, t) : t \in T\}$. Hence a typical element of G is of the form $(t_1, \dots, t_k)(\sigma, \dots, \sigma)\pi$, where the $t_i \in T$, $\sigma \in \text{Aut}(T)$ and $\pi \in S_k$. Let $w \in \Gamma(v)$. Then as G is quasiprimitive on Δ_2 of type CD, $N_w = D_1 \times \dots \times D_l$ where each D_i is a full diagonal subgroup of T^m where $k = ml$, $l \geq 2$. Thus $D_1 = \{t, t^{\varphi_{12}}, \dots, t^{\varphi_{1m}} : t \in T\}$ where $\varphi_{12}, \dots, \varphi_{1m} \in \text{Aut}(T)$ and the other D_i are conjugates of D_1 under elements of G . So we may assume that $D_i = \{t, t^{\varphi_{i2}}, \dots, t^{\varphi_{im}} : t \in T\}$ for some $\varphi_{ij} \in \text{Aut}(T)$, and $G \leq T^k \cdot (\text{Out}(T) \times (S_m \text{ wr } S_l))$. Thus every element $g \in G_v$ is of the form $(\sigma, \dots, \sigma)\tau$ for $\sigma \in \text{Aut}(T)$ and $\tau \in S_m \text{ wr } S_l$. Now $N_v \cap N_w = \{(t, \dots, t) : t \in C\} \triangleleft G_v \cap G_w$ where C is the centraliser in T of all the φ_{ij} . If all the $\varphi_{ij} = 1$, then $N_v \leq N_w$ and so as $G_v^{\Gamma(v)}$ is primitive it follows that $N_v^{\Gamma(v)} = 1$. Then Lemma 6.2 implies that $N_v = 1$, which contradicts $|N_v| = |T|$. Thus at least one of the φ_{ij} is nontrivial and $C \neq T$. Let $g \in G_v \cap G_w$. Then $g = (\sigma, \dots, \sigma)\tau$ where σ normalises C and so g normalises the subgroup $A = C_1 \times \dots \times C_k$ where each $C_i = \{(c, \dots, c) : c \in C\} < D_i$. Let $H = \langle G_v \cap G_w, A \rangle$. Then $H = A(G_v \cap G_w)$ and $H \cap N_w = A \neq N_v \cap N_w$. Thus $G_v \cap G_w < H < G_w$, contradicting $G_v \cap G_w$ being maximal in G_w . Hence Γ cannot be of type $\{\text{SD}, \text{CD}\}$. \square

In the locally (G, s) -arc transitive case we can prove even more.

Lemma 6.4. *There are no connected locally (G, s) -arc transitive graphs with $s \geq 2$ such that G acts faithfully and quasiprimitively on both orbits and of type HC or CD on one.*

Proof. Suppose such a graph Γ exists, and that G is of type HC or CD on Δ_1 . Then G has socle $N = T_1 \times \dots \times T_k \cong T^k$ for some finite nonabelian simple group T and $k \geq 2$. Let $v \in \Delta_1$. Then there exists an integer $l \geq 2$ dividing k such that without loss of generality, $N_v = D_1 \times \dots \times D_l$ where each D_i is a full diagonal subgroup of $T_{k(i-1)/l+1} \times \dots \times T_{ki/l}$. If G is of type HC, then $l = k/2$ and for either HC or CD type, G_v permutes D_1, \dots, D_l transitively by conjugation. Now as N does not act regularly on Δ_1 , Lemma 6.2 implies that

$$1 \neq N_v^{\Gamma(v)} \triangleleft G_v^{\Gamma(v)}.$$

Then by a theorem of Burnside [3, Theorem 4.3] we see that $N_v^{\Gamma(v)} \cong T$. If $\ker(N_v^{\Gamma(v)})$ denotes the kernel of the action of N_v on $\Gamma(v)$, then $\ker(N_v^{\Gamma(v)}) \cong T^{l-1}$. Also, since $\ker(N_v^{\Gamma(v)}) \triangleleft N_v$, it follows that $\ker(N_v^{\Gamma(v)})$ is a product of $l - 1$ of the D_i . However, as G_v acts transitively by conjugation on $\{D_1, \dots, D_l\}$ and G_v normalises $\ker(N_v^{\Gamma(v)})$, this is a contradiction. \square

We saw in Lemma 3.3 that there are locally (G, s) -arc transitive graphs with G of type HA, AS, TW and PA on both G -orbits and by Lemma 6.4 there are no such graphs of type $\{X, Y\}$ with HC or CD in $\{X, Y\}$. Therefore by Proposition 6.3 if such a graph has type $\{X, Y\}$ with $X \neq Y$, then $\{X, Y\} = \{\text{SD}, \text{PA}\}$. Example 4.1 provides an example of such a graph. We now complete the proof of Theorem 1.2 by showing that locally (G, s) -arc transitive graphs of type HS or SD do not exist. The argument is similar to that in [22].

Lemma 6.5. *There are no connected locally (G, s) -arc transitive graphs with G acting faithfully on both orbits and quasiprimitive of type HS or SD.*

Proof. Suppose that such a graph Γ does exist. Let X be the socle of G . Then $X \cong T^k$ for some finite nonabelian simple group T and $k \geq 2$. We can identify Δ_1 and Δ_2 with the elements of T^{k-1} such that the action of X on T^{k-1} is given by

$$(t_1, \dots, t_k) : (a_1, \dots, a_{k-1}) \mapsto (t_k^{-1}a_1t_1, \dots, t_k^{-1}a_{k-1}t_{k-1}).$$

Now $T \cong X_v \triangleleft G_v$. Also, by Lemma 6.2, $X_v^{\Gamma(v)} \neq 1$ and so is isomorphic to T . Furthermore, since G_v acts 2-transitively on $\Gamma(v)$, a theorem of Burnside (see [3, Theorem 4.3]) implies that the action of X_v on $\Gamma(v)$ is primitive.

Let $v = (1_T, \dots, 1_T) \in \Delta_1$. Then $X_v = \{(t, \dots, t) : t \in T\}$. Also for $u \in \Gamma(v)$ we have that $X_u = \{(t, t^{\varphi_2}, \dots, t^{\varphi_k}) : t \in T\}$ for some $\varphi_2, \dots, \varphi_k \in \text{Aut}(T)$. Now $X_{v,u} = \{(t, \dots, t) : t \in C_T(\varphi_i) \text{ for all } i\}$. As $X_v \neq X_{v,u}$, at least one φ_i is not the identity. Furthermore, as $X_{v,u}$ is a maximal subgroup of X_v , we have that $X_{v,u} \cong C_T(\varphi_i)$. However, G_v is a 2-transitive almost simple group on $\Gamma(v)$ and no such group exists where the point stabiliser of the socle is a centraliser of a (possibly outer) automorphism. (See [3, Section 7.4].) Thus no such graph Γ exists. \square

It remains to provide examples of G -locally primitive graphs to prove our assertions about the existence of such graphs for the remaining possible quasiprimitive types for G -locally primitive graphs which are not possible for locally (G, s) -arc transitive graphs, namely the types $\{\text{CD}, \text{PA}\}$ and the types $X = Y \in \{\text{HS}, \text{SD}, \text{HC}, \text{CD}\}$. The constructions given in Example 6.6 below therefore complete the proof of Theorem 1.2.

Example 6.6. (1) (Type $\{\text{CD}, \text{PA}\}$ of valency $\{n^2, |A_{n-1}|^2\}$) Let $T = A_n$ for $n \geq 6$ and let $G = T^4 : (S_2 \text{ wr } S_2) = T \text{ wr } (S_2 \text{ wr } S_2)$. The action of G on the set of right cosets of the subgroup

$$G_v = \{(t, t, s, s) : t, s \in T\} : (S_2 \text{ wr } S_2)$$

is primitive of type CD. The group G also acts on the set of right cosets of $G_w = A_{n-1}^4 : (S_2 \text{ wr } S_2)$ with a primitive action of type PA. Consider the graph $\Gamma = \text{Cos}(G, G_v, G_w)$. Here G acts primitively on its two orbits on vertices with type $\{\text{CD}, \text{PA}\}$. Now

$$G_v \cap G_w = \{(t, t, s, s) : t, s \in A_{n-1}\} : (S_2 \text{ wr } S_2)$$

which is a maximal subgroup of both G_v and G_w . Actually G_v is primitive on $\Gamma(v)$ of type PA, and G_w is primitive on $\Gamma(w)$ of type CD. Thus Γ is a G -locally primitive connected graph which is biregular of valency $\{n^2, |A_{n-1}|^2\}$.

(2) (Type HS and SD of valency $|T : C_T(\sigma)|$) Let T be a finite nonabelian simple group with automorphism σ such that $C_T(\sigma)$ is a maximal subgroup of T . For example, $T = A_n$ and σ is the automorphism induced by conjugation by $(1, 2)$.

Let $G = T \times T$, $G_v = \{(t, t) : t \in T\}$ and $G_w = \{(t, t^\sigma) : t \in T\}$. Consider the graph $\Gamma = \text{Cos}(G, G_v, G_w)$. Then the actions of G on $\Delta_1 = [G : G_v]$ and $\Delta_2 = [G : G_w]$ are primitive of type HS. Also $\langle G_v, G_w \rangle = G$, so Γ is connected. Now $G_v \cap G_w = \{(t, t) : t \in C_T(\sigma)\}$ which is a maximal subgroup of both G_v and G_w . Hence Γ is G -locally primitive with G of type HS and of valency $|T : C_T(\sigma)|$.

Suppose that σ has order two and let $\overline{G} = G : S_2$. Then $\overline{G} \leq \text{Aut}(\Gamma)$ and acts primitively on both Δ_1 and Δ_2 with type SD. Furthermore, Γ is also \overline{G} -locally primitive.

(3) (Type HC and CD of valency $|T : C_T(\sigma)|^2$) Take T to be a finite nonabelian simple group with an automorphism σ of order two such that $C_T(\sigma)$ is a maximal subgroup of T . Let $G = (T^2 \times T^2) : S_2$ where S_2 is generated by the element $(1, 2)(3, 4)$. Then G has two minimal normal subgroups each isomorphic to T^2 . In addition, let

$$G_v = \{(t, s, t, s) : t, s \in T\} : S_2$$

and

$$G_w = \{(t, s, t^\sigma, s^\sigma) : t, s \in T\} : S_2.$$

Consider the graph $\Gamma = \text{Cos}(G, G_v, G_w)$. Then as $\langle G_v, G_w \rangle = G$, Γ is connected and the actions of G on Δ_1 and Δ_2 are quasiprimitive of type HC. Now

$$G_v \cap G_w = \{(t, s, t, s) : t, s \in C_T(\sigma)\} : S_2$$

which is a maximal subgroup of both G_v and G_w . In fact G_v and G_w are both primitive permutation groups of type PA. Thus Γ is a G -locally primitive graph of valency $|T : C_T(\sigma)|^2$.

Now if we let $\overline{G} = (T^2 \times T^2) : (S_2 \text{ wr } S_2)$, where $S_2 \text{ wr } S_2$ preserves the partition $\{\{1, 3\}, \{2, 4\}\}$, we see that $\overline{G} \leq \text{Aut}(\Gamma)$. Also \overline{G} acts quasiprimitively of type CD on both Δ_1 and Δ_2 . Furthermore, Γ is \overline{G} -locally primitive.

7. QUASIPRIMITIVE ON ONLY ONE ORBIT

In this section we prove Theorem 1.3. We begin with the following lemma.

Lemma 7.1. *Let Γ be a G -edge transitive connected graph such that G acts faithfully on its two orbits Δ_1 and Δ_2 on vertices. Suppose that G has a nontrivial normal subgroup N such that $N_v^{\Gamma(v)} = 1$ for all $v \in \Delta_1$. If there exists $w \in \Delta_2$ such that $N_w^{\Gamma(w)} = 1$, then N acts semiregularly on $V\Gamma$.*

Proof. As G is transitive on Δ_2 , $N_w^{\Gamma(w)} = 1$ for all $w \in \Delta_2$. The connectivity of Γ then implies that N_v fixes every element of Δ_1 and Δ_2 . Then since G acts faithfully on Δ_1 , $N_v = 1$ and so N acts semiregularly on Δ_1 . Also for $w \sim v$, the subgroup N_w is contained in $N_v = 1$. Hence $N_w = 1$ and so N acts semiregularly on Δ_2 . \square

We can now prove Theorem 1.3.

Proof. Let Γ be a locally (G, s) -arc transitive connected graph with $s \geq 2$ such that G acts faithfully on its two orbits Δ_1 and Δ_2 but quasiprimitively on only Δ_1 . Suppose first that G is quasiprimitive of type HC, SD or CD. Let $X = \text{soc}(G)$. Then for $v \in \Delta_1$, X_v is a subdirect subgroup of X , so $X_v \cong T^r$ for some $r \geq 1$. Suppose that $X_v^{\Gamma(v)} = 1$. Then as X does not act regularly on Δ_1 , Lemma 7.1 implies that $X_w^{\Gamma(w)} \neq 1$ for all $w \in \Delta_2$. Now let $w \in \Gamma(v)$ such that X_w moves v , so $X_w \neq X_v$. Since $X_v^{\Gamma(v)} = 1$ it follows that $X_v < X_w$. Then as X_v is a subdirect

subgroup of X , so is X_w . Thus $X_w \cong T^l$ for some $l > r$. Furthermore, since $X_w^{\Gamma(w)}$ is a nontrivial normal subgroup of the 2-transitive group $G_w^{\Gamma(w)}$, a theorem of Burnside [3, Theorem 4.3] implies that $X_w^{\Gamma(w)} = T$ and is a primitive group. Thus $X_w \cong T^{r+1}$ and $(X_w)_v = X_v \cong T^r$. Since the kernel of X_w on $\Gamma(w)$ is contained in $(X_w)_v$, it follows that $(X_w)_v$ is equal to the kernel, that is, $X_w^{\Gamma(w)}$ is regular. This contradicts $X_w^{\Gamma(w)}$ being primitive, and we deduce that $X_v^{\Gamma(v)} \neq 1$.

Now X is transitive on Δ_1 , since G is quasiprimitive on Δ_1 . By Lemma 5.5 and the fact that $X_v^{\Gamma(v)} \neq 1$, it follows that X must also be transitive on Δ_2 . Since G is not quasiprimitive on Δ_2 it follows that $X = \text{soc}(G)$ is not a minimal normal subgroup of G and hence G has type HC. Since $X_v \cong T^r$ we have $X \cong T^{2r}$ and $r \geq 2$. We have seen that $X_v^{\Gamma(v)}$ is a nontrivial normal subgroup of the 2-transitive group $G_v^{\Gamma(v)}$. Thus our favourite theorem of Burnside implies that $X_v^{\Gamma(v)} \cong T$. However, X_v is a minimal normal subgroup of G_v and so since it acts nontrivially on $\Gamma(v)$, it acts faithfully on $\Gamma(v)$. Thus $X_v^{\Gamma(v)} \cong T^r$ which is a contradiction since $r \geq 2$. Thus G is of type HA, HS, AS, PA or TW. We saw in Section 4 that examples of locally s -arc transitive graphs for the five types HA, HS, AS, PA and TW do occur. This completes the proof of Theorem 1.3. \square

Although there are no locally $(G, 2)$ -arc transitive graphs satisfying the conditions of Theorem 1.3 with G quasiprimitive on Δ_1 of type HC, SD or CD, there are locally primitive graphs of these types as the following examples demonstrate.

Example 7.2. (1) (HC stars of valency $\{n^2, |A_{n-1}|^2\}$) Let Γ be as in the first example of Example 6.6 and let $G = T^4 : S_2$ where $T = A_n$ and S_2 is generated by the element $(1, 2)(3, 4)$. Then Γ is G -locally primitive such that the action of G on $\Delta_1 = [G : G_v]$ is primitive of type HC. Let $N = T \times T \times 1 \times 1 \triangleleft G$. Then N acts regularly on Δ_1 but has n^2 orbits on Δ_2 . Thus Γ_N is the star K_{1, n^2} .

(2) (Type SD of valency $\{3, |T|\}$) Let $G = T^3 : S_3$ for some nonabelian simple group T and consider the subgroups

$$G_v = \{(t, t, t) : t \in T\} \times S_3$$

and

$$G_w = \{(s, t, t) : s, t \in T\} : \langle (2, 3) \rangle.$$

Then the graph $\Gamma = \text{Cos}(G, G_v, G_w)$ is connected, and G acts primitively of type SD on $\Delta_1 = [G : G_v]$. Now

$$G_v \cap G_w = \{(t, t, t) : t \in T\} \times \langle (2, 3) \rangle$$

which is a maximal subgroup of both G_v and G_w . Thus Γ is G -locally primitive and biregular of valency $\{n^2, |A_{n-1}|^2\}$. The group G does not act quasiprimitively on Δ_2 , as the normal subgroup $N = T^3$ has three orbits. Thus Γ_N is the star $K_{1, 3}$.

(3) (Type CD of valency $\{9, |T|^2\}$) Let $G = T^6 : (S_3 \text{ wr } S_2)$ for some nonabelian simple group T . Let

$$G_v = \{(t, t, t, s, s, s) : t, s \in T\} : (S_3 \text{ wr } S_2)$$

and

$$G_w = [(T \times (\{(t, t) : t \in T\} \times S_2)) \times (T \times (\{(t, t) : t \in T\} \times S_2))] : S_2.$$

Then the graph $\Gamma = \text{Cos}(G, G_v, G_w)$ is connected and G acts primitively of type CD on $\Delta_1 = [G : G_v]$. Now

$$G_v \cap G_w = \{(t, t, t, s, s, s) : t, s \in T\} : (((\langle 2, 3 \rangle) \times \langle (2, 3) \rangle) : S_2)$$

which is a maximal subgroup of G_v and G_w . Thus Γ is G -locally primitive and biregular of valency $\{9, |T|^2\}$. Now G does not act quasiprimatively on Δ_2 as N has nine orbits, so Γ_N is the star $K_{1,9}$.

8. SOME OPEN PROBLEMS

The work of this paper directs our attention to studying the “basic” locally (G, s) -arc transitive graphs with $s \geq 2$, that is, upon the following two problems.

Problem 8.1. Further analyse the structure and properties of locally (G, s) -arc transitive graphs, for which G acts faithfully and quasiprimatively on both orbits of G on vertices.

Problem 8.2. Further analyse the structure and properties of locally (G, s) -arc transitive graphs, for which G acts faithfully on both orbits Δ_1 and Δ_2 on vertices, but G acts quasiprimatively only on Δ_1 .

For some of the quasiprimitive types given in the first problem it would seem feasible to determine, or at least provide a nice characterisation of, such graphs. The $\{\text{SD}, \text{PA}\}$ case is currently being considered by the authors. In the case where G acts quasiprimatively of type PA on both orbits, the only known examples are the standard double covers of (G, s) -arc transitive graphs as given in Lemma 3.3. It would be interesting to find examples of PA type which did not arise in this way.

The second problem has been looked at in [12] where a nice characterisation of such graphs has been found which enables the vertices of Δ_2 to be represented by certain subsets of Δ_1 . In the case where G is quasiprimitive of type HA, HS or TW on Δ_1 , we can identify Δ_1 with a minimal normal subgroup N of G and the problem is reduced to finding certain collections of p -subgroups of N .

The paper [10] determines all locally (G, s) -arc transitive graphs where G is a Ree simple group acting quasiprimatively on both of its orbits. Three new infinite families of examples are constructed.

Problem 8.3. Classify all locally (G, s) -arc transitive graphs for other families of small rank Lie type groups G .

Placing further restrictions on s decreases the possibilities for the “basic” graphs. For example, Lemma 5.6 implies that when $s \geq 4$, we need to study only the case where G acts quasiprimatively on both orbits. When G is quasiprimitive of type HA on both G -orbits, it was shown in [15, Lemma 3.2] that Γ has an automorphism τ of order two which interchanges Δ_1 and Δ_2 . Then $\langle N, \tau \rangle$ is a regular normal subgroup of $\overline{G} = \langle G, \tau \rangle$. Hence by [18, Proposition 3.2], Γ is not (\overline{G}, s) -arc transitive for $s \geq 4$ and so Γ is not locally (G, s) -arc transitive for $s \geq 4$.

Question 8.4. Given a locally (G, s) -arc transitive graph with $s \geq 3$ and G acting faithfully on both Δ_1 and Δ_2 and quasiprimatively on at least one, what are the possible quasiprimitive types for G ? Similarly, find the possible quasiprimitive types for G when $s \geq 4$.

Given a locally (G, s) -arc transitive graph Γ it would be interesting to determine the full automorphism group. This leads us to the following question.

Question 8.5. When is a locally (G, s) -arc transitive graph $\text{Aut}(\Gamma)$ -vertex transitive?

Clearly Γ has to be regular, and so determining when Γ is regular would be a good first step. In Section 3 we discussed some sufficient conditions when all vertex stabilisers were conjugate in G . A regular graph which is edge transitive but not vertex transitive is called *semisymmetric*. Such graphs have been studied extensively; see for example [9], [11] and [15]. The smallest number of vertices for a semisymmetric graph is 80, see [8], and there are two such graphs, one of which is a generalised 4-gon and hence locally 5-arc transitive.

In [31], Weiss conjectured that for a G -vertex transitive, G -locally primitive graph of valency k , the order $|G_v|$ of a vertex stabiliser is bounded above by a function of k . For the nonbipartite case, it was proved in [4] that the conjecture holds if and only if it holds in the case where G is an almost simple group. In [20] it was conjectured that the same result holds for the more general class of G -vertex transitive G -locally quasiprimitive graphs. It was mentioned in [31] that the case where Γ is not G -vertex transitive should also be investigated. As currently stated the conjecture does not hold even for locally 3-arc transitive graphs. Vertices in a G -locally primitive graph have valency k_1 if they lie in Δ_1 and k_2 if they lie in Δ_2 with k_1 and k_2 not necessarily equal. There are infinite families of graphs for which k_2 is constant while k_1 , and hence $|G_v|$ for some vertex v of valency k_2 , are unbounded. For example, the graphs in Example 4.2 constructed from the points and lines of $\text{AG}(d, q)$ for a fixed q and unbounded d . With this in mind we make the following conjecture.

Conjecture 8.6. *If Γ is a connected G -locally primitive graph such that G has two orbits on vertices and Γ is biregular of valency $\{k_1, k_2\}$, then the order $|G_e|$ of an edge stabiliser is bounded by a function of $\max\{k_1, k_2\}$.*

REFERENCES

- [1] Robert W. Baddeley, *Two-arc transitive graphs and twisted wreath products*, J. Algebraic Combin. **2** (1993), 215–237. MR **94h**:05037
- [2] I. Z. Bouwer and D. Ž. Djoković, *On regular graphs. III*, J. Combinatorial Theory (B) **14** (1973), 268–277. MR **47**:4858
- [3] Peter J. Cameron, *Permutation groups*, London Mathematical Society Student Texts, no. 45, Cambridge University Press, Cambridge, 1999. MR **2001c**:20008
- [4] Marston D. Conder, Cai Heng Li, and Cheryl E. Praeger, *On the Weiss conjecture for finite locally primitive graphs*, Proc. Edinburgh Math. Soc. **43** (2000), 129–138. MR **2001e**:05054
- [5] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson, *Atlas of finite groups*, Clarendon Press, Oxford, 1985. MR **88g**:20025
- [6] A. Delgado, D. Goldschmidt, and B. Stellmacher, *Groups and graphs: new results and methods*, Birkhäuser Verlag, Basel, 1985. MR **88a**:05076
- [7] P. Dembowski, *Finite geometries*, Springer-Verlag, Berlin, 1968. MR **38**:1597
- [8] Shaofei Du and Dragan Marušić, *Biprimitive graphs of smallest order*, J. Algebraic Combin. **9** (1999), 151–156. MR **2000b**:05069
- [9] Shaofei Du and Mingyao Xu, *A classification of semisymmetric graphs of order $2pq$* , Comm. Algebra **28** (2000), 2685–2715. MR **2001g**:05059
- [10] Xin Gui Fang, Cai Heng Li, and Cheryl E. Praeger, *On locally two-arc transitive graphs admitting a Ree simple group*, in preparation.
- [11] Jon Folkman, *Regular line-symmetric graphs*, J. Combinatorial Theory **3** (1967), 215–232. MR **37**:97
- [12] Michael Giudici, Cai Heng Li, and Cheryl E. Praeger, *Characterising finite locally s -arc transitive graphs with a star normal quotient*, in preparation.

- [13] ———, *Some locally 3-arc transitive graphs constructed from triality*, submitted.
- [14] David M. Goldschmidt, *Automorphisms of trivalent graphs*, Ann. of Math. (2) **111** (1980), 377–406. MR **82a**:05052
- [15] A. A. Ivanov and M. E. Iofinova, *Biprimitive cubic graphs*, Investigations in the algebraic theory of combinatorial objects, Kluwer, 1993, pp. 459–472.
- [16] A. A. Ivanov and Cheryl E. Praeger, *On finite affine 2-arc transitive graphs*, Europ. J. Combinatorics **14** (1993), 421–444. MR **94k**:05089
- [17] Cai Heng Li, *On finite moufang graphs*, in preparation.
- [18] ———, *Finite s -arc transitive graphs of prime-power order*, Bull. London Math. Soc. **33** (2001), 129–137. MR **2002d**:05064
- [19] ———, *The finite vertex-primitive and vertex-biprimitive s -transitive graphs for $s \geq 4$* , Trans. Amer. Math. Soc. **353** (2001), 3511–3529. MR **2002c**:05084
- [20] Cai Heng Li, Cheryl E. Praeger, Akshay Venkatesh, and Sanming Zhou, *Finite locally-quasiprimitive graphs*, Discrete Math. **246** (2002), 197–218. MR **2003c**:05110
- [21] Martin W. Liebeck, Cheryl E. Praeger, and Jan Saxl, *On the O’Nan–Scott Theorem for finite primitive permutation groups*, J. Austral. Math. Soc. (A) **44** (1988), 389–396. MR **89a**:20002
- [22] Cheryl E. Praeger, *On a reduction theorem for finite, bipartite 2-arc-transitive graphs*, Australas J. Comb. **7** (1993), 21–36. MR **93m**:05091
- [23] Cheryl E. Praeger, *An O’Nan–Scott theorem for finite quasiprimitive permutation groups and an application to 2-arc transitive graphs*, J. London Math. Soc. (2) **47** (1993), 227–239. MR **94f**:05068
- [24] Cheryl E. Praeger, *Finite quasiprimitive graphs*, Surveys in combinatorics, 1997. Proceedings of the 16th British combinatorial conference, London, UK, July 1997 (R. A. Bailey, ed.), Lond. Math. Soc. Lect. Note Ser., no. 241, Cambridge University Press, 1997, pp. 65–85. MR **99b**:05076
- [25] Gert Sabidussi, *Vertex-transitive graphs*, Monatsh. Math. **68** (1964), 426–438. MR **31**:91
- [26] D. H. Smith, *Primitive and imprimitive graphs*, Quart. J. Math. Oxford (2) **22** (1971), 551–557. MR **48**:5926
- [27] Bernd Stellmacher, *Locally s -transitive graphs*, unpublished.
- [28] Michio Suzuki, *Group theory. I*, Springer-Verlag, Berlin, 1982. MR **82k**:20001c
- [29] W. T. Tutte, *A family of cubical graphs*, Proc. Cambridge Philos. Soc. **43** (1947), 459–474. MR **9**:97g
- [30] ———, *On the symmetry of cubic graphs*, Canad. J. Math. **11** (1959), 621–624. MR **22**:679
- [31] R. Weiss, *s -transitive graphs*, Algebraic methods in graph theory, Vol. I, II (Szeged, 1978), North-Holland, Amsterdam, 1981, pp. 827–847. MR **83b**:05071
- [32] Richard Weiss, *The nonexistence of 8-transitive graphs*, Combinatorica **1** (1981), 309–311. MR **84f**:05050
- [33] Richard Weiss, *Generalized polygons and s -transitive graphs*, Finite geometries, buildings, and related topics (Pingree Park, CO, 1988), Oxford Univ. Press, New York, 1990, pp. 95–103. MR **91j**:51011

SCHOOL OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF WESTERN AUSTRALIA, 35 STIRLING HIGHWAY, CRAWLEY, WESTERN AUSTRALIA 6009, AUSTRALIA
E-mail address: `giudici@maths.uwa.edu.au`

SCHOOL OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF WESTERN AUSTRALIA, 35 STIRLING HIGHWAY, CRAWLEY, WESTERN AUSTRALIA 6009, AUSTRALIA
E-mail address: `li@maths.uwa.edu.au`

SCHOOL OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF WESTERN AUSTRALIA, 35 STIRLING HIGHWAY, CRAWLEY, WESTERN AUSTRALIA 6009, AUSTRALIA
E-mail address: `praeger@maths.uwa.edu.au`