STRONG L^p -SOLUTIONS TO THE NAVIER-STOKES FLOW PAST MOVING OBSTACLES: THE CASE OF SEVERAL OBSTACLES AND TIME DEPENDENT VELOCITY

EVA DINTELMANN, MATTHIAS GEISSERT, AND MATTHIAS HIEBER

ABSTRACT. Consider the Navier-Stokes flow past several moving or rotating obstacles with possible time-dependent velocity. It is shown that under suitable assumptions on the data, there exists a unique, local strong solution in the $L^p - L^q$ -setting for suitable $p, q \in (1, \infty)$. Moreover, it is proved that this strong solution coincides with the known mild solution in the very weak sense.

1. INTRODUCTION

The mathematical description of the Navier-Stokes flow past rotating or moving obstacles gained quite a bit of attention in the last years. The motion is hereby described by the equations of Navier-Stokes in an exterior domain depending on the time variable t . More precisely, consider the equation

$$(1.1) \quad \begin{cases} \partial_t v - \Delta v + (v \cdot \nabla)v + \nabla q &= f & \text{in } \Omega(t), & t \in (0, T), \\ \operatorname{div} v &= 0 & \text{in } \Omega(t), & t \in (0, T), \\ v(x, t) &= M_i(t)x & \text{on } \Gamma_i(t), & i = 1, \dots, m, \quad t \in (0, T), \\ v(x, 0) &= v_0(x) & \text{in } \Omega(0). \end{cases}$$

Here $v = v(x, t)$ and $q = q(x, t)$ denote the velocity and the pressure of the fluid, respectively. In this paper we consider time-dependent domains $\Omega(t)$ of the following form: let $\mathcal{O}_1, \dots, \mathcal{O}_m \subset \mathbb{R}^n$, $n \geq 2$, be compact sets with boundaries $\Gamma_1, \dots, \Gamma_m$ of class $C^{1,1}$. We denote by $\Omega := \mathbb{R}^n \setminus \bigcup_{i=1}^m \mathcal{O}_i$ the exterior domain. For time-dependent matrices

$$M_i \in C^\infty([0, T]; \mathbb{R}^{n \times n}),$$

with $\operatorname{tr} M_i(t) = 0$ for all $t \in [0, T]$, $i \in \{1, \dots, m\}$, we define the time dependent exterior domain

$$\Omega(t) := \mathbb{R}^n \setminus \bigcup_{i=1}^m \mathcal{O}_i(t)$$

with $\mathcal{O}_i(t) := \{y = G_{(i)}(t)x, x \in \mathcal{O}_i\}$, $\Gamma_i(t) := \{y = \overline{G_{(i)}(t)}x, x \in \Gamma_i\}$ for $t \in [0, T]$ and a suitably defined isomorphism $G_{(i)}(t) : \overline{\mathcal{O}_i} \rightarrow \overline{\mathcal{O}_i(t)}$; for details see (2.5). As the obstacles shall not collide, we require

$$\operatorname{dist} \left(\overline{\mathcal{O}_i(t)}, \overline{\mathcal{O}_j(t)} \right) > 0, \quad i \neq j, \quad t \in [0, T].$$

Received by the editors July 28, 2006.

2000 *Mathematics Subject Classification*. Primary 76D03, 35Q30, 35B30.

Key words and phrases. Navier-Stokes equations, rotating obstacles, strong L^p -solutions.

The boundary condition on $\Gamma_i(t)$ is the usual no-slip condition. It is the aim of this paper to construct a strong L^p -solution to (1.1).

It is interesting to compare our solution to problem (1.1) with the results which have recently been obtained by several different approaches. The situation of *one* obstacle rotating with *constant* angular velocity (i.e. M equals the rotation matrix) was first considered by Hishida [His99]. He proved the existence of a unique local *mild* solution to (1.1) in the context of L^2 .

Strong solutions, again in the L^2 -context and for one obstacle, were obtained by Galdi and Silvestre in [GS05] using Galerkin methods as well as by Cumsille and Tucsnak [CT06]. In the case of two dimensions, these strong solutions are even global in time under appropriate assumptions on the data; see [CT06].

The situation where the data belong to L^p for $1 < p < \infty$ was first considered in [GHH06a], where the existence of a unique, local, *mild* L^p -solution to (1.1) was established. It was shown by Hishida and Shibata in [HS06] that this solution is even a global one, provided the data are small enough.

In this paper, we consider the situation of *strong* L^p -solutions. We prove the existence of a local, strong solution to (1.1) in L^p even for *several* non-colliding obstacles which may rotate or move with a *time-dependent* angular velocity. One of the main tools in the proof of our results will be the maximal $L^p - L^q$ -regularity of the Stokes operator in exterior domains.

It is a natural question to ask whether the strong solution to (1.1) obtained in Theorem 3.1 below coincides with the mild solution constructed in [GHH06a]. We give an affirmative answer to this question in Theorem 3.3 below. Of course, we need to explain first the meaning of *coincides*, since mild and strong solutions are defined on different spaces. In this section we make use of the concept of *very weak solutions* which was introduced in [FJR72] for $\mathbb{R}^n$ and in [Ama00] for domains.

For more information about the Navier-Stokes equation in the rotating framework of all of $\mathbb{R}^n$ or $\mathbb{R}_+^n$, we refer to the papers [CM97], [BMN99], [HS05], [GIMM04] and [GIMMS05] dealing in particular with data non-decaying at infinity.

2. PRELIMINARIES

We start by transforming equation (1.1) defined in the time-dependent domain $\Omega(t)$ to an equation on a fixed cylindrical domain. More precisely, following the approach introduced by Inoue and Wakimoto [IW77], we introduce a change of coordinates which coincides in the special case of pure rotation, i.e. M equals the rotation matrix, with the rotation in a neighborhood of the rotating obstacle, but equals the identity far away from the rotating body; see also [CT06].

For the time being, assume there is only one moving obstacle. We then make the following assumption:

- (A1) Let $\mathcal{O} \subset \mathbb{R}^n$, $n \geq 2$, be a compact set (the obstacle) with boundary $\Gamma := \partial\mathcal{O}$ of class $C^{1,1}$. Denote by $\Omega := \mathbb{R}^n \setminus \mathcal{O}$ the exterior domain corresponding to $\mathcal{O}$. For $M \in \mathbb{R}^{n \times n}$ with $\text{tr } M = 0$ define $\Omega(t), \Gamma(t), \mathcal{O}(t)$ as $\Omega(t) := \{y = e^{tM}x, x \in \Omega\}$, $\Gamma(t) := \{y = e^{tM}x, x \in \Gamma\}$, $\mathcal{O}(t) := \{y = e^{tM}x, x \in \mathcal{O}\}$.

We then consider a ball $B_r(0)$ of radius $r > 0$ such that $B_r(0) \supset \mathcal{O}(t)$ for all $t \in [0, T]$ for some $T > 0$. Choose a cut-off function $\eta \in C_c^\infty(\mathbb{R}^n)$ satisfying $0 \leq \eta \leq 1$, $\eta = 1$ on $B_r(0)$ and $\eta = 0$ on $B_{2r}(0)^c$. Define $b : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by

$$(2.1) \quad b(y) := \eta(y)My - B_K((\nabla\eta)M\cdot)(y),$$

where $K := \text{supp}(\nabla\eta)$. Here B_K denotes the Bogovskiĭ operator. For the definition and properties of this operator, we refer e.g. to [Bog79], [Gal94], and [GHH06b]. Note that $b(y) = My$ for $y \in \overline{\mathcal{O}(t)}$. Moreover, since $\int_K (\nabla\eta)(y) My \, dy = 0$, thanks to $\text{tr } M = 0$, it follows by construction that $\text{div } b = 0$ in $\mathbb{R}^n$ and $b \in C_c^\infty(\mathbb{R}^n; \mathbb{R}^n)$.

Then consider the initial value problem

$$(2.2) \quad \begin{cases} \partial_t X(y, t) &= b(X(y, t)), & t > 0, & y \in \mathbb{R}^n, \\ X(y, 0) &= y, & & y \in \mathbb{R}^n. \end{cases}$$

Then, by standard theory of ODE's, there exists a unique vector field

$$X \in C^\infty(\mathbb{R}^n \times \mathbb{R}_+)$$

satisfying (2.2). Moreover, $X(\cdot, t)$ is a C^∞ -diffeomorphism from Ω onto $\Omega(t)$. Its inverse $Y(\cdot, t)$ is of the same class of regularity and satisfies the initial value problem

$$(2.3) \quad \begin{cases} \partial_t Y(x, t) &= -b(Y(x, t)), & t > 0, & x \in \mathbb{R}^n, \\ Y(x, 0) &= x, & & x \in \mathbb{R}^n. \end{cases}$$

In fact, we only need the restrictions of X and Y on $[0, T]$, nevertheless even (2.2) and (2.3) can be solved on the whole of $\mathbb{R}^n \times \mathbb{R}_+$.

Denote by $J_X(\cdot, t)$ and $J_Y(\cdot, t)$ the Jacobian of $X(\cdot, t)$ and $Y(\cdot, t)$, respectively. Since $\text{div } b = 0$, Liouville's theorem, see e.g. [Arn92], implies that

$$(2.4) \quad J_X(y, t) J_Y(X(y, t), t) = \text{id} \text{ and } \det J_X(y, t) = \det J_Y(x, t) = 1$$

for all $t \geq 0$ and $x, y \in \mathbb{R}^n$.

In the situation of several obstacles moving with time dependent velocity we make the following assumption:

- (A2) Let $\mathcal{O}_1, \dots, \mathcal{O}_m \subset \mathbb{R}^n$, $n \geq 2$, be compact sets with boundaries $\Gamma_1, \dots, \Gamma_m$ of class $C^{1,1}$. We denote by $\Omega := \mathbb{R}^n \setminus \bigcup_{i=1}^m \mathcal{O}_i$ the exterior domain. For time-dependent matrices

$$M_i \in C^\infty([0, T]; \mathbb{R}^{n \times n}),$$

with $\text{tr } M_i(t) = 0$ for all $t \in [0, T]$, $i \in \{1, \dots, m\}$, we define the time-dependent sets $\Omega(t)$, $\Gamma_i(t)$ and $\mathcal{O}_i(t)$ for $t \in [0, T]$ by aid of the unique solution $G_{(i)}(t) : \overline{\mathcal{O}_i} \rightarrow \overline{\mathcal{O}_i(t)}$ of the following ODE:

$$(2.5) \quad \begin{cases} G'_{(i)}(t)x &= M_i(t)G_{(i)}(t)x, & x \in \overline{\mathcal{O}_i}, & t \in (0, T), \\ G_{(i)}(0)x &= x, & x \in \overline{\mathcal{O}_i}. \end{cases}$$

Note that $G_{(i)}(t) : \overline{\mathcal{O}_i} \rightarrow \overline{\mathcal{O}_i(t)}$ is an isomorphism, as

$$\begin{cases} [G'_{(i)}(t)x]_k &= M_i(t)[G_{(i)}(t)x]_k, & x \in \overline{\mathcal{O}_i}, & t \in (0, T), \\ [G_{(i)}(0)x]_k &= x_k, & x \in \overline{\mathcal{O}_i}, \end{cases}$$

with $1 \leq k \leq n$, $i \in \{1, \dots, m\}$, has a unique fundamental system of linear independent solutions $G_{(i)}(t) = (G_{(i)}^1(t), \dots, G_{(i)}^n(t))$, due to the non-vanishing Wronski-determinant at zero. In particular, for the case of constant matrices $M_i \in \mathbb{R}^{n \times n}$, we are left with $G_{(i)}(t) = e^{tM_i}$ and inverse $G_{(i)}^{-1}(t) = e^{-tM_i}$.

As the obstacles shall not collide, we require

$$\text{dist} \left(\overline{\mathcal{O}_i(t)}, \overline{\mathcal{O}_j(t)} \right) > 0, \quad i \neq j, \quad t \in [0, T].$$

We now choose open sets $B_{1_i}, B_{2_i} \subset \mathbb{R}^n$, such that $\overline{\mathcal{O}_i} \subset B_{1_i} \subset \overline{B_{1_i}} \subset B_{2_i}$ and set

$$B_{k_i}(t) := \{y = G_{(i)}(t)x, x \in B_{k_i}\}, \quad t \in [0, T], k = 1, 2, i \in \{1, \dots, m\}.$$

Then

$$\overline{\mathcal{O}_i(t)} \subset B_{1_i}(t) \subset \overline{B_{1_i}(t)} \subset B_{2_i}(t)$$

for $t \in [0, T]$. Moreover, we demand the sets B_{2_i} to be so small that the sets $B_{2_i}(t)$ are pairwise disjoint for $t \in [0, T]$. Next, we introduce a time-dependent cut-off function $\eta \in C^\infty(\mathbb{R}^n \times [0, T])$, $0 \leq \eta \leq 1$, such that

$$\eta(y, t) := \begin{cases} 1 & \text{on } \bigcup_{i=1}^m B_{1_i}(t), \\ 0 & \text{on } \mathbb{R}^n \setminus \bigcup_{i=1}^m B_{2_i}(t), \end{cases} \quad t \in [0, T].$$

We define the sets

$$K_i(t) := (\text{supp } \nabla_y \eta(t, \cdot)) \cap \overline{B_{2_i}(t)}, \quad t \in [0, T], i \in \{1, \dots, m\},$$

and the time dependent vector fields $b, b^* : \mathbb{R}^n \times [0, T] \rightarrow \mathbb{R}^n$ by

$$\begin{aligned} b(y, t) &:= \eta(y, t) \sum_{i=1}^m M_i(t)y - \sum_{i=1}^m B_{K_i(t)}((\nabla_y \eta)(\cdot, t)M_i(t)\cdot)(y), \\ b^*(y, t) &:= \eta(y, t) \sum_{i=1}^m G_i^{-1}(t)M_i(t)G_i(t)y \\ &\quad - \sum_{i=1}^m B_{K_i(t)}((\nabla_y \eta)(\cdot, t)G_i^{-1}(t)M_i(t)G_i(t)\cdot)(y). \end{aligned}$$

Note, that due to $\text{tr } M_i(t) = 0$, $t \in [0, T]$, $i = 1, \dots, m$, and properties of the Bogovskiĭ operator, the vector field b is solenoidal for $t \in [0, T]$ on all of $\mathbb{R}^n$. Further, $b(y, t) = M_i(t)y$ for $y \in \overline{\mathcal{O}_i(t)}$, $t \in [0, T]$, $i \in \{1, \dots, m\}$. Note that for fixed $t^* \in [0, T]$

$$b(y, t^*) = \eta(y, t^*) \sum_{i=1}^m M_i(t^*)y - \sum_{i=1}^m B_{K_i^*}((\nabla_y \eta)(\cdot, t^*)M_i(t^*)\cdot)(y).$$

Since η is smooth in the first variable, the mapping properties of the Bogovskiĭ operator imply that $b(\cdot, t^*) \in C_c^\infty(\mathbb{R}^n)$. Now, freeze $y^* \in \mathbb{R}^n$ and consider

$$b(y^*, t) = \eta(y^*, t) \sum_{i=1}^m M_i(t)y^* - \sum_{i=1}^m B_{K_i^*}((\nabla_y \eta)(\cdot, t)M_i(t)\cdot)(y^*).$$

The smoothness of the cut-off function η and the matrices $M_i(\cdot)$ is inherited by b , and we thus see that

$$b \in C_{c,\sigma}^\infty(\mathbb{R}^n \times [0, T]; \mathbb{R}^n).$$

In particular, b is uniformly Lipschitz continuous with respect to the first variable and is bounded on $[0, T] \times \mathbb{R}^n$.

Thus, the ordinary differential equation

$$(2.6) \quad \begin{cases} \partial_t X(y, t) &= b(X(y, t), t), \quad t > 0, \quad y \in \mathbb{R}^n, \\ X(y, 0) &= y, \quad y \in \mathbb{R}^n, \end{cases}$$

admits a unique solution by the Picard-Lindelöf theorem. Moreover,

$$X \in C^\infty(\mathbb{R}^n \times \mathbb{R}_+).$$

As above, let $Y(\cdot, t)$ be the inverse of $X(\cdot, t)$. Then Y satisfies

$$(2.7) \quad \begin{cases} \partial_t Y(x, t) &= -(J_X)^{-1}(Y(x, t), t)b(x, t), & t > 0, & x \in \mathbb{R}^n, \\ Y(x, 0) &= x, & & x \in \mathbb{R}^n, \end{cases}$$

and (2.4). Again, only the restrictions of X, Y to $[0, T]$ will be relevant in the sequel.

We set

$$\begin{aligned} U(y, t) &:= J_Y(X(y, t), t)v(X(y, t), t), & y \in \Omega, & t \in [0, T], \\ \pi(y, t) &:= q(X(y, t), t), & y \in \Omega, & t \in [0, T]. \end{aligned}$$

Then (similar to [IW77], [CT06, Prop.3.5]), a function

$$v \in L^p(0, T; W^{2,q}(\Omega(\cdot))) \cap W^{1,p}(0, T; L^q(\Omega(\cdot))), \quad q \in L^p(0, T; \hat{W}^{1,q}(\Omega(\cdot)))$$

is a solution to (1.1) if and only if

$$U \in L^p(0, T; W^{2,q}(\Omega)) \cap W^{1,p}(0, T; L^q(\Omega)), \quad \pi \in L^p(0, T; \hat{W}^{1,q}(\Omega))$$

and (U, π) satisfies the following set of equations

$$(2.8) \quad \begin{cases} \partial_t U + (\mathcal{M} - \mathcal{L})U &= f - \mathcal{N}U - \mathcal{G}\pi, & \text{in } \Omega \times (0, T), \\ \operatorname{div} U &= 0, & \text{in } \Omega \times (0, T), \\ U &= G_i^{-1}(t)M_i(t)G_i(t)y, & \text{on } \Gamma_i \times (0, T), \quad i = 1, \dots, m, \\ U(0) &= v_0, & \text{in } \Omega. \end{cases}$$

Here

$$\begin{aligned} (\mathcal{L}U)_i &:= \sum_{j,k=1}^n \partial_j(g^{jk}\partial_k U_i) + 2 \sum_{j,k,l=1}^n g^{kl}\Gamma_{jk}^i \partial_l U_j \\ &\quad + \sum_{j,k,l=1}^n \left(\partial_k(g^{kl}\Gamma_{jl}^i) + \sum_{m=1}^n g^{kl}\Gamma_{jl}^m \Gamma_{km}^i \right) U_j, \\ (\mathcal{N}U)_i &:= \sum_{j=1}^n U_j \partial_j U_i + \sum_{j,k=1}^n \Gamma_{jk}^i U_j U_k, \\ (\mathcal{M}U)_i &:= \sum_{j=1}^n \dot{Y}_j \partial_j U_i + \sum_{j,k=1}^n \left(\Gamma_{jk}^i \dot{Y}_k + (\partial_k Y_i)(\partial_j \dot{X}_k) \right) U_j, \\ (\mathcal{G}\pi)_i &:= \sum_{j=1}^n g^{ij} \partial_j \pi, \end{aligned}$$

with the metric contravariant tensor

$$g^{ij} = \sum_{k=1}^n (\partial_k Y_i)(\partial_k Y_j),$$

the metric covariant tensor

$$g_{ij} = \sum_{k=1}^n (\partial_i X_k)(\partial_j X_k)$$

and Christoffel's symbol

$$\Gamma_{ij}^k = 1/2 \sum_{l=1}^n g^{kl} (\partial_j g_{il} + \partial_i g_{jl} - \partial_l g_{ij}).$$

Note that $\mathcal{L}$ is the transformed Stokes operator, while $\mathcal{M}$ arises from transforming the time derivative. The non-linearity $\mathcal{N}$ and modified gradient $\mathcal{G}$ correspond to $(v \cdot \nabla)v$ and ∇ , respectively.

Setting $u(y, t) := U(y, t) - b(y, t)$, we see that

$$U \in L^p(0, T; W^{2,q}(\Omega)) \cap W^{1,p}(0, T; L^q(\Omega)), \quad \pi \in L^p(0, T; \hat{W}^{1,q}(\Omega)),$$

is a solution of (2.8) if and only if

$$u \in L^p(0, T; W^{2,q}(\Omega)) \cap W^{1,p}(0, T; L^q(\Omega)), \quad \pi \in L^p(0, T; \hat{W}^{1,q}(\Omega)),$$

and (u, π) solves

$$(2.9) \quad \begin{cases} \partial_t u - \Delta u + \nabla \pi &= F - \mathcal{N}u - \mathcal{B}u + (\mathcal{L} - \Delta)u \\ &\quad - \mathcal{M}u + (\nabla - \mathcal{G})\pi & \text{in } \Omega \times (0, T), \\ \operatorname{div} u &= 0 & \text{in } \Omega \times (0, T), \\ u &= 0 & \text{on } \Gamma_i \times (0, T), \quad i = 1, \dots, m, \\ u(0) &= v_0 - b(0) & \text{in } \Omega, \end{cases}$$

with

$$(\mathcal{B}u)_i = \sum_{j=1}^n (u_j \partial_j b_i^* + b_j^* \partial_j u_i) + 2 \sum_{j,k=1}^n \Gamma_{jk}^i b_k^* u_j, \quad F = f - \partial_t b^* - (\mathcal{M} - \mathcal{L})b^* - \mathcal{N}b^*.$$

In the sequel, maximal L^p -regularity of the Stokes operator in $L_\sigma^q(\Omega)$ plays an important role. More precisely, for $1 < q < \infty$, we define the Stokes operator A_q in $L_\sigma^q(\Omega)$ by

$$\begin{cases} A_q u &:= P_q \Delta u, \\ D(A_q) &:= W^{2,q}(\Omega) \cap W_0^{1,q}(\Omega) \cap L_\sigma^q(\Omega). \end{cases}$$

As usual, P_q denotes the Helmholtz projection on $L^q(\Omega)$.

Let $T_0 > 0$, $1 < p < \infty$, $T \in (0, T_0)$, $f \in L^p(0, T; L_\sigma^q(\Omega))$ and $u_0 \in (L_\sigma^q(\Omega), D(A_q))_{1-\frac{1}{p}, p}$. Then it follows from a classical result of Solonnikov [Sol77] (also see [Gig81], [Frö02] or [GHHSS06]) that there exists a unique solution

$$u \in X_{p,q}^T := W^{1,p}(0, T; L_\sigma^q(\Omega)) \cap L^p(0, T; W^{2,q}(\Omega) \cap W_0^{1,q}(\Omega) \cap L_\sigma^q(\Omega))$$

of the inhomogeneous Stokes problem

$$\begin{cases} u'(t) - A_q u(t) &= f(t), & t \in (0, T), \\ u(0) &= u_0. \end{cases}$$

Moreover, there exists $C > 0$, independent of T , f and u_0 , such that

$$\|u\|_{X_{p,q}^T} \leq C(\|f\|_{L^p(0,T;L^q(\Omega))} + \|u_0\|_{(L_\sigma^q(\Omega), D(A_q))_{1-\frac{1}{p}, p}}).$$

Setting $\nabla \pi := (\operatorname{Id} - P_q)\Delta u$, we see that (u, π) is a solution to

$$\begin{cases} u'(t) - \Delta u(t) - \nabla \pi(t) &= f(t), & t \in (0, T), \\ u(0) &= u_0, \end{cases}$$

satisfying

$$(2.10) \quad \|u\|_{X_{p,q}^T} + \|\pi\|_{Y_{p,q}^T} \leq C(\|f\|_{L^p(0,T;L^q(\Omega))} + \|u_0\|_{(L_\sigma^q(\Omega), D(A_q))_{1-\frac{1}{p}, p}}),$$

where $Y_{p,q}^T := L^p(0, T; \hat{W}^{1,q}(\Omega))$ and $C > 0$ is a constant independent of $T \in (0, T_0)$, f and u_0 .

For the rest of this section, assume that $M = -M^t$, (A1) holds and consider

$$(2.11) \quad \begin{cases} \partial_t v - \Delta v + (v \cdot \nabla)v + \nabla q = 0 & \text{in } \Omega(t), \quad t \in (0, T), \\ \operatorname{div} v = 0 & \text{in } \Omega(t), \quad t \in (0, T), \\ v(x, t) = Mx & \text{on } \Gamma(t), \quad t \in (0, T), \\ v(x, 0) = v_0(x) & \text{in } \Omega(0). \end{cases}$$

It was shown in [GHH06a] that (2.11) admits a unique, local, mild solution w . In order to compare the solutions obtained in [GHH06a] and in Corollary 3.2, we introduce the notion of very weak solutions to equation (2.11). To this end, let $1 < p, q < \infty$. A function $u \in L^p(0, T_0; L^q(\Omega(\cdot)))$ satisfying $\operatorname{div} u(t) = 0$ in $\Omega(t)$ is called a *very weak solution* to (2.11) if

$$\begin{aligned} \langle v_0, \varphi \rangle_{\Omega} &- \int_0^{T_0} \langle u(t), \varphi'(t) + \Delta \varphi(t) + (u(t) \cdot \nabla) \varphi(t) \rangle_{\Omega(t)} dt \\ &+ \int_0^{T_0} \langle Mx, (\nu(t) \cdot \nabla) \varphi(t) \rangle_{\Gamma(t)} dt = 0, \end{aligned}$$

for $\varphi \in D$, where

$$D := \left\{ \varphi \in C^1 \left([0, T_0]; C_c^\infty(\overline{\Omega(\cdot)}) \right) : \varphi(T_0) = 0, \operatorname{div} \varphi(t) = 0, \right. \\ \left. \varphi(t)|_{\Gamma(t)} = 0 \text{ for all } t \in [0, T_0] \right\}$$

and $\nu(t)$ is the outer normal. We say that two very weak solutions

$$u, v \in L^p(0, T_0; L^q(\Omega(\cdot)))$$

to (2.11) with initial value $v_0 \in L^q(\Omega)$ coincide in the *very weak sense* if there exists $T \in (0, T_0)$ such that $u(t) = v(t)$ for a.a. $t \in (0, T)$.

In [GHH06a], problem (2.11) is transformed into the equation

$$(2.12) \quad \begin{cases} \hat{u}' - A_{\Omega, b} \hat{u} + (\hat{u} \cdot \nabla) \hat{u} = F_2, & t \in (0, T), \\ \hat{u}(0) = v_0 - b, \end{cases}$$

where $\hat{u}(x, t) := e^{-tM} v(e^{tM} x, t) - b(x)$. Here,

$$A_{\Omega, b} \hat{u} := P_q(\Delta \hat{u} + Mx \cdot \nabla \hat{u} - M\hat{u} - b \cdot \nabla \hat{u} - \hat{u} \cdot \nabla b)$$

with $D(A_{\Omega, b}) := \left\{ \hat{u} \in W^{2, q}(\Omega) \cap W_0^{1, q}(\Omega) \cap L_\sigma^p(\Omega) : Mx \cdot \nabla \hat{u} \in L^q(\Omega) \right\}$ and $F_2 := \Delta b + Mx \cdot \nabla b - Mb - b \cdot \nabla b$. It is proved that, for $v_0 - b \in L_\sigma^r(\Omega)$ with $r \geq n$, there exists a *mild solution* $\hat{u}$ to (2.12), i.e. $\hat{u} \in C([0, T]; L_\sigma^r(\Omega))$ satisfies the integral equation

$$(2.13) \quad \hat{u}(t) = T_{\Omega, b}(v_0 - b) - \int_0^t T_{\Omega, b}(t-s) P_q(\hat{u} \cdot \nabla \hat{u})(s) ds + \int_0^t T_{\Omega, b} P_q F_2(s) ds, \quad t \in (0, T).$$

Here, $(T_{\Omega, b}(t))_{t \geq 0} := (e^{tA_{\Omega, b}})_{t \geq 0}$ is the semigroup generated by $(A_{\Omega, b}, D(A_{\Omega, b}))$ in $L_\sigma^r(\Omega)$. Moreover, this solution $\hat{u}$ satisfies:

$$(2.14) \quad t \mapsto t^{\frac{n}{2}(\frac{1}{r} - \frac{1}{q})} \hat{u}(\cdot) \in C([0, T]; L_\sigma^q(\Omega)),$$

$$(2.15) \quad t \mapsto t^{\frac{n}{2}(\frac{1}{r} - \frac{1}{q}) + \frac{1}{2}} \nabla \hat{u}(\cdot) \in C([0, T]; L^q(\Omega)).$$

We show in Section 4 that w given by

$$(2.16) \quad w(x, t) := e^{tM} \hat{u}(e^{-tM} x, t) + e^{tM} b(e^{-tM} x),$$

with $\hat{u}$ given by the variation-of-constants formula, (2.13) is a very weak solution to (2.11) for suitable choices of p, q and n . We are in the position to state our main results.

3. MAIN RESULTS

Our existence and uniqueness result for equation (1.1) reads as follows.

Theorem 3.1. *Assume (A2) and let $p, q \in (1, \infty)$ such that $\frac{n}{2q} + \frac{1}{p} \leq \frac{3}{2}$. Assume that*

- a) $f \in L^p(0, T; L_\sigma^q(\Omega(\cdot)))$,
- b) $v_0 - b(\cdot, 0) \in (L_\sigma^q(\Omega), W_0^{1,q}(\Omega) \cap W^{2,q}(\Omega) \cap L_\sigma^q(\Omega))_{1-1/p, p}$.

Then there exists a $T > 0$ such that the problem (1.1) admits a unique strong solution

$$v \in L^p(0, T; W^{2,q}(\Omega(\cdot))) \cap W^{1,p}(0, T; L_\sigma^q(\Omega(\cdot))), \quad q \in L^p(0, T; \hat{W}^{1,q}(\Omega(\cdot))).$$

Moreover, one may choose $T > 0$ such that either $T = +\infty$ or the function

$$t \mapsto \|v(t)\|_{(L_\sigma^q(\Omega(t)), W_0^{1,q}(\Omega(t)) \cap W^{2,q}(\Omega(t)) \cap L_\sigma^q(\Omega(t)))_{1-1/p, p}}$$

is unbounded on its maximal interval of existence $[0, T)$.

Corollary 3.2. *Assume (A1) and let $p, q \in (1, \infty)$ such that $\frac{n}{2q} + \frac{1}{p} \leq \frac{3}{2}$. Assume that*

- a) $f \in L^p(0, T; L_\sigma^q(\Omega(\cdot)))$,
- b) $v_0 - b \in (L_\sigma^q(\Omega), W_0^{1,q}(\Omega) \cap W^{2,q}(\Omega) \cap L_\sigma^q(\Omega))_{1-1/p, p}$.

Then there exists a $T > 0$ such that the problem (2.11) admits a unique strong solution

$$v \in L^p(0, T; W^{2,q}(\Omega(\cdot))) \cap W^{1,p}(0, T; L_\sigma^q(\Omega(\cdot))), \quad q \in L^p(0, T; \hat{W}^{1,q}(\Omega(\cdot))).$$

Moreover, either $T = +\infty$ or

$$t \mapsto \|v(t)\|_{(L_\sigma^q(\Omega(t)), W_0^{1,q}(\Omega(t)) \cap W^{2,q}(\Omega(t)) \cap L_\sigma^q(\Omega(t)))_{1-1/p, p}}$$

is unbounded on its maximal interval of existence $[0, T)$.

The following theorem says that the two solutions v and w coincide in the very weak sense. More precisely, the following holds true.

Theorem 3.3. *Let $1 < p, q < \infty$ such that $\frac{n}{2q} + \frac{1}{p} \leq \frac{3}{2}$. Assume that*

- a) $v_0 - b \in (L_\sigma^q(\Omega), D(A_q))_{1-\frac{1}{p}, p}$,
- b) $v_0 - b \in L_\sigma^r(\Omega)$ for some $r > n$ provided $\frac{n}{2q} + \frac{1}{p} = \frac{3}{2}$.

Then w given by (2.16) and v given in Corollary 3.2 coincide in the very weak sense.

4. PROOF OF THE FIRST MAIN RESULT

Note that in order to prove our main result it suffices to construct a unique solution (u, π) to (2.9). The strategy of the proof is as follows: first, we derive estimates on the coefficients of the operators $\mathcal{N}, \mathcal{B}, \mathcal{L}, \mathcal{M}$ and $\mathcal{G}$ in problem (2.9). Then, a fixed point argument in a suitable closed subspace of $X_{p,q}^T \times Y_{p,q}^T$ yields a unique solution $(u, \pi) \in X_{p,q}^T \times Y_{p,q}^T$ via maximal L^p -regularity of the Stokes operator.

Observe that for a multi-index α and $k \in \mathbb{N}$ there is some constant $K_{|\alpha|,k,T} > 0$ such that

$$\|\partial_y^\alpha \partial_t^k b\|_{L^\infty(\mathbb{R}^n \times [0,T])} \leq K_{|\alpha|,k,T}, \quad |\alpha| + k > 0.$$

The following lemma yields estimates of the transformation mappings X and Y , respectively, that are defined by (2.6) and (2.7). Clearly, the assertions remain true in the case of b being independent from the time variable and X, Y defined as in (2.2) and (2.3).

Lemma 4.1. *Let $T_0 > 0$, $k \in \mathbb{N}$ and α be a multi-index satisfying $|\alpha| + k > 0$. Then there exists $C_{|\alpha|,k,T_0} > 0$ such that*

$$\|\partial_y^\alpha \partial_t^k X\|_{L^\infty(\mathbb{R}^n \times [0,T_0])} \leq C_{|\alpha|,k,T_0}.$$

The above estimates remain valid when $X(\cdot, t)$ is replaced by its inverse $Y(\cdot, t)$.

Proof. Let $T_0 > 0$, $k \in \mathbb{N}$ and α be a multi-index satisfying $|\alpha| + k > 0$. By a direct calculation, we see that $X(t, y) = y$ for $y \notin \text{supp } b$. Hence,

$$\|\partial_y^\alpha \partial_t^k X\|_{L^\infty((\text{supp } b)^c \times [0,T_0])} \leq 1.$$

Since $\text{supp } b$ is compact and $X \in C^\infty(\mathbb{R}^n \times \mathbb{R}_+)$, there exists $C_{|\alpha|,k,T_0}$ such that

$$\|\partial_y^\alpha \partial_t^k X\|_{L^\infty((\text{supp } b) \times [0,T_0])} \leq C_{|\alpha|,k,T_0}.$$

□

It follows from the definition of g^{ij} , g_{ij} and Γ_{ij}^k and the previous lemma that all coefficients of $\mathcal{L}, \mathcal{M}, \mathcal{N}, \mathcal{B}$ and $\mathcal{G}$ are smooth and bounded on finite time intervals $[0, T_0]$ for any $T_0 > 0$. Moreover, by the mean value theorem, for $x \in \mathbb{R}^n$ we have $g^{ij}(x, t) - \delta_{ij} = t \partial_\tau g^{ij}(x, \tau)$ for some $\tau \in (0, t)$. Hence, it follows from Lemma 4.1 that

$$(4.1) \quad \|g^{ij} - \delta_{ij}\|_{L^\infty(\mathbb{R}^n \times [0,T_0])} \leq t \|\partial_t g^{ij}\|_{L^\infty(\mathbb{R}^n \times [0,T_0])} \leq Ct, \quad t \in (0, T_0).$$

The following embedding of $X_{p,q}^T$ in $L^k(0, T; W^{s,m}(\Omega))$ is needed to cope with the gradient terms. It mainly relies on the mixed derivative theorem. Precisely, the following lemma holds:

Lemma 4.2. *Let $1 < p, q < \infty$, $T_0 > 0$ and $s = 0$ or $s = 1$. Assume that $k, m \in (1, \infty)$ obey $\frac{2-s}{2} + \frac{n}{2m} - \frac{n}{2q} \geq \frac{1}{p} - \frac{1}{k}$. Then $X_{p,q}^T$ is continuously embedded in $L^k(0, T; W^{s,m}(\Omega))$ for any $T \in (0, T_0)$. Moreover, there exists $C_{T_0} > 0$ such that*

$$\|u\|_{L^k(0,T;W^{s,m}(\Omega))} \leq C_{T_0} \|u\|_{X_{p,q}^T}, \quad T \in (0, T_0), \quad u \in X_{p,q}^T, \quad u(0) = 0.$$

Proof. By the mixed derivative theorem (see [Sob75] or [DHP07]), for $\theta \in (0, 1)$ there exists $C > 0$ such that

$$\|u\|_{H^{\theta,p}(0,T_0;H^{2-2\theta,q}(\Omega))} \leq C \|u\|_{X_{p,q}^{T_0}}, \quad u \in X_{p,q}^{T_0}.$$

It then follows from Sobolev embeddings that

$$\|u\|_{L^k(0,T;W^{s,m}(\Omega))} \leq C_{T_0} \|u\|_{X_{p,q}^{T_0}}, \quad u \in X_{p,q}^{T_0}.$$

Let $S : \{u \in X_{p,q}^T : u(0) = 0\} \rightarrow X_{p,q}^{T_0}$ be defined by

$$Su(t) := \begin{cases} u(t - (T_0 - T)), & t \geq T_0 - T, \\ 0, & 0 \leq t < T_0 - T. \end{cases}$$

Then $\|Su\|_{X_{p,q}^{T_0}} = \|u\|_{X_{p,q}^T}$ and $\|Su\|_{L^k(0,T;W^{s,m}(\Omega))} = \|u\|_{L^k(0,T_0;W^{s,m}(\Omega))}$. We thus obtain

$$\|u\|_{L^k(0,T_0;W^{s,m}(\Omega))} = \|Su\|_{L^k(0,T;W^{s,m}(\Omega))} \leq C_{T_0} \|Su\|_{X_{p,q}^{T_0}} = C_{T_0} \|u\|_{X_{p,q}^T}$$

for all $u \in X_{p,q}^T$ satisfying $u(0) = 0$. $\square$

Next, we prove estimates for the terms on the right-hand side of (2.9).

Lemma 4.3. *Fix $T_0 > 0$ and $u \in X_{p,q}^{T_0}$. Then, for $T \in (0, T_0)$ there exists $C > 0$, independent of T and $K(T) > 0$ with $\lim_{T \rightarrow 0} K(T) = 0$, such that for $(v_1, q_1), (v_2, q_2) \in X_{p,q}^T \times Y_{p,q}^T$ satisfying $v_1(0) = v_2(0) = 0$,*

$$(a) \quad \|\mathcal{N}u\|_{L^p(0,T;L^q(\Omega))} \leq (K(T))^2,$$

$$(b) \quad \|\mathcal{N}(v_1 + u) - \mathcal{N}(v_2 + u)\|_{L^p(0,T;L^q(\Omega))} \leq C(K(T) + \sum_{i=1}^2 \|v_i\|_{X_{p,q}^T}) \|v_1 - v_2\|_{X_{p,q}^T},$$

$$(c) \quad \|\mathcal{B}(v_1 - v_2)\|_{L^p(0,T;L^q(\Omega))} \leq CT^{\frac{1}{3p}} \|v_1 - v_2\|_{X_{p,q}^T},$$

$$(d) \quad \|\mathcal{M}(v_1 - v_2)\|_{L^p(0,T;L^q(\Omega))} \leq CT^{\frac{1}{3p}} \|v_1 - v_2\|_{X_{p,q}^T},$$

$$(e) \quad \|(\mathcal{L} - \Delta)(v_1 - v_2)\|_{L^p(0,T;L^q(\Omega))} \leq C(T + T^{\frac{1}{3p}}) \|v_1 - v_2\|_{X_{p,q}^T},$$

$$(f) \quad \|(\nabla - \mathcal{G})(q_1 - q_2)\|_{L^p(0,T;L^q(\Omega))} \leq CT \|q_1 - q_2\|_{Y_{p,q}^T}.$$

The estimates above are valid even if $v_1(0) \neq 0$ and $v_2(0) \neq 0$. However, in this case, C depends on T as well.

Proof. Set $k = 3p$, $k' = 3p/2$, $m = 3q$, $m' = 3q/2$. By Hölder's inequality, we obtain for the first term of $\mathcal{N}u$

$$\|u \cdot \nabla u\|_{L^p(0,T;L^q(\Omega))} \leq \|u\|_{L^k(0,T;L^m(\Omega))} \|u\|_{L^{k'}(0,T;W^{1,m'}(\Omega))}.$$

We now define

$$K(T) := \|u\|_{L^k(0,T;L^m(\Omega))} + \|u\|_{L^{k'}(0,T;W^{1,m'}(\Omega))}.$$

Then,

$$\|u \cdot \nabla u\|_{L^p(0,T;L^q(\Omega))} \leq (K(T))^2.$$

Since $\lim_{T \rightarrow 0} K(T) = 0$, assertion (a) follows.

Next, set $w_1 = v_1 + u$ and $w_2 = v_2 + u$. By Hölder's inequality, we similarly obtain for the first term of $\mathcal{N}w_1 - \mathcal{N}w_2$

$$\begin{aligned} & \|w_1 \cdot \nabla w_1 - w_2 \cdot \nabla w_2\|_{L^p(0,T;L^q(\Omega))} \\ & \leq \| (v_1 - v_2) \cdot \nabla v_1 + v_2 \cdot (\nabla v_1 - \nabla v_2) \|_{L^p(0,T;L^q(\Omega))} \\ & \quad + \| (v_1 - v_2) \cdot \nabla u + u \cdot (\nabla v_1 - \nabla v_2) \|_{L^p(0,T;L^q(\Omega))} \\ & \leq \left(\|v_1\|_{L^{k'}(0,T;W^{1,m'}(\Omega))} + \|u\|_{L^{k'}(0,T;W^{1,m'}(\Omega))} \right) \|v_1 - v_2\|_{L^k(0,T;L^m(\Omega))} \\ & \quad + (\|v_2\|_{L^k(0,T;L^m(\Omega))} + \|u\|_{L^k(0,T;L^m(\Omega))}) \|v_1 - v_2\|_{L^{k'}(0,T;W^{1,m'}(\Omega))}. \end{aligned}$$

Hence, by Lemma 4.2, there exists $C > 0$, independent of $T \in (0, T_0)$ and v_1, v_2 , such that

$$\|w_1 \cdot \nabla w_1 - w_2 \cdot \nabla w_2\|_{L^p(0,T;L^q(\Omega))} \leq C \left(\|v_1\|_{X_{p,q}^T} + \|v_2\|_{X_{p,q}^T} + K(T) \right) \|v_1 - v_2\|_{X_{p,q}^T}.$$

Since the second term of $\mathcal{N}w_1 - \mathcal{N}w_2$ can be estimated similarly, assertion (b) follows.

Similarly, (c) and (d) follow from the estimates

$$\|\mathcal{B}(v_1 - v_2)\|_{L^p(0,T;L^q(\Omega))} \leq CT^{\frac{1}{k}} \|v_1 - v_2\|_{L^{k'}(0,T;W^{1,q}(\Omega))} \leq CT^{\frac{1}{k}} \|v_1 - v_2\|_{X_{p,q}^T}$$

and

$$\|\mathcal{M}(v_1 - v_2)\|_{L^p(0,T;L^q(\Omega))} \leq CT^{\frac{1}{k}} \|v_1 - v_2\|_{L^{k'}(0,T;W^{1,q}(\Omega))} \leq CT^{\frac{1}{k}} \|v_1 - v_2\|_{X_{p,q}^T},$$

respectively, where $C > 0$ is independent of $T \in (0, T_0)$ and v_1, v_2 .

As all coefficients of $\mathcal{L}$ are smooth we may rewrite it in non-divergence form, and, by (4.1), we obtain

$$\begin{aligned} & \|(\mathcal{L} - \Delta)(v_1 - v_2)\|_{L^p(0,T;L^q(\Omega))} \\ & \leq C \|g^{jk}\|_{L^\infty(\mathbb{R}^n \times (0,T))} \|D^2(v_1 - v_2)\|_{L^p(0,T;L^q(\Omega))} \\ & \quad + CT^{\frac{1}{k}} \|v_1 - v_2\|_{L^p(0,T;W^{1,q}(\Omega))} \\ & \leq C(T + T^{\frac{1}{k}}) \|v_1 - v_2\|_{X_{p,q}^T}, \quad T \in (0, T_0), \quad v_1, v_2 \in X_{p,q}^T, \quad v_1(0) = 0, \quad v_2(0) = 0. \end{aligned}$$

Hence (e) follows. Assertion (f) similarly follows from (4.1). $\square$

Let $T_0 > 0$, $T \in (0, T_0)$ and $p, q \in (1, \infty)$ satisfy $\frac{n}{2q} + \frac{1}{p} \leq \frac{3}{2}$. Next, we introduce the space for the fixed point argument. In order to do this, consider

$$\begin{cases} \partial_t u^* - \Delta u^* + \nabla \pi^* &= \hat{F}(0, 0) & \text{in } \Omega \times (0, T), \\ \operatorname{div} u^* &= 0 & \text{in } \Omega \times (0, T), \\ u^* &= 0 & \text{on } \Gamma \times (0, T), \\ u^*(0) &= u_0 & \text{in } \Omega, \end{cases}$$

where $\hat{F} : X_{p,q}^{T_0} \times Y_{p,q}^{T_0} \rightarrow L^p(0, T_0; L^q(\Omega))$ is defined by

$$\hat{F}(v, q) := F - \mathcal{N}v - \mathcal{B}v + (\mathcal{L} - \Delta)v - \mathcal{M}v + (\nabla - \mathcal{G})q.$$

Note that $\hat{F}$ is well-defined by Lemma 4.3 thanks to $\frac{n}{2q} + \frac{1}{p} \leq \frac{3}{2}$. Hence, by (2.10), there exists a unique solution $(u^*, \pi^*) \in X_{p,q}^T \times Y_{p,q}^T$. We equivalently rewrite (2.9)

in terms of a fixed point problem by

$$(4.2) \quad \begin{cases} \partial_t \tilde{u} - \Delta \tilde{u} + \nabla \tilde{\pi} &= F + \hat{F}(v + u^*, q + \pi^*) & \text{in } \Omega \times (0, T), \\ \operatorname{div} \tilde{u} &= 0 & \text{in } \Omega \times (0, T), \\ \tilde{u} &= 0 & \text{on } \Gamma \times (0, T), \\ \tilde{u}(0) &= 0 & \text{in } \Omega, \end{cases}$$

where we used the notation

$$u := \tilde{u} - u^*, \quad \pi := \tilde{\pi} - \pi^*.$$

In view of the Banach fixed point theorem we define for a given radius $R > 0$ and $T \in (0, T_0)$ the closed set

$$\mathcal{K}_{R,T} := \left\{ (v, q) \in X_{p,q}^T \times Y_{p,q}^T : v(0) = 0 \text{ and } \|v\|_{X_{p,q}^T} + \|q\|_{Y_{p,q}^T} \leq R \right\}$$

and a mapping

$$\Phi_{R,T} : \begin{cases} \mathcal{K}_{R,T} & \rightarrow X_{p,q}^T \times Y_{p,q}^T, \\ (v, q) & \mapsto (\tilde{u}, \tilde{\pi}) \end{cases} \quad \text{such that (4.2) holds.}$$

In order to apply the Banach fixed point theorem to $\Phi_{R,T}$ we have to show that the mapping is well-defined, maps $\mathcal{K}_{R,T}$ into itself and is a contraction.

Note that $\phi_{R,T}$ is well-defined due to Lemma 4.3 and (2.10). The next lemma shows that for suitable choices of $R > 0$ and $T > 0$ the closed set $\mathcal{K}_{R,T}$ is mapped by $\Phi_{R,T}$ into itself.

Lemma 4.4. *There exist $R > 0$ and $T_1 > 0$, such that $\Phi_{R,T} : \mathcal{K}_{R,T} \rightarrow \mathcal{K}_{R,T}$ for all $T \in (0, T_1)$.*

Proof. By (2.10) and Lemma 4.3, we obtain

$$\begin{aligned} \|\Phi_{R,T}(v, q)\|_{X_{p,q}^T \times Y_{p,q}^T} &\leq C \|F + \hat{F}(v + u^*, q + \pi^*)\|_{L^p(0,T;L^q(\Omega))} \\ &\leq C (\|F\|_{L^p(0,T;L^q(\Omega))} + \|\mathcal{N}(v + u^*) - \mathcal{N}u^*\|_{L^p(0,T;L^q(\Omega))} + \|\mathcal{N}u^*\|_{L^p(0,T;L^q(\Omega))} \\ &\quad + \|\mathcal{B}(v + u^*)\|_{L^p(0,T;L^q(\Omega))} + \|(\mathcal{L} - \Delta)(v + u^*)\|_{L^p(0,T;L^q(\Omega))} \\ &\quad + \|\mathcal{M}(v + u^*)\|_{L^p(0,T;L^q(\Omega))} + \|(\nabla - \mathcal{G})(q + \pi^*)\|_{L^p(0,T;L^q(\Omega))}) \\ &\leq C (\|F\|_{L^p(0,T;L^q(\Omega))} + K(T)R + R^2 + (K(T))^2 + \|u^*\|_{X_{p,q}^T} + \|\pi^*\|_{Y_{p,q}^T} \\ &\quad + T^{\frac{1}{3p}}R + TR), \quad (v, q) \in \mathcal{K}_{R,T}. \end{aligned}$$

Since

$$\lim_{T \rightarrow 0} \|F\|_{L^p(0,T;L^q(\Omega))} + K(T) + \|u^*\|_{X_{p,q}^T} + \|\pi^*\|_{Y_{p,q}^T} = 0,$$

we obtain $\|\Phi_{R,T}(v, q)\|_{X_{p,q}^T \times Y_{p,q}^T} \leq R$ provided R and T are small enough. $\square$

Lemma 4.5. *There exists $T_0 > 0$ such that $\Phi_{R,T} : \mathcal{K}_{R,T} \rightarrow \mathcal{K}_{R,T}$ is a contraction for all $T \in (0, T_0)$.*

Proof. Again, by (2.10) and Lemma 4.3, we obtain

$$\begin{aligned}
& \|\Phi_{R,T}(v_1, q_1) - \Phi_{R,T}(v_2, q_2)\|_{X_{p,q}^T \times Y_{p,q}^T} \\
& \leq C \left(\|\mathcal{N}(v_1 + u^*) - \mathcal{N}(v_2 + u^*)\|_{L^p(0,T;L^q(\Omega))} + \|\mathcal{B}(v_1 - v_2)\|_{L^p(0,T;L^q(\Omega))} \right. \\
& \quad + \|(\mathcal{L} - \Delta)(v_1 - v_2)\|_{L^p(0,T;L^q(\Omega))} + \|\mathcal{M}(v_1 - v_2)\|_{L^p(0,T;L^q(\Omega))} \\
& \quad \left. + \|(\nabla - \mathcal{G})(q_1 - q_2)\|_{L^p(0,T;L^q(\Omega))} \right) \\
& \leq C(K(T) + R + T + T^{\frac{1}{3p}}) \|(v_1, q_1) - (v_2, q_2)\|_{X_{p,q}^T \times Y_{p,q}^T}, \\
& \quad (v_1, q_1), (v_2, q_2) \in \mathcal{K}_{R,T}.
\end{aligned}$$

Choosing T and R small enough, we obtain $C(K(T) + R + T + T^{\frac{1}{3p}}) < 1$. $\square$

Proof of Theorem 3.1. The existence of a unique strong solution now follows from Lemma 4.4, Lemma 4.5 and the Banach fixed point theorem. Now, the theorem follows in a standard way from the fact that $T > 0$ is uniform with respect to v_0 , provided

$$\|v_0\|_{(L_\sigma^q(\Omega), W_0^{1,q}(\Omega) \cap W^{2,q}(\Omega) \cap L_\sigma^q(\Omega))_{1-1/p,p}} < C_0,$$

and the continuous embedding

$$X_{p,q}^T \hookrightarrow C\left(0, T; \left(W_0^{1,q}(\Omega) \cap W^{2,q}(\Omega) \cap L_\sigma^q(\Omega), L_\sigma^q(\Omega)\right)_{1-1/p,p}\right).$$

$\square$

5. COMPARISON OF STRONG AND MILD SOLUTIONS

In this section we prove Theorem 3.3. For the notion of *very weak* solutions we refer back to Section 2. In a first step, we show that a mild solution is a very weak solution.

Lemma 5.1. *Let $(v_0 - b) \in L_\sigma^q(\Omega)$ for some $q \geq n$ and denote the mild solution to (2.12) on $[0, T]$ for some $T > 0$ by $\hat{u}$ (for a representation by the variation-of-constants formula see (2.13)). Then w , defined by (2.16), is a very weak solution to (2.11).*

Proof. By (2.14) and (2.15), we obtain

$$\hat{u} \in C([0, T]; L_\sigma^q(\Omega)) \quad \text{and} \quad \nabla \hat{u} \in L^1(0, T; L^q(\Omega)).$$

Choose $\hat{u}_n^1 \in C^1([0, T]; C_{c,\sigma}^\infty(\Omega))$ and $\hat{u}_n^2 \in C^1([0, T]; C_c^\infty(\overline{\Omega}))$ such that

$$(5.1) \quad \lim_{n \rightarrow \infty} \|\hat{u}_n^1 - \hat{u}\|_{C([0,T];L^q(\Omega))} = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \|\nabla \hat{u}_n^2 - \nabla \hat{u}\|_{L^1(0,T;L^q(\Omega))} = 0.$$

Then, by [Paz83, Chapter 4, Theorem 2.9], there exists a solution

$$\hat{v}_n \in L^1(0, T; D(A_{\Omega,b})) \cap W^{1,1}(0, T; L_\sigma^q(\Omega))$$

satisfying

$$\begin{cases} \hat{v}_n'(t) - A_{\Omega,b}\hat{v}_n(t) + P_q((\hat{u}_n^1 \cdot \nabla \hat{u}_n^2)(t)) &= P_q F_2(t), & t \in (0, T), \\ \hat{v}_n(0) &= \hat{u}_n^1(0). \end{cases}$$

By the representation of $\hat{v}_n$ via the variation-of-constants formula, we obtain

$$(5.2) \quad \lim_{n \rightarrow \infty} \|\hat{v}_n - \hat{u}\|_{L^1(0,T;L^{\frac{q}{2}}(K))} = 0$$

for any compact $K \subset \overline{\Omega}$. Setting

$$\begin{aligned} \nabla \hat{\pi}_n &= (\text{Id} - P_q)[F_2(t) + \hat{u}_n^1 \cdot \nabla \hat{u}_n^2(t) - \Delta \hat{v}_n(t) - Mx \cdot \nabla \hat{v}_n(t) \\ &\quad + M\hat{v}_n(t) + b \cdot \nabla \hat{v}_n(t) + \hat{v}_n(t) \cdot \nabla b] \end{aligned}$$

for $t \in (0, T)$, we see that $(\hat{v}_n, \hat{\pi}_n)$ solves

$$\left\{ \begin{array}{ll} \hat{v}_n' - \Delta \hat{v}_n - Mx \cdot \nabla \hat{v}_n + M\hat{v}_n \\ \quad + b \cdot \nabla \hat{v}_n + \hat{v}_n \cdot \nabla b + \hat{u}_n^1 \cdot \nabla \hat{u}_n^2 + \nabla \hat{\pi}_n &= F_2 \quad \text{in } \Omega, \quad t \in (0, T), \\ \text{div } \hat{v}_n &= 0 \quad \text{in } \Omega, \quad t \in (0, T), \\ \hat{v}_n &= 0 \quad \text{on } \Gamma, \quad t \in (0, T), \\ \hat{v}_n(0) &= \hat{u}_n^1(0) \quad \text{in } \Omega. \end{array} \right.$$

Hence, $(v_n, \pi_n) := (e^{tM} \hat{v}_n(e^{-tM}x, t) + e^{tM}b(e^{-tM}x), \hat{\pi}(e^{-tM}x, t))$ is a solution to

$$\left\{ \begin{array}{ll} \partial_t v_n - \Delta v_n + u_n^1 \cdot \nabla u_n^2 + (b \cdot \nabla)(v_n - u_n^2) \\ \quad + ((v_n - u_n^1) \cdot \nabla)b + \nabla \pi_n &= 0 \quad \text{in } \Omega(t), \quad t \in (0, T), \\ \text{div } v_n &= 0 \quad \text{in } \Omega(t), \quad t \in (0, T), \\ v_n &= Mx \quad \text{on } \Gamma(t), \quad t \in (0, T), \\ v_n(0) &= u_n^1(0) \quad \text{in } \Omega, \end{array} \right.$$

where

$$u_n^1 := e^{tM} \hat{u}_n^1(e^{-tM}x, t) + e^{tM}b(e^{-tM}x)$$

and

$$u_n^2 := e^{tM} \hat{u}_n^2(e^{-tM}x, t) + e^{tM}b(e^{-tM}x).$$

Moreover,

$$(v_n, \pi_n) \in (L^1(0, T; W^{2,q}(\Omega(\cdot))) \cap W^{1,1}(0, T; L^q(\Omega(\cdot)))) \times L^1(0, T; \dot{W}^{1,q}(\Omega(\cdot)))$$

thanks to $\hat{v}_n \in L^1(0, T; D(A_{\Omega,b})) \cap W^{1,1}(0, T; L^q(\Omega))$. Therefore, integration by parts yields

$$\begin{aligned} \langle v_n(0), \varphi \rangle_{\Omega} &- \int_0^T \langle v_n(t), \varphi'(t) + \Delta \varphi(t) \rangle + \langle u_n^2(t), (u_n^1(t) \cdot \nabla) \varphi(t) \rangle_{\Omega(t)} dt \\ &+ \int_0^T \langle (b \cdot \nabla)(v_n - u_n^2) + ((v_n - u_n^1) \cdot \nabla)b, \varphi \rangle_{\Omega(t)} dt \\ &+ \int_0^T \langle Mx, \nu(t) \cdot \nabla \varphi(t) \rangle_{\Gamma(t)} dt = 0, \quad \varphi \in D. \end{aligned}$$

Letting $n \rightarrow \infty$, it follows from (5.1) and (5.2) that u is a very weak solution. $\square$

In a second step, the following lemma shows uniqueness of the very weak solution for suitable values of p and q . Especially, we will conclude coincidence of the mild and strong notion of solutions, as stated in Theorem 3.3.

Lemma 5.2. *Let $1 < p, q < \infty$ be such that $\frac{n}{2q} + \frac{1}{p} \leq \frac{1}{2}$ and let $v_1, v_2 \in L^p(0, T_0; L^q(\Omega(\cdot)))$ be two very weak solutions to (2.11) for some $T_0 > 0$ with initial value $v_0 \in L^q(\Omega)$. Assume that $v_1 - v_2 \in L^p(0, T_0; L^q_\sigma(\Omega(\cdot)))$. Then there exists $T \in (0, T_0)$ such that $v_1(t) = v_2(t)$ for a.e. $t \in (0, T)$.*

Proof. For the time being, let us assume that for some $T \in (0, T_0)$ and all $f \in L^{p'}(0, T; L^{q'}(\Omega(\cdot)))$, $\frac{1}{p} + \frac{1}{p'} = 1$, $\frac{1}{q} + \frac{1}{q'} = 1$, there exists a solution

$$(\varphi, \pi) \in D_{\text{ext}} := \left(W^{1,p'}(0, T; L^{q'}_\sigma(\Omega(\cdot))) \cap L^{p'}(0, T; W^{2,q'}(\Omega(\cdot))) \right) \\ \times L^{p'}(0, T; \hat{W}^{1,q'}(\Omega(\cdot)))$$

to the dual backward problem

$$(5.3) \quad \begin{cases} -\partial_t \varphi - \Delta \varphi + \nabla \pi - (v_1 + v_2) \cdot \nabla \varphi &= f & \text{in } \Omega(t), \quad t \in (0, T), \\ \operatorname{div} \varphi &= 0 & \text{in } \Omega(t), \quad t \in (0, T), \\ \varphi &= 0 & \text{on } \Gamma(t), \quad t \in (0, T), \\ \varphi(T) &= 0 & \text{in } \Omega. \end{cases}$$

Then, we obtain for all $f \in L^{p'}(0, T; L^{q'}(\Omega(\cdot)))$

$$\begin{aligned} & \int_0^T \langle v_1(t) - v_2(t), f(t) \rangle_{\Omega(t)} dt \\ &= \int_0^T \langle v_1(t) - v_2(t), (-\partial_t \varphi - \Delta \varphi + \nabla \pi - (v_1 + v_2) \cdot \nabla \varphi)(t) \rangle_{\Omega(t)} dt \\ &= \int_0^T \langle v_1(t) - v_2(t), (-\partial_t \varphi - \Delta \varphi - (v_1 + v_2) \cdot \nabla \varphi)(t) \rangle_{\Omega(t)} dt = 0. \end{aligned}$$

Here, we have used that φ can be approximated by functions in D . This implies $v_1 = v_2$ in $(0, T)$.

It thus remains to show that for $f \in L^{p'}(0, T; L^{q'}(\Omega(\cdot)))$, there exists a solution $(\varphi, \pi) \in D_{\text{ext}}$ to (5.3). In order to do so, we first consider the forward problem

$$(5.4) \quad \begin{cases} \partial_t \varphi - \Delta \varphi + \nabla \pi - (v_1^T + v_2^T) \cdot \nabla \varphi &= f^T & \text{in } \Omega(t), \quad t \in (0, T), \\ \operatorname{div} \varphi &= 0 & \text{in } \Omega(t), \quad t \in (0, T), \\ \varphi &= 0 & \text{on } \Gamma(t), \quad t \in (0, T), \\ \varphi(0) &= 0 & \text{in } \Omega, \end{cases}$$

where $v_1^T(t) = v_1(T - t)$, $v_2^T(t) = v_2(T - t)$ and $f^T(t) = f(T - t)$ for $t \in (0, T)$. Note first that by a scaling argument, we may assume that $\|f^T\|_{L^{p'}(0, T; L^{q'}(\Omega(\cdot)))}$ is arbitrarily small. Then, similar to the proof of Theorem 3.1, it follows that there exists $T > 0$, independent of f^T , and a solution $(\varphi, \pi) \in D_{\text{ext}}$ to (5.3). Indeed, we have an additional term coming from f^T , which is no problem since f^T is arbitrarily small. Moreover, the non-linear term has to be replaced by a term coming from $(v_1^T + v_2^T) \cdot \nabla \varphi$. For the convenience of the reader, we will give the estimates for $(\tilde{v}_1^T + \tilde{v}_2^T) \cdot \nabla \tilde{\varphi}$. We choose $r, s \in (1, \infty)$ such that $\frac{1}{p'} = \frac{1}{p} + \frac{1}{r}$ and $\frac{1}{q'} = \frac{1}{q} + \frac{1}{s}$.

Then, it follows from Hölder's inequality and Lemma 4.2 that

$$\begin{aligned} \|(\tilde{v}_1^T - \tilde{v}_2^T) \cdot \nabla \tilde{\varphi}\|_{L^{p'}(0,T;L^{q'}(\Omega))} &\leq \|\tilde{v}_1^T - \tilde{v}_2^T\|_{L^p(0,T;L^q(\Omega))} \|\nabla \tilde{\varphi}\|_{L^r(0,T;L^s(\Omega))} \\ &\leq C \|\tilde{v}_1^T - \tilde{v}_2^T\|_{L^p(0,T;L^q(\Omega))} \|\tilde{\varphi}\|_{X_{p',q'}^T}. \end{aligned}$$

Since $\|\tilde{v}_1^T - \tilde{v}_2^T\|_{L^p(0,T;L^q(\Omega))} \rightarrow 0$ for $T \rightarrow 0$, we get a solution (φ^T, π^T) to (5.4) for some $T > 0$. Finally, (φ, π) , where $\varphi(t) := \varphi^T(T-t)$ and $\pi(t) := \pi^T(T-t)$ for $t \in (0, T)$, is a solution to (5.3). $\square$

We finally prove Theorem 3.3.

Proof of Theorem 3.3. Let us first assume that $\frac{n}{2q} + \frac{1}{p} < \frac{3}{2}$. Then, by Sobolev embeddings,

$$(L_\sigma^q(\Omega), D(A_q))_{1-\frac{1}{p}, p} \hookrightarrow L_\sigma^{\tilde{r}}(\Omega)$$

for some $\tilde{r} > n$. Hence, $v, w \in C([0, T_0]; L^{\tilde{r}}(\Omega))$ for some $T_0 > 0$. In particular, v, w satisfy the assumption of Lemma 5.2. Hence, the assertion follows from iteration in this case.

Let us now assume that $\frac{n}{2q} + \frac{1}{p} = \frac{3}{2}$ and $v_0 - b \in L_\sigma^r(\Omega)$ for some $r > n$. By Sobolev embeddings, we have

$$v \in L^s(0, T; L^{\tilde{s}}(\Omega))$$

for some $\tilde{s} \in (n, r)$ and $1 < s < \infty$ satisfying $\frac{n}{2\tilde{s}} + \frac{1}{s} \leq \frac{1}{2}$. Moreover, since $v_0 - b \in L_\sigma^{\tilde{s}}(\Omega)$, we have $v \in C([0, T]; L^{\tilde{s}}(\Omega))$. Hence, in this case the assertion follows similar to the above. $\square$

REFERENCES

- [Ama00] H. Amann, On the strong solvability of the Navier-Stokes equations, *J. Math. Fluid Mech.* **2** (2000), 16–98. MR1755865 (2002b:76028)
- [Arn92] V. Arnol'd, *Ordinary Differential Equations*, Springer, Berlin, 1992. MR1162307 (93b:34001)
- [Bal77] J. M. Ball, Strongly continuous semigroups, weak solutions, and the variation of constants formula, *Proc. Amer. Math. Soc.* **63** (1977), 370–373. MR0442748 (56:1128)
- [BMN99] A. Babin, A. Mahalov and B. Nicolaenko, Global regularity of 3D rotating Navier-Stokes equations for resonant domains, *Indiana Univ. Math. J.* **48** (1999), 1133–1176. MR1736966 (2001b:35241)
- [Bog79] M. E. Bogovskiĭ, Solution of the first boundary value problem for an equation of continuity of an incompressible medium, *Dokl. Akad. Nauk SSSR* **248** (1979), 1037–1040. MR0553920 (82b:35135)
- [CT06] P. Cumsille and M. Tucsnak, Wellposedness for the Navier-Stokes flow in the exterior of a rotating obstacle, *Math. Methods Appl. Sci.* **29** (2006), 595–623. MR2205973 (2007b:35250)
- [CM97] Z. Chen and T. Miyakawa, Decay properties of weak solutions to a perturbed Navier-Stokes system in $\mathbb{R}^n$, *Adv. Math. Sci. Appl.* **7** (1997), 741–770. MR1476275 (98k:35147)
- [DHP03] R. Denk, M. Hieber and J. Prüss, $\mathcal{R}$ -boundedness, *Fourier Multipliers and Problems of Elliptic and Parabolic Type*, Mem. Amer. Math. Soc. (2003). MR2006641 (2004i:35002)
- [DHP07] ———, Optimal L^p - L^q -estimates for parabolic boundary value problems with inhomogeneous data, *Math. Z.* **257** (2007), 193–224. MR2318575 (2008f:35166)
- [FJR72] E.B. Fabes, B.F. Jones and N.M. Riviere, The initial value problem for the Navier-Stokes equations with data in L^p , *Arch. Rational Mech. Anal.* **45** (1972), 222–240. MR0316915 (47:5463)
- [Frö02] A. Fröhlich, Maximal regularity for the non-stationary Stokes system in an aperture domain, *J. Evol. Equ.* **4** (2002), 471–493. MR1941038 (2003h:35200)

- [Gal94] G. P. Galdi, *An Introduction to the Mathematical Theory of the Navier-Stokes Equations*. Vol. I, Springer, New York, 1994. MR1284205 (95i:35216a)
- [Gal03] G.P. Galdi, Steady flow of a Navier-Stokes fluid around a rotating obstacle, *J. Elasticity* **71** (2003), 1–31. MR2042672 (2005c:76030)
- [GS05] P. Galdi, A. Silvestre, Strong solutions to the Navier-Stokes equations around a rotating obstacle, *Arch. Rational Mech. Anal.* **176** (2005), 331–350. MR2185661 (2006i:35276)
- [GHH06a] M. Geissert, H. Heck, M. Hieber, L^p -theory of the Navier-Stokes flow past rotating or moving obstacles, *J. Reine Angew. Math.* **596** (2006), 45–62. MR2254804 (2007d:35208)
- [GHH06b] ———, *On the equation $\operatorname{div} u = f$ and the Bogovskii Operator*, Oper. Theory Adv. Appl., 168, Birkhäuser, Basel, 2006. MR2240056 (2007k:35034)
- [GHHSS06] M. Geissert, M. Hess, M. Hieber, C. Schwarz and K. Stavrakidis, Maximal $L^p - L^q$ -estimates for the Stokes equation: a short proof of Solonnikov’s theorem, *J. Math. Fluid Mechanics*, to appear.
- [Gig81] Y. Giga, Analyticity of the semigroup generated by the Stokes operator in L_r spaces, *Math. Z.* **178** (1981), 297–329. MR0635201 (83e:47028)
- [GIMM04] Y. Giga, K. Inui, A. Mahalov, S. Matsui, Navier-Stokes equations in a rotating frame in $\mathbb{R}^3$ with initial data nondecreasing at infinity, *Hokkaido Math. J.* **35** (2006), 321–364. MR2254655 (2007f:35217)
- [GIMMS05] Y. Giga, K. Inui, A. Mahalov, S. Matsui, J. Saal, Rotating Navier-Stokes equations in $\mathbb{R}_+^3$ with initial data nondecreasing at infinity: the Ekman boundary layer problem, *Arch. Rat. Mech. Anal.*, **186** (2007), 177–224. MR2342201
- [HS05] M. Hieber and O. Sawada, The Navier-Stokes equations in R^n with linearly growing initial data, *Arch. Rational Mech. Anal.* **175** (2005), 269–285. MR2118478 (2006k:35216)
- [His99] T. Hishida, An existence theorem for the Navier-Stokes flow in the exterior of a rotating obstacle, *Arch. Ration. Mech. Anal.* **150** (1999), 307–348. MR1741259 (2001b:76024)
- [HS06] T. Hishida, Y. Shibata, Stability of the Navier-Stokes flow past a rotating obstacle, preprint (2006).
- [IW77] A. Inoue and M. Wakimoto, On existence of solutions of the Navier-Stokes equation in a time dependent domain, *J. Fac. Sci. Univ. Tokyo, Sect. IA Math.* **24** (1977), 303–319. MR0481649 (58:1750)
- [Paz83] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer, New York, 1983. MR0710486 (85g:47061)
- [Sob75] P. E. Sobolevskii, Fractional powers of coercively positive sums of operators, *Dokl. Akad. Nauk SSSR* **225** (1975), 1271–1274. MR0482314 (58:2387)
- [Sol77] V. Solonnikov, Estimates for solutions of nonstationary Navier-Stokes equations, *J. Sov. Math.* **8** (1977), 467–529.

FACHBEREICH MATHEMATIK, TECHNISCHE UNIVERSITÄT DARMSTADT, SCHLOSSGARTENSTR. 7,
D-64289 DARMSTADT, GERMANY

E-mail address: dintelmann@mathematik.tu-darmstadt.de

FACHBEREICH MATHEMATIK, TECHNISCHE UNIVERSITÄT DARMSTADT, SCHLOSSGARTENSTR. 7,
D-64289 DARMSTADT, GERMANY

E-mail address: geissert@mathematik.tu-darmstadt.de

FACHBEREICH MATHEMATIK, TECHNISCHE UNIVERSITÄT DARMSTADT, SCHLOSSGARTENSTR. 7,
D-64289 DARMSTADT, GERMANY

E-mail address: hieber@mathematik.tu-darmstadt.de