ON THE COHOMOLOGY GROUPS OF HOLOMORPHIC BANACH BUNDLES

LÁSZLÓ LEMPert

ABSTRACT. We consider a compact complex manifold M and introduce the notion of two holomorphic Banach bundles E, F over M being compact perturbations of one another. Given two such bundles we show that if the cohomology groups $H^q(M, E)$ are finite dimensional, then so are the cohomology groups $H^q(M, F)$; as well as a more precise result in the same spirit.

INTRODUCTION

A basic result in complex geometry is the finiteness theorem of Cartan and Serre [CS], according to which the cohomology groups of a finite rank holomorphic vector bundle $E \rightarrow M$ over a compact base are finite dimensional. (More generally, the same holds for the cohomology groups of coherent sheaves over M .) It is no surprise that holomorphic *Banach* bundles $E \rightarrow M$ over a compact base can easily have infinite dimensional cohomology groups. For example, J. Kim observes that if E is the trivial bundle with fiber a Banach space A , then $H^q(M, E) \approx H^q(M, \mathcal{O}) \otimes A$ (tensor product over $\mathbb{C}$); the special case when $H^q(M, \mathcal{O}) = 0$ is due to Leiterer [K, Li2]. In particular, $\dim H^q(M, E) = \infty$ if $\dim A = \infty$ and $H^q(M, \mathcal{O}) \neq 0$.

However, already in [G], Gohberg was led to a class of holomorphic (first Hilbert, then) Banach bundles, for which finiteness theorems can be proved. The terminology is *bundles of compact type*, meaning their structure group can be reduced to the group of invertible operators of the form $\text{id} + \text{compact}$. Let $F \rightarrow \mathbb{P}_1$ be a holomorphic Banach bundle of compact type over the Riemann sphere. The Riemann–Hilbert type result of [G, GL], coupled with Bungart’s work [B], implies that $F \approx T \oplus V$, where T is trivial and V has finite rank. Hence

$$H^1(\mathbb{P}_1, F) \approx H^1(\mathbb{P}_1, T) \oplus H^1(\mathbb{P}_1, V) \approx H^1(\mathbb{P}_1, V)$$

is finite dimensional by what has been said above. More recently Leiterer in [Li2] considers an arbitrary compact complex manifold M and holomorphic Banach bundles $F \rightarrow M$ of compact type, $V \rightarrow M$ of finite rank. Assuming $H^q(M, V) = 0$ for some $q \geq 0$, and putting a mild restriction on the fiber type of F , he proves $\dim H^q(M, F \otimes V) < \infty$.

Our goal in this paper is to generalize and thereby better understand the phenomenon discovered by Gohberg and Leiterer. Instead of bundles of compact type, we shall work with a bundle and its compact perturbation, by which we mean

Received by the editors May 4, 2007.

2000 *Mathematics Subject Classification*. Primary 32L05, 32L10, 58B15.

This research was partially supported by NSF grants DMS0203072 and 0700281.

©2009 American Mathematical Society
 Reverts to public domain 28 years from publication

the following. Let N be a complex manifold and $E, F \rightarrow N$ holomorphic Banach bundles.

Definition 1.1. (a) A fredomorphism $E \rightarrow F$ is a homomorphism $\varphi: E \rightarrow F$ such that $\varphi|_{E_x}: E_x \rightarrow F_x$ is a Fredholm operator for each $x \in N$.

(b) We say that F is a compact perturbation of E if there are an open cover $\mathfrak{U}$ of N and fredomorphisms $\varphi_U: E|_U \rightarrow F|_U$ such that $\varphi_U - \varphi_V|_{E_x}: E_x \rightarrow F_x$ is compact whenever $U, V \in \mathfrak{U}$ and $x \in U \cap V$.

It is easy to see (cf. Proposition 4.2) that being a compact perturbation is a symmetric and transitive relation—for example, any two finite rank bundles $E, F \rightarrow N$ are compact perturbations of each other, the zero homomorphism $E \rightarrow F$ being a (global) fredomorphism. Further, if F is a holomorphic Banach bundle of compact type whose fibers are isomorphic to a Banach space A , then F is a compact perturbation of the trivial bundle $A \times N \rightarrow N$.

We recall that the sheaf cohomology groups $H^q(N, E)$ of holomorphic Banach bundles are topological vector spaces; see Section 2.

Theorem 1.2. *Let M be a compact complex manifold, E, F holomorphic Banach bundles over M that are compact perturbations of one another, and $q = 0, 1, \dots$. If $H^{q+1}(M, E)$ is Hausdorff (in particular if $\dim H^{q+1}(M, E) < \infty$) and if $\dim H^q(M, E) < \infty$, then $H^q(M, F)$ is also finite dimensional (and Hausdorff).*

The assumption that $H^{q+1}(M, E)$ be Hausdorff is not automatically satisfied, and without it the conclusion would not necessarily hold; see the discussion at the end of the paper.

There should be a sheaf version of this theorem, involving cohesive sheaves from [LP] or Leiterer's Banach coherent analytic Fréchet sheaves from [Li1], but so far we have only been able to prove an extension of Theorem 1.2 to certain very special subsheaves of E, F above.

Compact perturbations can be defined more generally for topological Banach bundles. If M is a compact topological space and $E, F \rightarrow M$ are Banach bundles, any system of fredomorphisms $\varphi_U: E|_U \rightarrow F|_U$, $U \in \mathfrak{U}$ as in Definition 1.1b induces a virtual vector bundle of finite rank over M , i.e., an element of the group $K(M)$. In particular, one can speak of characteristic classes in this situation. Perhaps even in the holomorphic category one should regard compact perturbations as generalizations of holomorphic vector bundles of finite rank; Theorem 1.2 seems to point in this direction.

The finiteness results of Gohberg and Leiterer are special cases of Theorem 1.2. Indeed, as said, a bundle F of compact type is a compact perturbation of a trivial bundle $E = A \times M \rightarrow M$, and $F \otimes V$ is a compact perturbation of $E \otimes V$, when $\text{rk } V < \infty$. Now in [K], J. Kim proves $H^p(M, E \otimes V) \approx H^p(M, V) \otimes A$ as topological vector spaces; in particular all $H^p(M, E \otimes V)$ are Hausdorff. (In general there are many natural ways to topologize the tensor product $Y \otimes A$ of Banach spaces; but there is no ambiguity when $\dim Y = n < \infty$ as above. In this case $Y \otimes A \approx A \oplus \dots \oplus A$, n times.) The Cartan–Serre theorem, at least for bundles over manifolds, is also a special case.

In analysis and geometry finite dimensionality results typically depend on functional analysis: Fredholm theory or Schwartz's theorem on compact perturbations of Fréchet space operators. We shall also help ourselves to these tools, but the present proof will have a nonlinear ingredient as well, Michael's selection theorem

[M], whose application in complex geometry is pretty much the only novelty in this work.

2. HOLOMORPHIC FRÉCHET BUNDLES AND THEIR COHOMOLOGY GROUPS. GENERALITIES

By a Fréchet space we shall mean here a complete, metrizable locally convex topological vector space; a sequence of seminorms defining the topology need not be specified. Similarly, by a Banach space we mean a completely normable locally convex space. A continuous linear map between locally convex spaces will be called a homomorphism. A (locally trivial) holomorphic Fréchet bundle over a finite dimensional complex manifold M is a holomorphic map $\pi: E \rightarrow M$ of a complex Fréchet manifold E on M , with each fiber $E_x = \pi^{-1}(x)$, $x \in M$, endowed with the structure of a complex vector space. It is required that for every $x \in M$ there be a neighborhood $U \subset M$ of x , a complex Fréchet space A , and a biholomorphism (trivialization) $E|U = \pi^{-1}U \rightarrow A \times U$ mapping the fibers E_y linearly on $A \times \{y\} \cong A$, $y \in U$. If A above is a Banach space, we speak of a Banach bundle. For more about these notions, see [Le1, Sections 1,2]. We fix a holomorphic Fréchet bundle $E \rightarrow M$ and in this section we describe basic properties of its sheaf cohomology groups $H^q(M, E)$ as locally convex spaces. In addition to the notation and definitions introduced in this section, the results that we will need later on are Lemma 2.5 and Corollary 2.6.

Let $\varphi_U: E|U \rightarrow A \times U$ be a trivialization. For each compact $K \subset U$ and continuous seminorm p on A we define seminorms

$$\|e\|_{K,p} = \sup_{x \in K} p(\varphi_U e(x)), \quad e \in \Gamma(U, E),$$

on the space $\Gamma(U, E)$ of holomorphic sections. They endow $\Gamma(U, E)$ with the structure of a complete locally convex space, in fact a Fréchet space, since a countable collection of K, p suffices to define the topology. Clearly, this Fréchet space structure is independent of the choice of the trivialization.

If $q = 0, 1, \dots$, and $\mathfrak{U}$ is a countable open cover of M such that $E|U$ is trivial for each $U \in \mathfrak{U}$, then the space of (not necessarily alternating) q -cochains

$$C^q(\mathfrak{U}, E) = \prod_{U_0, \dots, U_q \in \mathfrak{U}} \Gamma\left(\bigcap_{i=0}^q U_i, E\right)$$

is again a Fréchet space; we use the convention that the space of functions (and later, forms) on the empty set is the 0-dimensional space (0). The coboundary operator

$$\delta = \delta_{\mathfrak{U}}^q : C^q(\mathfrak{U}, E) \rightarrow C^{q+1}(\mathfrak{U}, E)$$

is continuous, $Z^q(\mathfrak{U}, E) = \text{Ker} \delta_{\mathfrak{U}}^q$ is a closed subspace, and the cohomology groups

$$H^q(\mathfrak{U}, E) = Z^q(\mathfrak{U}, E) / \delta_{\mathfrak{U}}^{q-1} C^{q-1}(\mathfrak{U}, E)$$

are locally convex vector spaces, complete and semimetrizable, but possibly non-Hausdorff. A refinement map $\mathfrak{V} \rightarrow \mathfrak{U}$ between two covers induces homomorphisms $\rho_{\mathfrak{V}}^{\mathfrak{U}} : H^q(\mathfrak{U}, E) \rightarrow H^q(\mathfrak{V}, E)$. We endow the vector space direct limit $H^q(M, E)$ of the system $\{H^q(\mathfrak{U}, E), \rho_{\mathfrak{V}}^{\mathfrak{U}}\}$ with the finest locally convex topology for which the canonical maps $H^q(\mathfrak{U}, E) \rightarrow H^q(M, E)$ are continuous. (It can be seen from Theorem 2.2 below that this is simply the finest topology which makes the canonical maps continuous.)

A variant of this construction provides better control of the topology obtained. If $\varphi_U: E|U \rightarrow A \times U$ is a trivialization as above and $W \Subset U$ is open, then for each continuous seminorm p on A and $e \in \Gamma(W, E)$ we let $\|e\|_p = \sup_{x \in W} p(\varphi_U e(x)) \leq \infty$ and define the space of bounded holomorphic sections by

$$\Gamma_b(W, E) = \{e \in \Gamma(W, E) : \|e\|_p < \infty \text{ for each } p\}.$$

This space, with the seminorms $\|\cdot\|_p$ is a Fréchet space, whose topology is again independent of the choice of $U \supset W$ and the trivialization φ_U . Obviously, the embedding $\Gamma_b(W, E) \rightarrow \Gamma(W, E)$ is continuous. If $\mathfrak{W}$ is a countable open cover of M consisting of such W , then the space of bounded q -cochains,

$$C_b^q(\mathfrak{W}, E) = \prod_{W_0, \dots, W_q \in \mathfrak{W}} \Gamma_b\left(\bigcap_{i=0}^q W_i, E\right),$$

is a Fréchet space, and the coboundary δ restricts to a homomorphism $C_b^q(\mathfrak{W}, E) \rightarrow C_b^{q+1}(\mathfrak{W}, E)$. Denoting the kernel of $\delta|C_b^q(\mathfrak{W}, E)$ by $Z_b^q(\mathfrak{W}, E) \subset C_b^q(\mathfrak{W}, E)$, the bounded cohomology groups

$$H_b^q(\mathfrak{W}, E) = Z_b^q(\mathfrak{W}, E) / \delta_{\mathfrak{W}}^{q-1} C_b^{q-1}(\mathfrak{W}, E)$$

are complete semimetrizable locally convex spaces. With a cover $\mathfrak{U}$ finer than $\mathfrak{W}$, restriction induces homomorphisms

$$(2.1) \quad H_b^q(\mathfrak{W}, E) \rightarrow H^q(\mathfrak{U}, E).$$

Composing these with the canonical homomorphisms $H^q(\mathfrak{U}, E) \rightarrow H^q(M, E)$ we obtain canonical homomorphisms

$$(2.2) \quad H_b^q(\mathfrak{W}, E) \rightarrow H^q(M, E).$$

Definition 2.1. We say that a countable open cover $\mathfrak{W}$ of M is special (with respect to E) if (a) for every $W \in \mathfrak{W}$ there is a biholomorphism of a neighborhood of $\overline{W}$ into some $\mathbb{C}^n$ that maps W on a bounded, strongly pseudoconvex domain with smooth boundary; also E is trivial on this neighborhood, and (b) the boundaries of $W \in \mathfrak{W}$ are in general position. This latter means that if $k \in \mathbb{N}$ and ρ_i are smooth defining functions of $W_i \in \mathfrak{W}$, $i = 1, \dots, k$, then $d\rho_1, \dots, d\rho_k$ are linearly independent at each point of the set $\{\rho_1 = \dots = \rho_k = 0\}$.

Sard's lemma implies that M has arbitrarily fine special covers.

Theorem 2.2. *If $\mathfrak{W}$ is a special cover of M , then (2.2) is an isomorphism; if, in addition, $\mathfrak{U}$ consists of Stein open sets, then (2.1) is also an isomorphism.*

The proof uses routine sheaf theory plus certain deeper facts of several complex variables. We start by introducing a few more spaces generalizing $\Gamma(U, E)$, $\Gamma_b(W, E)$, $C^q(\mathfrak{U}, E)$, etc. Fix smooth vector fields $\xi_1, \dots, \xi_m$ that span each tangent space $T_x M$. Let $\varphi_U: E|U \rightarrow A \times U$ be a trivialization over an open $U \subset M$. We endow the space $\Omega^{0,r}(U, E)$ of smooth E -valued $(0, r)$ forms with seminorms $\|\cdot\|_{K,p,j}$, one for each compact $K \subset U$, continuous seminorm p on A , and $j = 0, 1, 2, \dots$:

$$(2.3) \quad \|f\|_{K,p,j} = \max_K \max_p \left(\xi_{i_1} \xi_{i_2} \dots \xi_{i_j} (\varphi_U f(\xi_{i_{j+1}}, \dots, \xi_{i_{j+r}})) \right),$$

the outer max being taken over all tuples $1 \leq i_1, \dots, i_{j+r} \leq m$. If $\mathfrak{U}$ is a countable cover of M consisting of such open sets, we define a double complex

$$C^{qr}(\mathfrak{U}) = \prod_{U_0, \dots, U_q \in \mathfrak{U}} \Omega_b^{0,r} \left(\bigcap_{i=0}^q U_i, E \right), \quad q, r \geq 0,$$

of Fréchet spaces. We let

$$\delta = \delta^{qr} : C^{qr}(\mathfrak{U}) \rightarrow C^{q+1,r}(\mathfrak{U}), \quad \bar{\partial} = \bar{\partial}^{qr} : C^{qr}(\mathfrak{U}) \rightarrow C^{q,r+1}(\mathfrak{U})$$

denote the Čech coboundary, resp. the homomorphism obtained by applying $\bar{\partial}$ componentwise. Set

$$Z^{qr}(\mathfrak{U}) = \text{Ker } \bar{\partial}^{qr} \cap \text{Ker } \delta^{qr}.$$

With U as above and $W \Subset U$ open, we also consider the space $\Omega_b^{0,r}(W, E) \subset \Omega^{0,r}(W, E)$ of bounded $(0, r)$ forms f , meaning that for each continuous seminorm p on A , $\|f\|_p = \max \sup_W p(\varphi_U f(\xi_{i_1}, \dots, \xi_{i_r})) < \infty$, where the max is taken over all $1 \leq i_1, \dots, i_r \leq m$. We endow $\Omega_b^{0,r}(W, E)$ with these seminorms $\|\cdot\|_p$ to obtain a metrizable locally convex space, no longer complete. Observe that the embedding $\Omega_b^{0,r}(W, E) \rightarrow \Omega^{0,r}(W, E)$ is not continuous when $r > 0$. Nevertheless, given a countable cover $\mathfrak{W}$ consisting of such W there are the spaces

$$C_b^{qr}(\mathfrak{W}) = \prod_{W_0, \dots, W_q \in \mathfrak{W}} \Omega_b^{0,r} \left(\bigcap_{i=0}^q W_i, E \right) \subset C^{qr}(\mathfrak{W})$$

of bounded cochains, again with a metrizable locally convex topology. Thus δ^{qr} restricts to a homomorphism $C_b^{qr}(\mathfrak{W}) \rightarrow C_b^{q+1,r}(\mathfrak{W})$ (but $\bar{\partial}^{qr}$ is not continuous from $C_b^{qr}(\mathfrak{W})$ to $C_b^{q,r+1}(\mathfrak{W})$). Finally, we let $Z_b^{qr}(\mathfrak{W}) = Z^{qr}(\mathfrak{W}) \cap C_b^{qr}(\mathfrak{W})$ and endow it with the topology inherited from $C_b^{qr}(\mathfrak{W})$.

Lemma 2.3. *If $\mathfrak{W}$ is a special cover of M , $r \geq 1$, and $W_0, \dots, W_q \in \mathfrak{W}$, then there is a homomorphism*

$$(2.4) \quad T = T_{W_0 \dots W_q} : \Omega_b^{0,r} \left(\bigcap_{i=0}^q W_i, E \right) \rightarrow \Omega_b^{0,r-1} \left(\bigcap_{i=0}^q W_i, E \right)$$

such that $\bar{\partial} T f = f$ if $\bar{\partial} f = 0$.

Proof. Because of our hypotheses, we can assume that $M = \mathbb{C}^n$ and E is the trivial bundle $A \times \mathbb{C}^n \rightarrow \mathbb{C}^n$. The key is a result of Range and Siu, who construct, when $A = \mathbb{C}$, homomorphisms T as in (2.4) with the property that $\bar{\partial} T + T \bar{\partial} = \text{id}$; see [RS, (3.9)]. In fact, their operator T is a (locally uniform) limit of integral operators T^δ ; see [RS, (2.8)–(2.9)]. Now the integrals defining T^δ make sense if instead of $\mathbb{C}$ -valued forms we substitute in them A -valued forms, and by the Banach–Hahn theorem they satisfy the same estimates as the scalar-valued operators. It follows that the A -valued integral operators converge to a homomorphism T claimed by the lemma.

A simple consequence is

Theorem 2.4. *If M is Stein and $E = A \times M \rightarrow M$ is a trivial Fréchet bundle, then the Dolbeault cohomology groups $H_{\bar{\partial}}^{0,r}(M, E)$ vanish for $r \geq 1$.*

The only surprising thing about this theorem is that it cannot be generalized much. Vogt in [V] shows that $H_{\bar{\partial}}^{0,1}(M, E)$ need not vanish even for $M = \mathbb{C}$ but with a trivial bundle E with general complete locally convex fibers, and Leiterer explained to me that nontrivial Fréchet bundles over Stein manifolds may have non-vanishing $H_{\bar{\partial}}^{0,1}$. Such Fréchet bundles can be constructed from locally trivial fiber bundles with Stein base and fibers whose total space is not Stein (“counterexamples to Serre’s conjecture”; see [Sk]).

Proof. First assume that M is an open subset of some $\mathbb{C}^k$ and exhaust it by holomorphically convex compact sets $K_1, K_2, \dots$, with $K_i \subset \text{int } K_{i+1}$. It follows from Lemma 2.3 that if f is a smooth $\bar{\partial}$ -closed E -valued $(0, r)$ form in a neighborhood of K_i , then on a possibly smaller neighborhood, f is $\bar{\partial}$ -exact, provided $r \geq 1$. We shall also need the following approximation result, true for any $r \geq 0$: the above f can be approximated by $\bar{\partial}$ -closed forms in $\Omega^{0,r}(M, E)$, the approximation taking place in any seminorm $\|\cdot\|_{K_{i-1}, p, j}$; see (2.3).

Indeed, if $r \geq 1$, this follows by writing $f = \bar{\partial}g$ and approximating g in $\|\cdot\|_{K_{i-1}, p, j+1}$ by forms $g' \in \Omega^{0, r-1}(M, E)$, so that $f' = \bar{\partial}g'$ will be the required approximation of f . If $r = 0$, we are talking about approximating holomorphic functions, and the result is a rather simple special case of [Le2, Théorème 1.1].

Now let $r \geq 1$ and $f \in \Omega^{0,r}(M, E)$ be $\bar{\partial}$ -closed. Fix seminorms $p_1 \leq p_2 \leq \dots$ on A defining its topology. In a neighborhood of K_i write $f = \bar{\partial}g_i$. We can arrange recursively that

$$(2.5) \quad \|g_{i+1} - g_i\|_{K_{i-1}, p_i, i} < 2^{-i}, \quad i = 1, 2, \dots$$

Indeed, we note that in general $\bar{\partial}(g_{i+1} - g_i) = 0$ in a neighborhood of K_i ; hence there is a closed $h \in \Omega^{0, r-1}(M, E)$ such that

$$\|g_{i+1} - g_i - h\|_{K_{i-1}, p_i, i} < 2^{-i}.$$

Therefore replacing g_{i+1} by $g_{i+1} - h$ we achieve (2.5), which in turn implies that $g = \lim g_i$ is in $\Omega^{0, r-1}(M, E)$ and satisfies $\bar{\partial}g = f$. Thus we are done if M is open in $\mathbb{C}^k$.

The general case then follows, since a connected Stein manifold M can be embedded in a Stein domain $M' \subset \mathbb{C}^k$ as a holomorphic retract, and the holomorphic retraction induces a monomorphism $H_{\bar{\partial}}^{0,r}(M, E) \rightarrow H_{\bar{\partial}}^{0,r}(M', E')$, where E' is the trivial bundle $A \times M' \rightarrow M'$.

Now we return to a general complex manifold and a Fréchet bundle $E \rightarrow M$. If $\mathfrak{U}, \mathfrak{V}$ are open covers of M for which we defined the spaces $C^{qr}(\mathfrak{U}), C^{qr}(\mathfrak{V})$, and if $\mathfrak{U} \rightarrow \mathfrak{V}$ is a refinement map, then restriction induces homomorphisms $C^{qr}(\mathfrak{V}) \rightarrow C^{qr}(\mathfrak{U})$. The image of $f \in C^{qr}(\mathfrak{V})$ will be denoted $f|_{\mathfrak{U}} \in C^{qr}(\mathfrak{U})$.

Lemma 2.5. *If $\mathfrak{V}$ is a special cover of M , $\mathfrak{U}$ a countable refinement of $\mathfrak{V}$ consisting of Stein open sets, and $q, r \geq 0$, then there is a homomorphism*

$$\varepsilon = \varepsilon^{qr} : Z^{qr}(\mathfrak{U}) \rightarrow Z_b^{qr}(\mathfrak{V})$$

with the following properties:

- (a) if $f \in Z^{0r}(\mathfrak{U})$, then $\varepsilon f|_{\mathfrak{U}} = f$;
- (b) if $q \geq 1$ and $f \in Z^{qr}(\mathfrak{U})$, then there is an $h \in C^{q-1, r}(\mathfrak{U})$ such that

$$\bar{\partial}h = 0 \quad \text{and} \quad \varepsilon f|_{\mathfrak{U}} - f = \delta h;$$

- (c) the restriction of ε to $Z_b^{qr}(\mathfrak{U})$ is continuous;
 (d) if $\mathfrak{U}$ is also special and $f \in Z_b^{qr}(\mathfrak{U})$, then h in (b) can be chosen in $C_b^{q-1,r}(\mathfrak{U})$.

Note that (c) is not automatic, as the topology of $Z_b^{qr}(\mathfrak{U})$ is inherited from $C_b^{qr}(\mathfrak{U})$ and is different from the one inherited from $Z^{qr}(\mathfrak{U}) \subset C^{qr}(\mathfrak{U})$ when $r > 0$.

Proof. Define homomorphisms

$$R = R^{qr} : Z^{qr}(\mathfrak{U}) \rightarrow C^{q-1,r}(\mathfrak{U}), \quad S = S^{qr} : Z_b^{qr}(\mathfrak{V}) \rightarrow C_b^{q,r-1}(\mathfrak{V})$$

as follows. Let $\{\chi_U : U \in \mathfrak{U}\}$ be a smooth partition of unity subordinate to $\mathfrak{U}$; then for $f = (f_{U_0 \dots U_q}) \in Z^{qr}(\mathfrak{U})$, $q \geq 1$,

$$(2.6) \quad (Rf)_{U_1 \dots U_q} = \sum_{U \in \mathfrak{U}} \chi_U f_{UU_1 \dots U_q}.$$

If $g = (g_{V_0 \dots V_q}) \in Z_b^{qr}(\mathfrak{V})$, $r \geq 1$, apply the $\bar{\partial}$ solution operator of Lemma 2.3 componentwise to construct $Sg = (T_{V_0 \dots V_q} g_{V_0 \dots V_q})$. Thus $\delta Rf = f$ and $\bar{\partial} Sg = g$.

We prove the lemma by induction on q . The base case $q = 0$ holds since both $Z^{0r}(\mathfrak{U})$ and $Z_b^{0r}(\mathfrak{V})$ are naturally identified with the space of $\bar{\partial}$ -closed elements of $\Omega^{0,r}(M, E)$, the topology coming from $Z^{0r}(\mathfrak{U})$ being finer than the topology coming from $Z_b^{0r}(\mathfrak{V})$. If $q \geq 1$, assume that $\varepsilon^{q-1,r}$ has been constructed for every r , and let

$$\varepsilon^{qr} = \delta S^{q-1,r+1} \varepsilon^{q-1,r+1} \bar{\partial} R^{qr}.$$

To see that this is indeed a homomorphism $Z^{qr}(\mathfrak{U}) \rightarrow Z_b^{qr}(\mathfrak{V})$, notice that $\delta \bar{\partial} R = \bar{\partial} \delta R = 0$ and $\bar{\partial} \delta S = \delta \bar{\partial} S = 0$, so that $\bar{\partial} R^{qr}$ maps $Z^{qr}(\mathfrak{U})$ in $Z^{q-1,r+1}(\mathfrak{U})$ and $\delta S^{q-1,r+1}$ maps $Z_b^{q-1,r+1}(\mathfrak{V})$ in $Z_b^{qr}(\mathfrak{V})$. If $f \in Z^{qr}(\mathfrak{U})$, let

$$(2.7) \quad h_1 = (S^{q-1,r+1} \varepsilon^{q-1,r+1} \bar{\partial} R^{qr} f)|\mathfrak{U} - R^{qr} f \in C^{q-1,r}(\mathfrak{U}).$$

Then $\delta h_1 = \varepsilon^{qr} f|\mathfrak{U} - f$, and by the inductive assumption

$$\bar{\partial} h_1 = (\varepsilon^{q-1,r+1} \bar{\partial} R^{qr} f)|\mathfrak{U} - \bar{\partial} R^{qr} f$$

either vanishes (if $q = 1$), and we take $h = h_1$ in (b); or else there is $h_2 \in C^{q-2,r+1}(\mathfrak{U})$ such that $\bar{\partial} h_2 = 0$ and $\bar{\partial} h_1 = \delta h_2$. By Theorem 2.4 there is an $h_3 \in C^{q-2,r}(\mathfrak{U})$ with $\bar{\partial} h_3 = h_2$. Then $h = h_1 - \delta h_3$ satisfies $\bar{\partial} h = \bar{\partial} h_1 - \delta \bar{\partial} h_3 = 0$ and $\delta h = \varepsilon^{qr} f|\mathfrak{U} - f$, which completes the proof of (b).

Next (c) follows since $\bar{\partial} R$ is continuous from $Z_b^{qr}(\mathfrak{U})$ to $Z^{q-1,r+1}(\mathfrak{U})$, as one checks by applying $\bar{\partial}$ to (2.6). Finally, if $\mathfrak{U}$ is also special and $f \in Z_b^{qr}(\mathfrak{U})$, $q \geq 1$, then by induction in (2.7) $h_1 \in C_b^{q-1,r}(\mathfrak{U})$ and h_2 above is in $C_b^{q-2,r+1}(\mathfrak{U})$. By Lemma 2.3 one can choose $h_3 \in C_b^{q-2,r}(\mathfrak{U})$, whence $h = h_1 - \delta h_3 \in C_b^{q-1,r}(\mathfrak{U})$, which proves (d).

Proof of Theorem 2.2. First we show that (2.2) and (2.1) are monomorphisms. Being a special cover is stable under small C^∞ perturbations of the boundaries of the covering sets. Therefore we can construct another special cover $\mathfrak{V}$ such that the closure of each $W \in \mathfrak{V}$ is contained in some $V \in \mathfrak{V}$. Suppose $f \in Z_b^q(\mathfrak{V}, E) = Z_b^{q0}(\mathfrak{V})$ represents a cohomology class $[f] \in H_b^q(\mathfrak{V}, E)$ that (2.2) sends to $0 \in H^q(M, E)$. We apply Lemma 2.5a,b with $\mathfrak{U} = \mathfrak{V}$ to conclude that the image of $[\varepsilon^{q0} f] \in H^q(\mathfrak{V}, E)$ in $H^q(M, E)$ factors through $[f] \in H_b^q(\mathfrak{V}, E)$; hence this image is zero. On the other hand, Theorem 2.4 and the Dolbeault isomorphism show that $\mathfrak{V}$ is a Leray cover for the bundle E , so that $H^q(\mathfrak{V}, E) \rightarrow H^q(M, E)$ is a bijection. Hence

$[\varepsilon^{q_0} f] = 0$, i.e., $\varepsilon^{q_0} f = \delta g$ with some $g \in C^{q-1,0}(\mathfrak{W})$. Since $g|_{\mathfrak{W}} \in C_b^{q-1,0}(\mathfrak{W})$ and $\varepsilon^{q_0} f|_{\mathfrak{W}} = \delta(g|_{\mathfrak{W}})$, Lemma 2.5a,d imply f is the coboundary of a bounded cochain: $[f] = 0$. Thus (2.2) is indeed a monomorphism, and with a Stein refinement $\mathfrak{U}$ of $\mathfrak{W}$, so is (2.1), since the canonical homomorphism $H^q(\mathfrak{U}, E) \rightarrow H^q(M, E)$ is a bijection, for the same reason as $H^q(\mathfrak{V}, E) \rightarrow H^q(M, E)$ was.

Next we take an arbitrary Stein refinement $\mathfrak{U}$ of $\mathfrak{W}$ and apply Lemma 2.5 with $\mathfrak{V} = \mathfrak{W}$. Suppose $f \in Z^q(\mathfrak{U}, E) = Z^{q_0}(\mathfrak{U})$ is δ -exact. Again, Lemma 2.5a,b imply that the image of $[\varepsilon^{q_0} f] \in H_b^q(\mathfrak{W}, E)$ in $H^q(M, E)$ is 0; hence by what has been proved, $[\varepsilon^{q_0} f] = 0$. Thus ε^{q_0} maps exact cocycles in $Z^{q_0}(\mathfrak{U})$ to exact cocycles in $Z_b^{q_0}(\mathfrak{W})$. It induces a homomorphism

$$H^q(\mathfrak{U}, E) \rightarrow H_b^q(\mathfrak{W}, E),$$

which, by Lemma 2.5a,b is a right inverse to the restriction homomorphism in (2.1). Passing to the direct limit we obtain a right inverse $H^q(M, E) \rightarrow H_b^q(\mathfrak{W}, E)$ of (2.2). Since (2.1), (2.2) are monomorphisms, the right inverses are left inverses as well.

Corollary 2.6. *If $\mathfrak{W}$ is a special cover of M and $\mathfrak{U}$ is a Stein refinement of $\mathfrak{W}$, then*

$$H^q(M, E) \approx H_b^q(\mathfrak{W}, E) \approx H^q(\mathfrak{U}, E).$$

Corollary 2.7. *If M is compact and $E \rightarrow M$ is a holomorphic Banach bundle, then the cohomology groups $H^q(M, E)$ are complete, seminormable topological vector spaces.*

Proof. Let $\mathfrak{W}$ be a finite special cover of M . Then $H_b^q(\mathfrak{W}, E)$ is a quotient of a Banach space, hence complete and seminormable; it is also isomorphic to $H^q(M, E)$.

3. σ -COMPACT SETS IN FRÉCHET SPACES

Recall that a topological space is σ -compact if it is the countable union of compact subspaces. A homomorphism of Fréchet spaces is compact if the image of some nonempty open set has compact closure.

Lemma 3.1. *The range of a compact homomorphism of Fréchet spaces is contained in a σ -compact set. Conversely, any σ -compact subset K of a Fréchet space Z is contained in the range of a compact homomorphism $Y \rightarrow Z$, where Y is a Banach space.*

Proof. The first part being obvious, we prove the converse only. Assume first that K is compact; by passing to its hull, we may as well assume K is convex and balanced. Such a set defines a Banach space $Y = Z_K = \bigcup \{\lambda K : \lambda \geq 0\}$, endowed with the norm $\|z\| = \min\{\lambda \geq 0 : z \in \lambda K\}$. Then the map $\varphi(z) = z$ from Y to Z is a compact homomorphism and $\varphi(Y) \supset K$. In general $K = \bigcup K_j$ with $K_j \subset Z$ compact, $j \in \mathbb{N}$. For each j there are Banach spaces $(Y_j, \|\cdot\|_j)$ and compact homomorphisms $\varphi_j : Y_j \rightarrow Z$ such that $\varphi_j(Y_j) \supset K_j$. Choose seminorms $p_1 \leq p_2 \leq \dots$ on Z defining its topology; we can assume that $p_j(\varphi_j(y)) \leq \|y\|_j / 2^j$ for $y \in Y_j$. The ℓ^1 -sum of the Y_j 's,

$$Y = \{y = (y_j) : y_j \in Y_j, \|y\| = \sum_j \|y_j\|_j < \infty\},$$

with the norm $\|\cdot\|$ is a Banach space, $\varphi: Y \rightarrow Z$ given by

$$\varphi(y) = \sum_j \varphi_j(y_j), \quad y = (y_j) \in Y,$$

is compact, and $\varphi(Y) \supset \bigcup K_j = K$.

Lemma 3.2. *Let C, Z be Fréchet spaces and $\delta: C \rightarrow Z$ a homomorphism. If there is a σ -compact $L \subset Z$ such that $Z = L + \delta C$, then δC is closed and $\dim Z/\delta C < \infty$.*

Proof. By Lemma 3.1 there exist a Banach space Y and a compact homomorphism $\varphi: Y \rightarrow Z$ with $\varphi Y \supset L$. Define $\psi, \theta: Y \oplus C \rightarrow Z$ by

$$\psi(y, c) = \varphi(y), \quad \theta(y, c) = \delta(c),$$

so that $\psi + \theta: Y \oplus C \rightarrow Z$ is onto. Since ψ is compact, by Schwartz's theorem [Sc], $\theta(Y \oplus C) = \delta C$ is indeed closed and of finite codimension.

4. COMPACT PERTURBATIONS

Let M be a complex manifold and E, F holomorphic Banach bundles over M . If $U \subset M$ is open, we shall say that a homomorphism $k: E|U \rightarrow F|U$ is fiberwise compact if $k|E_x \in \text{Hom}(E_x, F_x)$ is compact for every $x \in U$.

Proposition 4.1. *Suppose $U \Subset V \subset M$ are open and $E|V, F|V$ are trivial. If $k \in \text{Hom}(E|V, F|V)$ is fiberwise compact, then the homomorphism*

$$\alpha: \Gamma(V, E) \ni f \mapsto kf|U \in \Gamma(U, F)$$

is compact.

This would not be true for Fréchet bundles, which is essentially the only reason why Theorem 1.2 is restricted to Banach bundles.

Proof. Let $(A, \|\cdot\|_A), (B, \|\cdot\|_B)$ be Banach spaces such that $E|V, F|V$ are (isomorphic to) $A \times U$, resp. $B \times U$. The set

$$\mathcal{F} = \{f \in \Gamma(V, E) : \sup_{x \in U} \|f(x)\|_A < 1\}$$

is open in $\Gamma(V, E)$. Its image under α is a locally equicontinuous family on U by Cauchy estimates; also $\{\alpha f(x) : f \in \mathcal{F}\}$ has compact closure in B for every $x \in U$. By the Arzelà–Ascoli theorem any sequence in $\alpha(\mathcal{F})$ has a subsequence convergent in the topology of $\Gamma(U, F)$, and so α is indeed compact.

Proposition 4.2. *If F is a compact perturbation of E , then E is also a compact perturbation of F . More precisely, if $\mathfrak{U}$ is an open cover of M and $\varphi_U: E|U \rightarrow F|U$ are freedomorphisms such that $\varphi_U - \varphi_V$ are fiberwise compact for $U, V \in \mathfrak{U}$, then there exist an open cover $\mathfrak{U}'$ of M and freedomorphisms $\psi_{U'}: F|U' \rightarrow E|U'$ for $U' \in \mathfrak{U}'$ such that $\psi_{U'} - \psi_{V'}, \psi_{U'}\varphi_U - id_{E|U' \cap U}$, and $\varphi_U\psi_{U'} - id_{F|U' \cap U}$ are fiberwise compact, $U \in \mathfrak{U}, U', V' \in \mathfrak{U}'$.*

Proof. Given $x \in U \in \mathfrak{U}$, there is a finite codimensional subspace $E'_x \subset E_x$ on which φ_U is injective. In fact, E'_x can be extended to a holomorphic subbundle E' of E , over a neighborhood $N(x) \subset U$ of x , and $\varphi_U|E'$ is still injective. It follows that $F' = \varphi_U(E') \subset F|N(x)$ is a holomorphic subbundle of finite corank, and φ_U restricts to an isomorphism $\varphi': E' \rightarrow F'$. Upon shrinking $N(x)$ we can assume F' has a complementary bundle $F'' \subset F|N(x)$, and in fact, at the price of refining $\mathfrak{U}$

we can assume that the bundles E', F', F'' are defined over all of $U = N(x)$. If $p': F|U = F' \oplus F'' \rightarrow F'$ denotes the projection, then

$$\psi_U = \varphi'^{-1}p' \in \text{Hom}(F|U, E|U)$$

is a fredomorphism. Furthermore $\psi_U \varphi_U - \text{id}_{E|U} = k_U$ and $\varphi_U \psi_U - \text{id}_{F|U} = l_U$ are fiberwise compact (in fact, of finite rank). With ψ_U , $U \in \mathfrak{U} = \mathfrak{U}'$ thus constructed,

$$\psi_V - \psi_U = \psi_U(\varphi_U - \varphi_V)\psi_V - k_U\psi_V + \psi_U l_V$$

are also fiberwise compact, whence the claim follows.

From now on we assume M is compact. It will be convenient to use indexed covers $\mathfrak{U} = \{U_i : i \in I\}$, write $U_{ij\dots} = U_i \cap U_j \cap \dots$, and denote the components $f_{U_i U_j \dots}$ of a cochain by $f_{ij\dots}$.

Let $\mathfrak{U} = \{U_i : i \in I\}$ and $\mathfrak{V} = \{V_i : i \in I\}$ be finite open covers, $U_i \subseteq V_i$. Assume that $E|V_i$ and $F|V_i$ are trivial, and that there are fredomorphisms $\varphi_i : E|V_i \rightarrow F|V_i$ such that $\varphi_i - \varphi_j$ are fiberwise compact. Define homomorphisms $\Phi^q : C^q(\mathfrak{V}, E) \rightarrow C^q(\mathfrak{U}, F)$ between spaces of holomorphic cochains by

$$(4.1) \quad (\Phi^q f)_{i_0\dots i_q} = \varphi_{i_0} f_{i_0\dots i_q}|_{U_{i_0\dots i_q}}, \quad f \in C^q(\mathfrak{V}, E).$$

Proposition 4.3. $\Phi^q \delta - \delta \Phi^{q-1} : C^{q-1}(\mathfrak{V}, E) \rightarrow C^q(\mathfrak{U}, F)$ is compact for $q \geq 1$.

Proof. If $f = (f_{i_1\dots i_q}) \in C^{q-1}(\mathfrak{V}, E)$, then

$$(\Phi^q \delta f)_{i_0\dots i_q} = \varphi_{i_0} \sum_{j=0}^q (-1)^j f_{i_0\dots \hat{i}_j \dots i_q}|_{U_{i_0\dots i_q}}$$

and

$$(\delta \Phi^{q-1} f)_{i_0\dots i_q} = \varphi_{i_1} f_{i_1\dots i_q} + \varphi_{i_0} \sum_{j=1}^q (-1)^j f_{i_0\dots \hat{i}_j \dots i_q}|_{U_{i_0\dots i_q}}.$$

Hence the claim follows from Proposition 4.1, as

$$(\Phi^q \delta f - \delta \Phi^{q-1} f)_{i_0\dots i_q} = (\varphi_{i_0} - \varphi_{i_1}) f_{i_1\dots i_q}|_{U_{i_0\dots i_q}}.$$

Proof of Theorem 1.2. Construct three finite Stein covers $\mathfrak{U} = \{U_i : i \in I\}$, $\mathfrak{V} = \{V_i : i \in I\}$, and $\mathfrak{W} = \{W_i : i \in I\}$ of M , $U_i \subseteq V_i \subseteq W_i$. Make sure $\mathfrak{W}$ is special for both E and F (see Definition 2.1) and so fine that there are fredomorphisms $\varphi_i : E|W_i \rightarrow F|W_i$ and $\psi_i : F|W_i \rightarrow E|W_i$ such that

$$\varphi_i - \varphi_j, \quad \psi_i - \psi_j, \quad \varphi_i \psi_j - \text{id}_{F|W_{ij}}, \quad \text{and} \quad \psi_i \varphi_j - \text{id}_{E|W_{ij}}$$

are all fiberwise compact; see Proposition 4.2. Define homomorphisms Φ^q as in (4.1) and $\Psi^q : C^q(\mathfrak{W}, F) \rightarrow C^q(\mathfrak{U}, E)$ by

$$(\Psi^q f)_{i_0\dots i_q} = \psi_{i_0} f_{i_0\dots i_q}|_{V_{i_0\dots i_q}}, \quad f \in C^q(\mathfrak{W}, F).$$

It follows from Proposition 4.1 that

$$(4.2) \quad \Phi^q \Psi^q f = f|_{\mathfrak{U}} + \kappa f, \quad \text{where} \quad \kappa : C^q(\mathfrak{W}, F) \rightarrow C^q(\mathfrak{U}, F)$$

is compact. By the case $r = 0$ of Lemma 2.5 there are homomorphisms

$$\varepsilon = \varepsilon^q : Z^q(\mathfrak{U}, F) \rightarrow Z^q(\mathfrak{W}, F)$$

such that $\varepsilon f|_{\mathfrak{U}}$ is cohomologous to f for every $f \in Z^q(\mathfrak{U}, F)$. In view of (4.2), then

$$(4.3) \quad \Phi^q \Psi^q \varepsilon \equiv \text{id}_{Z^q(\mathfrak{U}, F)} + \kappa \varepsilon \quad \text{mod } \delta C^{q-1}(\mathfrak{U}, F).$$

By Corollary 2.6, $H^{q+1}(\mathfrak{V}, E) \approx H^{q+1}(M, E)$ is Hausdorff, that is, $\delta C^q(\mathfrak{V}, E) \subset Z^{q+1}(\mathfrak{V}, E)$ is closed. Michael's selection theorem [M, Theorem 3.2''] implies there is a continuous map

$$\mu: \delta C^q(\mathfrak{V}, E) \rightarrow C^q(\mathfrak{V}, E)$$

with $\delta\mu = \text{id}_{\delta C^q(\mathfrak{V}, E)}$. (Typically, Michael's theorem is formulated for Banach, rather than Fréchet spaces. However, it has always been understood that the proof applies for Fréchet spaces as well; see e.g. the last paragraph on page 364 in [M].) Consider the continuous map

$$(4.4) \quad \beta = \Psi^q \varepsilon - \mu \delta \Psi^q \varepsilon: Z^q(\mathfrak{U}, F) \rightarrow C^q(\mathfrak{V}, E).$$

In fact, β maps into $Z^q(\mathfrak{V}, E)$, since $\delta\beta = \delta\Psi^q \varepsilon - \delta\mu \delta\Psi^q \varepsilon = 0$.

Now we come to the main point: for every σ -compact $K \subset Z^q(\mathfrak{V}, E)$ there is a σ -compact $L \subset Z^q(\mathfrak{U}, F)$ such that

$$(4.5) \quad \beta^{-1}(K + \delta C^{q-1}(\mathfrak{V}, E)) \subset L + \delta C^{q-1}(\mathfrak{U}, F).$$

Indeed, by (4.3), (4.4), and since $\delta|Z^q(\mathfrak{V}, F) = 0$,

$$\begin{aligned} \Phi^q \beta &= \Phi^q \Psi^q \varepsilon - \Phi^q \mu \delta \Psi^q \varepsilon \\ &\equiv \text{id}_{Z^q(\mathfrak{U}, F)} + \kappa \varepsilon + \Phi^q \mu (\Psi^{q+1} \delta - \delta \Psi^q) \varepsilon \quad \text{mod } \delta C^{q-1}(\mathfrak{U}, F). \end{aligned}$$

Hence by Lemma 3.1 and the analog of Proposition 4.3 for Ψ , for any $S \subset Z^q(\mathfrak{U}, F)$ we have

$$(4.6) \quad S \subset \Phi^q \beta S + H + \delta C^{q-1}(\mathfrak{U}, F),$$

with some σ -compact $H \subset C^q(\mathfrak{U}, F)$. If now S stands for the left hand side of (4.5), then

$$\begin{aligned} \Phi^q \beta S &= \Phi^q (K + \delta C^{q-1}(\mathfrak{V}, E)) \\ &\subset \Phi^q K + (\Phi^q \delta - \delta \Phi^{q-1}) C^{q-1}(\mathfrak{V}, E) + \delta C^{q-1}(\mathfrak{U}, F). \end{aligned}$$

The first term in the last line is σ -compact, and by Lemma 3.1 and Proposition 4.3, the second is contained in a σ -compact set. Combining this with (4.6) we obtain a σ -compact $L \subset C^q(\mathfrak{U}, F)$ such that

$$S \subset L + \delta C^{q-1}(\mathfrak{U}, F).$$

Since L can be replaced by $L \cap Z^q(\mathfrak{U}, F)$, (4.5) is verified.

By hypothesis, $H^q(M, E) \approx H^q(\mathfrak{V}, E)$ is finite dimensional. We take for $K \subset Z^q(\mathfrak{V}, E)$ a finite dimensional complementary subspace to $\delta C^{q-1}(\mathfrak{V}, E)$. In this case (4.5) gives

$$Z^q(\mathfrak{U}, F) = L + \delta C^{q-1}(\mathfrak{U}, F)$$

with some σ -compact L . Hence Corollary 2.6 and Lemma 3.2 imply

$$H^q(M, F) \approx H^q(\mathfrak{U}, F) = Z^q(\mathfrak{U}, F) / \delta C^{q-1}(\mathfrak{U}, F)$$

is indeed Hausdorff and finite dimensional.

Theorem 1.2 also holds on compact manifolds with strongly pseudoconvex boundaries, by essentially the same proof, and, as said, it should have a sheaf version as well. But in the setting of this paper there does not seem to be much room to improve on it. Without the assumption that $H^{q+1}(M, E)$ is Hausdorff it would not hold. The reasoning to follow was inspired by an idea that the author learned from

Leiterer, who attributed it to Văjăitu. Suppose M is Kähler and $H^1(M, \mathcal{O}) \neq 0$, so that there is a sequence of nontrivial line bundles $\Lambda_k \rightarrow M$ converging to the trivial line bundle. If E is (a suitable Hilbertian completion of) $\bigoplus_1^\infty \Lambda_k$, and F the trivial Hilbert bundle $l^2 \times M \rightarrow M$, then E, F are compact perturbations of one another, $\dim H^0(M, E) < \infty$ but $\dim H^0(M, F) = \infty$. In more detail, let $\mathfrak{U}$ be a Stein cover of M and $g = (g_{UV}) \in Z^1(\mathfrak{U}, \mathcal{O})$ not exact. The multiplicative cocycles $(e^{g_{UV}/k}) \in Z^1(\mathfrak{U}, \mathcal{O}^*)$ are not exact either for large k ; by rescaling g we can assume they are not exact for any $k \in \mathbb{N}$. Thus the line bundles Λ_k they determine are holomorphically nontrivial. Since topologically they nevertheless are trivial, $H^0(M, \Lambda_k) = 0$. The infinite diagonal matrices h_{UV} with diagonal entries $e^{g_{UV}/k}$, $k \in \mathbb{N}$, define a cocycle $h = (h_{UV}) \in Z^1(\mathfrak{U}, \mathrm{GL}(l^2))$ and so a Hilbert space bundle $E \rightarrow M$. Each Λ_k canonically embeds in E , $\bigoplus \Lambda_k$ is dense in E , and there are holomorphic projections $E \rightarrow \Lambda_k$. It follows that $H^0(M, E) = 0$. However, the trivial bundle $F = l^2 \times M \rightarrow M$, a compact perturbation of E , has $\dim H^0(M, F) = \infty$.

By Theorem 1.2 we conclude that $H^1(M, E)$ is not Hausdorff. This example reveals one more notable thing: while all groups $H^q(M, F)$ are Hausdorff, a compact perturbation E has a non-Hausdorff cohomology group—non-Hausdorff cohomology groups of holomorphic Hilbert bundles over compact manifolds were first constructed by Erat in [E]. His construction is similar, being based on a finite rank vector bundle over $\mathbb{P}_1$ that can be deformed into a nonisomorphic bundle.

REFERENCES

- [B] L. Bungart, *On analytic fiber bundles. I. Holomorphic fiber bundles with infinite dimensional fibers*, Topology **7** (1967), 55–68. MR0222338 (36:5390)
- [CS] H. Cartan, J.P. Serre, *Un théorème de finitude concernant les variétés analytiques compactes*, C. R. Acad. Sci. Paris **237** (1953), 128–130. MR0066010 (16:517e)
- [E] M. Erat, *The cohomology of Banach space bundles over 1-convex manifolds is not always Hausdorff*, Math. Nachr. **248/249** (2003), 97–101. MR1950717 (2004a:32016)
- [G] I. Gohberg, *A factorization problem for operator functions (Russian)*, Izv. Akad. Nauk SSSR **28** (1964), 1055–1082. MR0174994 (30:5182)
- [GL] I. Gohberg, J. Leiterer, *General theorems on the factorization of operator-valued functions with respect to a contour, I, (Russian)*, Acta Sci. Math **34** (1973), 103–120; *II*, Acta Sci. Math **35** (1973), 39–59. MR0333790 (48:12113); MR0333791 (48:12114)
- [K] J. Kim, *Holomorphic Banach bundles over compact bases*, Math. Z., to appear.
- [Li1] J. Leiterer, *Banach coherent analytic Fréchet sheaves*, Math. Nachr. **85** (1978), 91–109. MR517643 (80b:32026)
- [Li2] J. Leiterer, *A finiteness theorem for holomorphic Banach bundles*, Ann. Scuola Norm. Sup. Pisa **VI** (2007), 15–37. MR2341512
- [Le1] L. Lempert, *The Dolbeault complex in infinite dimensions, I*, J. Amer. Math. Soc. **11** (1998), 485–520. MR1603858 (99f:58007)
- [Le2] L. Lempert, *Approximation de fonctions holomorphes d'un nombre infini de variables*, Ann. Inst. Fourier Grenoble **49** (1999), 1293–1304. MR1703089 (2001d:32027)
- [LP] L. Lempert and I. Patyi, *Analytic sheaves in Banach spaces*, Ann. Sci. École Norm. Sup. **40** (2007), 453–4863.
- [M] E. Michael, *Continuous selections, I*, Ann. of Math. (2) **63** (1956), 361–382. MR0077107 (17:990e)
- [RS] R.M. Range, Y.-T. Siu, *Uniform estimates for the $\bar{\partial}$ -equation on domains with piecewise smooth strictly pseudoconvex boundaries*, Math. Ann. **206** (1973), 325–354. MR0338450 (49:3214)
- [Sc] L. Schwartz, *Homomorphismes et applications complètement continues*, C. R. Acad. Sci. Paris **236** (1953), 2472–2473. MR0057457 (15:233b)

- [Sk] H. Skoda, *Fibrés holomorphes à base et à fibre de Stein*, Invent. Math. **43** (1977), 97–107. MR0508091 (58:22657)
- [V] D. Vogt, *Vektorwertige Distributionen als Randverteilungen holomorpher Funktionen*, Manuscripta Math. **17** (1975), 267–290. MR0407593 (53:11365)

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, INDIANA 47907-1395