CONVEXITY AND HORIZONTAL SECOND FUNDAMENTAL FORMS FOR HYPERSURFACES IN CARNOT GROUPS

LUCA CAPOGNA, SCOTT D. PAULS, AND JEREMY T. TYSON

ABSTRACT. We use a Riemannian approximation scheme to give a characterization for smooth convex functions on a Carnot group (in the sense of Danielli–Garofalo–Nhiu or Lu–Manfredi–Stroffolini) in terms of the positive semidefiniteness of the horizontal second fundamental form of their graph.

1. INTRODUCTION

In 1996, Caffarelli proposed a notion of convexity for functions on the Heisenberg group in terms of the standard one-dimensional convexity of their restriction to horizontal lines through any fixed point. This notion resurfaced in 2002 when it was independently discovered by Danielli–Garofalo–Nhiu [9] and generalized to arbitrary Carnot groups. At the same time, Lu–Manfredi–Stroffolini [23] (see also [20]) proposed an equivalent definition based on the notion of viscosity subsolutions. For related work, see also Balogh–Rickly [3], Gutiérrez–Montanari [16], [17], Garofalo–Tournier [14], Wang [28] and Magnani [24]. A notion of convexity for sets was introduced in [9], where the relationship between convexity of a function and convexity of its epigraph is studied.

In this paper we propose a notion of convexity for hypersurfaces in Carnot groups, in terms of the *horizontal second fundamental form* II_0 of their graphs (see Definition 3.7). To relate this definition to the previous literature we show that the graphs of (suitably regular) functions have positive definite symmetrized horizontal second fundamental form $(II_0)^*$ if and only if the functions in question are convex (in the sense of Caffarelli, [9], [23], or [20]) and provide quantitative statements of this fact. More precisely, we prove

Theorem 1.1. *Let $\mathbb{G}$ be a Carnot group and denote by $\Gamma^2(\mathbb{G})$ the class of functions twice continuously differentiable along horizontal directions.*

Received by the editors May 15, 2006 and, in revised form, November 14, 2007.

2000 *Mathematics Subject Classification.* Primary 43A80, 53C17; Secondary 22E30, 35H20, 52A41, 53A35.

Key words and phrases. Convexity, Carnot group, second fundamental form, Riemannian approximation.

The authors were partially supported by the National Science Foundation: the first author was partially supported by NSF DMS-0134318; the second author was partially supported by NSF DMS-0306752; the third author was partially supported by NSF DMS-0228807, DMS-0555869; all authors were partially supported by NSF DMS-0503695. Part of the research for this paper was done while the first and third authors were visitors of the Mathematics Department at Dartmouth College in Spring 2006. They gratefully acknowledge the hospitality.

©2010 American Mathematical Society
 Reverts to public domain 28 years from publication

- (i) Let $u \in \Gamma^2(\mathbb{G})$. At every point of $\mathbb{G}$, we have

$$\Lambda \geq (D_0^2 u)^* \geq \lambda \text{ if and only if } \Lambda' \geq (\Pi_0^{\mathcal{G}(u)})^* \geq \lambda',$$

where $(D_0^2 u)^*$ denotes the symmetrized horizontal Hessian of u and $(\Pi_0^{\mathcal{G}(u)})^*$ denotes the symmetrized horizontal second fundamental form of $\mathcal{G}(u)$, the graph of u .

- (ii) $u \in \Gamma^2(\mathbb{G})$ is convex if and only if $(\Pi_0^{\mathcal{G}(u)})^*$ is positive semi-definite.
 (iii) Let $U = \{(x, s) : u(x) \leq s\}$ be the epigraph of a function $u \in \Gamma^2(\mathbb{G})$. Then U is convex (as a set, in the sense of [9]) if and only if the symmetrized horizontal second fundamental form of its boundary, $(\Pi_0^{\partial U})^*$, is positive semi-definite.

In part (i), the bounds Λ' and Λ depend only on each other and on the norm of the horizontal gradient $\nabla_0 u$ of u . A similar statement holds for λ' and λ . We have $\Lambda < \infty$ if and only if $\Lambda' < \infty$, and $\lambda > 0$ if and only if $\lambda' > 0$. See Definition 2.5 and Definition 3.7, respectively, for definitions of the symmetrized horizontal Hessian and the symmetrized horizontal second fundamental form.

The results in Theorem 1.1 are new even in the simplest setting of the first Heisenberg group. The main contribution of the paper, however, lies not in the results themselves but rather in the method of proof, which is based on a careful study of cancellation properties arising from differentiating certain horizontal tensors in the Riemannian approximation scheme.

There is a big gap in terms of regularity of u between our *differential geometric* definition of convexity and the definition in [9], [23], and [20]. While we require two derivatives along the horizontal directions, the original definition can be applied to any lower-semicontinuous function and eventually yields Lipschitz regularity along the horizontal directions. For more details, see [3], [23], [24], [20] or [28].

Our motivations for providing a more geometric understanding of convexity are twofold:

- (i) In the Riemannian setting, the second fundamental form encodes a wealth of critical geometric information on the behavior of the Gauss map; likewise, the study of the horizontal second fundamental form for submanifolds of Carnot groups will allow for a better understanding of the horizontal Gauss map and possibly lead to an approach to the analog of the Alexandrov-Bakelman-Pucci maximum principle for subelliptic linear equations in non-divergence form.
- (ii) Convexity of manifolds evolving by curvature flows (for instance mean curvature flow [19]) is based on applications of maximum principles to certain non-linear evolution equations which describe the behavior of the second fundamental form. If one wants to extend such an analysis to the sub-Riemannian context (and there are plenty of reasons to do so; see [7]), then it is crucial to link convexity to some notion of second fundamental form.

Sub-Riemannian analogs of the second fundamental form have recently been introduced for level sets by Danielli, Garofalo and Nhieu [8], [10] in terms of restrictions of the defining function to horizontal planes. In this paper, we follow a different approach and use the approximation of the sub-Riemannian geometry with a family of Riemannian metrics. The equivalence of the two definitions can be found in [6] and [7]. The horizontal second fundamental form in Definition 3.7 was

proposed originally by Hladky and Pauls [18]. Here we relate it for the first time to the Riemannian approximation scheme and use systematically its symmetrization.

In closing we also want to mention related work of Arcozzi and Ferrari [1] (who studied the Hessian of the distance function in $\mathbb{H}^n$), and Calin and Mangione [5], [4] (who studied the second fundamental form in the Riemannian approximants to $\mathbb{H}$).

After a brief section where we recall basic definitions and results concerning Carnot groups, we analyze the relation between the symmetrized horizontal Hessian of a function and the symmetrized horizontal second fundamental form of its level set. Next we study the particular case of graphs and conclude by presenting concrete examples in the Heisenberg group. In our analysis we first consider only smooth objects (functions and hypersurfaces) and then reduce the regularity assumption to the Folland-Stein class Γ^2 using group mollifiers.

2. NOTATION AND SETUP

The notation and terminology in this paper draws heavily from our monograph [7].

Definition 2.1. Let $\mathbb{G}$ be a Lie group with Lie algebra $\mathfrak{g}$. We say that $\mathbb{G}$ is a *graded nilpotent Lie group* if there exists $k < \infty$ and vector subspaces $\mathcal{V}_i \subset \mathfrak{g}$ so that

$$(2.1) \quad \mathfrak{g} = \mathcal{V}_0 \oplus \mathcal{V}_1 \oplus \cdots \oplus \mathcal{V}_k$$

with the property that $[\mathcal{V}_0, \mathcal{V}_i] = \mathcal{V}_{i+1}$ for $0 \leq i < k$ and $[\mathcal{V}_i, \mathcal{V}_k] = 0$ for $0 \leq i \leq k$.

Given a graded nilpotent Lie group $\mathbb{G}$ equipped with a Riemannian metric g , we say that g is *compatible*¹ with the grading of $\mathfrak{g}$ if (2.1) is an orthogonal splitting of the Lie algebra with respect to g . In this paper, we consider only compatible Riemannian metrics. For a fixed $(\mathbb{G}, g)$, let $\{X_i\}_{i=1}^{m+1} \cup \{Y_j\}_{j=1}^{n+1}$ be an orthonormal set of left-invariant vector fields on $\mathbb{G}$ with the following properties:²

- (1) The span of $\{X_i\}_{i=1}^{m+1}$ is $\mathcal{V}_0$ and the span of $\{Y_j\}_{j=1}^{n+1}$ is $\mathcal{V}_1 \oplus \cdots \oplus \mathcal{V}_k$.
- (2) For each $1 \leq j \leq n+1$, there exists an integer $d(j)$, $1 \leq d(j) \leq k$, so that $Y_j \in \mathcal{V}_{d(j)}$.

We denote by $\exp$ the exponential map from $\mathfrak{g}$ to $\mathbb{G}$.

Definition 2.2. Given a graded nilpotent Lie group $\mathbb{G}$ equipped with a compatible Riemannian metric g , and vector fields $\{X_i\}_{i=1}^{m+1} \cup \{Y_j\}_{j=1}^{n+1}$ as above, we define the left-invariant frame

$$\mathcal{F}_1^{(\mathbb{G}, g)} = \{X_1, \dots, X_{m+1}, Y_1, \dots, Y_{n+1}\}.$$

When the metric is clear from the context, we will simply write this as $\mathcal{F}_1^{\mathbb{G}}$.

We call the $\{X_i\}$ *horizontal vector fields* and call their span, denoted $H\mathbb{G}$, the *horizontal bundle*. We call the $\{Y_j\}$ *vertical vector fields* and call their span, denoted $V\mathbb{G}$, the *vertical bundle*. Then $T\mathbb{G} = H\mathbb{G} \oplus V\mathbb{G}$.

¹In contrast with the intrinsic approach used in [25], where the metric is only assigned on the horizontal distribution at first and then extended to a *taming* metric on the whole ambient space.

²Here we identify left-invariant vector fields with elements of $\mathfrak{g}$.

Definition 2.3. Given a graded nilpotent Lie group $\mathbb{G}$, we define a Carnot–Carathéodory metric on $\mathbb{G}$ by specifying an inner product, $\langle \cdot, \cdot \rangle$, on $\mathcal{V}_0$ and defining the metric by

$$d_{cc}(x, y) = \inf_{\gamma \in \mathcal{A}} \left\{ \int_0^1 \langle \dot{\gamma}(s), \dot{\gamma}(s) \rangle^{\frac{1}{2}} ds \mid \gamma(0) = x, \gamma(1) = y \right\},$$

where $\mathcal{A}$ is the set of all absolutely continuous paths whose derivative, when it exists, lies in $H\mathbb{G}$.

Definition 2.4. A *Carnot group* is a connected, simply connected graded nilpotent Lie group equipped with a Carnot–Carathéodory metric.

Given a Carnot group $\mathbb{G}$ equipped with a compatible Riemannian metric g and basis $\mathcal{F}_1^{\mathbb{G}}$ we denote by $\Gamma^k(\mathbb{G})$, $k \in \mathbb{N}$, the Folland–Stein space of functions which are k times continuously differentiable along horizontal directions (see [13] and [27]).

Definition 2.5. The *horizontal gradient operator* is

$$\nabla_0 = (X_1, \dots, X_{m+1}).$$

In other words,

$$\nabla_0 u = (X_1 u) X_1 + \dots + (X_{m+1} u) X_{m+1}$$

for $u \in \Gamma^1(\mathbb{G})$. For $u \in \Gamma^2(\mathbb{G})$ we define the *horizontal Hessian* of u to be

$$D_0^2 u = (X_i X_j u)_{i,j=1,\dots,m+1}$$

and the *symmetrized horizontal Hessian* of u to be

$$(D_0^2 u)^* = \left(\frac{1}{2} (X_i X_j u + X_j X_i u) \right)_{i,j=1,\dots,m+1}.$$

Various definitions for convexity have been proposed and studied in the setting of Carnot groups. For our purposes the following definition is convenient. See [9], [23], [20], [3], [28], [16], [17] and the notes and bibliographies in these papers for a complete history and detailed list of pertinent references.

Definition 2.6. A function $u \in \Gamma^2(\mathbb{G})$ is *convex* if its symmetrized horizontal Hessian $(D_0^2 u)^*$ is positive semi-definite. A set $A \subset \mathbb{G}$ is *convex* if, for every $x \in A$, the intersection of A with $\exp H_x \mathbb{G}$ is the image under the exponential map of a set in the Lie algebra $\mathfrak{g}$ which is (Euclidean) starlike with respect to x .

Remark 2.7. We note that, as a special case of Proposition 7.6 of [9], a function u on $\mathbb{G}$ is convex if and only if its epigraph $\{(x, s) \in \mathbb{G} \times \mathbb{R} : u(x) \leq s\}$ is convex in the Carnot group $\mathbb{G} \times \mathbb{R}$ (equipped with a product sub-Riemannian structure as in section 4). See also Corollary 4.7.

Next, we construct a family of compatible Riemannian metrics which approximate the Carnot–Carathéodory metric. Compare Korányi [21], Korányi and Reimann [22], Pansu [26] and Gromov [15].

Definition 2.8. Let $(\mathbb{G}, g)$ be a graded nilpotent Lie group with fixed Riemannian metric and coordinate frame $\mathcal{F}_1^{(\mathbb{G}, g)}$. For each $L > 0$ we define Riemannian metrics g_L , the *anisotropic dilations of the metric g* , by setting $g_L(X_i, X_j) = g(X_i, X_j) = \delta_{ij}$, $g_L(X_i, Y_j) = g(X_i, Y_j) = 0$ and $g_L(Y_i, Y_j) = L^{2/d(j)} \delta_{ij}$. Here $d(j)$ is the quantity

defined in the discussion preceding Definition 2.2.³ We define a new, rescaled frame orthonormal with respect to g_L :

$$\mathcal{F}_1^{(\mathbb{G}, g_L)} = \{X_1, \dots, X_{m+1}, \tilde{Y}_1, \dots, \tilde{Y}_{n+1}\},$$

where $X_i, Y_j \in \mathcal{F}_1^{(\mathbb{G}, g)}$ and $\tilde{Y}_j = L^{-1/d(j)} Y_j$.

Example 2.9. The simplest non-abelian example is the first Heisenberg group $\mathbb{H} = \mathbb{H}^1$. We use coordinates (x_1, x_2, x_3) in $\mathbb{H}$ and denote the standard frame in $T\mathbb{H}$ by $X_1 = \partial_{x_1} - \frac{1}{2}x_2\partial_{x_3}$, $X_2 = \partial_{x_2} + \frac{1}{2}x_1\partial_{x_3}$, and $Y_1 = \partial_{x_3}$. Then $\tilde{Y}_1 = L^{-1/2}\partial_{x_3}$. In this frame the metric g_L takes the form

$$(2.2) \quad g_L(x) = \begin{pmatrix} 1 + \frac{1}{4}x_2^2L & -\frac{1}{4}x_1x_2L & -\frac{1}{2}x_2L \\ -\frac{1}{4}x_1x_2L & 1 + \frac{1}{4}x_1^2L & \frac{1}{2}x_1L \\ -\frac{1}{2}x_2L & \frac{1}{2}x_1L & L \end{pmatrix}.$$

The metrics g_L approximate the sub-Riemannian structure of $\mathbb{G}$ in the following sense. Here we denote by d_L the distance function on $\mathbb{G}$ induced by g_L .

Lemma 2.10. *As $L \rightarrow \infty$, the metric spaces $(\mathbb{G}, d_L)$ converge in the Gromov-Hausdorff sense to $(\mathbb{G}, d_{cc})$.*

See [25, p. 18], [15, p. 144, 1.4.D], and [7, Theorem 2.9] for related statements.

Our notation for the Levi-Civita connection is ∇ ; we will use the same notation without regard to the metric g_L . Given a smooth function $u : \mathbb{G} \rightarrow \mathbb{R}$, we denote its Riemannian gradient with respect to g_L by

$$\nabla_L u = (X_1 u) X_1 + \dots + (X_{m+1} u) X_{m+1} + (\tilde{Y}_1 u) \tilde{Y}_1 + \dots + (\tilde{Y}_{n+1} u) \tilde{Y}_{n+1}.$$

Given vectors $U, V \in T\mathbb{G}$, we let $\langle U, V \rangle_L = g_L(U, V)$ and $|U|_L^2 = g_L(U, U)$. We remark that if $U \in H\mathbb{G}$, then $|U|_L$ is independent of U . In this case, we will simply write $|U|$ instead of $|U|_L$. We note directly from the definition that

$$(2.3) \quad \lim_{L \rightarrow \infty} \nabla_L u = \nabla_0 u.$$

We recall the Kozul formula for the Levi-Civita connection [12, p. 55]:

$$(2.4) \quad \langle \nabla_U V, W \rangle_L = \frac{1}{2} \left\{ U \langle V, W \rangle_L + V \langle W, U \rangle_L - W \langle U, V \rangle_L \right. \\ \left. - \langle W, [V, U] \rangle_L - \langle U, [V, W] \rangle_L - \langle V, [U, W] \rangle_L \right\},$$

for vector fields U, V, W . A standard result in sub-Riemannian geometry (see for instance [7, Proposition 3.1]) asserts that the restriction to the horizontal bundle of the Levi-Civita connections for the metrics g_L do not depend on L . We tacitly use this fundamental fact throughout the paper.

The following lemma describes the vertical component of the restriction of ∇ to the horizontal bundle.

Lemma 2.11. *If $U = \sum_{i=1}^{m+1} p_i X_i$, $V = \sum_{i=1}^{m+1} q_i X_i$, $W = \sum_{j=1}^{n+1} r_j \tilde{Y}_j$ and U, V, W are of unit length and mutually g_L -orthogonal, then*

$$(2.5) \quad \langle \nabla_U V, W \rangle_L = -\frac{1}{2} \langle [V, U], W \rangle_L.$$

³A rougher rescaling, with $g_L(Y_i, Y_j) = L\delta_{ij}$, would work just as well for our purposes.

Proof. Observe that

$$\begin{aligned} [U, W] &= \sum_{i=1}^{m+1} \sum_{j=1}^{n+1} \left(p_i X_i r_j \tilde{Y}_j + p_i r_j X_i \tilde{Y}_j \right) - \sum_{i=1}^{m+1} \sum_{j=1}^{n+1} \left(r_j \tilde{Y}_j p_i X_i + r_j p_i \tilde{Y}_j X_i \right) \\ &= \sum_{l=1}^{m+1} \sum_{k=1}^{n+1} \left(p_l (X_l r_k) \tilde{Y}_k + p_l r_k [X_l, \tilde{Y}_k] - r_k (\tilde{Y}_k p_l) X_l \right), \end{aligned}$$

while $[V, W]$ is given by the same expression with q_i replacing p_i . A direct computation yields $\langle [U, W], V \rangle_L = -\sum_{i,j=1}^{m+1} r_j q_i (\tilde{Y}_j p_i)$ and $\langle [V, W], U \rangle_L = -\sum_{i,j=1}^{m+1} r_j p_i (\tilde{Y}_j q_i)$. Here we used the fact that $\langle [X_i, Y_j], X_k \rangle_L = 0$. Summing the two identities gives

$$\begin{aligned} \langle [U, W], V \rangle_L + \langle [V, W], U \rangle_L &= -\sum_{i=1}^{m+1} \sum_{j=1}^{n+1} r_j (p_i \tilde{Y}_j q_i + q_i \tilde{Y}_j p_i) \\ &= -\sum_{i=1}^{m+1} \sum_{j=1}^{n+1} r_j \tilde{Y}_j (p_i q_i) = -\sum_{j=1}^{n+1} r_j \tilde{Y}_j \langle U, V \rangle_L = 0. \end{aligned}$$

Now using the orthonormality of $\{U, V, W\}$ and (2.4) we have the desired result. $\square$

3. HYPERSURFACES IN CARNOT GROUPS

We fix a Carnot group, $\mathbb{G}$, and a family of metrics g_L associated to a fixed Riemannian metric g as in the previous section. We note that the (topological) dimension of $\mathbb{G}$ is $N = m + n + 2$. Let M be a smooth hypersurface in $\mathbb{G}$ given by

$$M = \{x \in \mathbb{G} : u(x) = 0\},$$

where $u : \mathbb{G} \rightarrow \mathbb{R}$ is a smooth function with non-vanishing (Riemannian with respect to any g_L metric) gradient in a neighborhood of M .

Denote by $\Sigma(M) = \{x \in M \mid T_x M \supset H_x \mathbb{G}\}$ the *characteristic set* of M . The (Riemannian) surface measure of Σ is zero (see [11] or [2] as well as [7] for further references and more precise statements).

Definition 3.1. For any non-characteristic point, the *unit horizontal normal* to M is defined as the normalized projection of the Riemannian normal to the horizontal subbundle. In the basis $\mathcal{F}_1^{\mathbb{G}}$, it is given as

$$(3.1) \quad \mathbf{n}_0 = \frac{(X_1 u) X_1 + \cdots + (X_{m+1} u) X_{m+1}}{|(X_1 u) X_1 + \cdots + (X_{m+1} u) X_{m+1}|}.$$

Note that the definition of $\mathbf{n}_0$ does not depend on L . Letting

$$(3.2) \quad \mathbf{n}_L = \frac{\nabla_L u}{|\nabla_L u|_L}$$

denote the Riemannian unit normal (with respect to g_L), we note that $\lim_{L \rightarrow \infty} \mathbf{n}_L = \mathbf{n}_0$ uniformly in compact subsets of $M \setminus \Sigma(M)$.

We next consider a basis for $T\mathbb{G}|_M$ adapted to the submanifold M :

$$\mathcal{F}_2^{\mathbb{G}} = \{Z_1, \dots, Z_{m+n+1}, \mathbf{n}_L\},$$

where $\{Z_i\}$ is an orthonormal basis for TM in the metric g_L .

Definition 3.2. We denote the Riemannian second fundamental form of M in the coordinate frame $\mathcal{F}_2^{\mathbb{G}}$ in (M, g_L) by

$$\Pi_L^{M, \mathcal{F}_2^{\mathbb{G}}} = (h_{ij}^L) = (\langle \nabla_{Z_i} \mathbf{n}_L, Z_j \rangle_L)_{i,j=1, \dots, m+n+1}$$

and the Riemannian Hessian of a smooth function $u : \mathbb{G} \rightarrow \mathbb{R}$ with respect to any orthonormal basis $\mathcal{F}^{\mathbb{G}} = \{W_1, \dots, W_{m+n+2}\}$ of $T\mathbb{G}$ is

$$\text{Hess}_L^{\mathcal{F}^{\mathbb{G}}}(u) = (\langle \nabla_{W_i}(\nabla_L u), W_j \rangle_L)_{i,j=1, \dots, m+n+2}$$

at any point in $\mathbb{G}$. We will omit the frame from the notation whenever we wish to emphasize coordinate independence.

The following lemma is a simple computation using (3.2) and Definition 3.2.

Lemma 3.3. *Let M be a smooth hypersurface in $\mathbb{G}$ given as a level set of a smooth function u on $\mathbb{G}$. Then for any point in M , and for $i, j = 1, \dots, m+n+1$ we have*

$$|\nabla_L u|_L \left(\Pi_L^{M, \mathcal{F}_2^{\mathbb{G}}} \right)_{ij} = \left(\text{Hess}_L^{\mathcal{F}_2^{\mathbb{G}}}(u) \right)_{ij}.$$

In the following we view $\text{Hess}_L(u)$ as a bilinear form on $T\mathbb{G}$ by the equation

$$\text{Hess}_L(u)(W, V) = \langle \nabla_W(\nabla_L u), V \rangle_L,$$

for all $V, W \in T\mathbb{G}$. When we want to consider (as we do here) only the restriction of this form to the subspace TM we will use the notation $\text{Hess}_L(u)|_{TM}(W, V)$, $W, V \in TM$. Lemma 3.3 can be rephrased as

$$\text{Hess}_L(u)|_{TM} = |\nabla_L u|_L \Pi_L^M,$$

or more explicitly,⁴

$$\begin{aligned} \text{Hess}_L(u)(W, V) &= \sum_{i,j=1}^{m+n+1} \left(\text{Hess}_L^{\mathcal{F}_2^{\mathbb{G}}}(u) \right)_{ij} a_i b_j \\ &= |\nabla_L u|_L \sum_{i,j=1}^{m+n+1} \left(\Pi_L^{M, \mathcal{F}_2^{\mathbb{G}}} \right)_{ij} a_i b_j = \Pi_L^M(W, V), \end{aligned}$$

where $W = \sum_{i=1}^{m+n+1} a_i Z_i$ and $V = \sum_{i=1}^{m+n+1} b_i Z_i$.

To take advantage of this identity in the limit as $L \rightarrow \infty$ we need to extract *horizontal data* out of it. To accomplish this, we construct a specific basis of the form of $\mathcal{F}_2^{\mathbb{G}}$ that will be useful for our purposes. To facilitate this, we make several definitions concerning a decomposition of $\mathbf{n}_L$ at non-characteristic points.

Definition 3.4. Let $\mathbf{n}_L$ be the Riemannian unit normal and $\mathbf{n}_0$ be the unit horizontal normal to M . At any non-characteristic point we set

$$\begin{aligned} T_0 &= \frac{\mathbf{n}_L - \langle \mathbf{n}_L, \mathbf{n}_0 \rangle_L \mathbf{n}_0}{|\mathbf{n}_L - \langle \mathbf{n}_L, \mathbf{n}_0 \rangle_L \mathbf{n}_0|_L}, \\ a_L &= \langle \mathbf{n}_L, \mathbf{n}_0 \rangle_L = \frac{|\nabla_0 u|}{|\nabla_L u|_L}, \\ b_L &= \langle \mathbf{n}_L, T_0 \rangle_L. \end{aligned}$$

Thus

$$\mathbf{n}_L = a_L \mathbf{n}_0 + b_L T_0.$$

⁴Note that the summation on the left-hand side does not extend over the full Hessian matrix.

We may rewrite b_L in a more precise manner as follows:

$$\begin{aligned} b_L^2 &= 1 - a_L^2 = |\nabla_L u|_L^{-2} (|\nabla_L u|_L^2 - |\nabla_0 u|^2) \\ &= |\nabla_L u|_L^{-2} \left(\sum_{j=1}^{n+1} (\tilde{Y}_j u)^2 \right) \leq \frac{1}{L^{2/k} |\nabla_L u|_L^2} \left(\sum_{j=1}^{n+1} (Y_j u)^2 \right), \end{aligned}$$

where k is as in (2.1). To summarize, using (2.3) we have the following lemma:

Lemma 3.5. *At non-characteristic points, $\lim_{L \rightarrow \infty} a_L = 1$ and $\lim_{L \rightarrow \infty} b_L = 0$.*

We now define a new basis:

Definition 3.6. Let $(\mathbb{G}, g_L)$ be as above and M be a smooth hypersurface in $\mathbb{G}$ given as a level set of a function $u : \mathbb{G} \rightarrow \mathbb{R}$. Then

$$\mathcal{F}_3^{(\mathbb{G}, g_L)} = \{e_0, e_1, \dots, e_m, T_1, \dots, T_n, \mathbf{n}_L\}$$

is a basis for $T\mathbb{G}|_{M \setminus \Sigma(M)}$ of the form of $\mathcal{F}_2^{\mathbb{G}}$, where

$$e_0 = b_L \mathbf{n}_0 - a_L T_0,$$

$\{e_1, \dots, e_m\}$ is an orthonormal basis for $HM = TM \cap H\mathbb{G}$, and $\{T_1, \dots, T_n\}$ is an orthonormal basis for $VM = TM \cap V\mathbb{G}$. Again, when the metric is understood, we will suppress it in the notation.

We emphasize that e_0 is not a horizontal vector field. Moreover, as $L \rightarrow \infty$, we see that $b_L \rightarrow 0$ and hence $e_0 \rightarrow -T_0$.

We now introduce a central concept of this paper, the horizontal second fundamental form of a hypersurface in a Carnot group.

Definition 3.7. Given a smooth hypersurface $M \subset (\mathbb{G}, g_L)$ with adapted basis $\mathcal{F}_3^{\mathbb{G}}$, the *horizontal second fundamental form* at a non-characteristic point is

$$\Pi_0^M = (h_{ij}^0) = (\langle \nabla_{e_i} \mathbf{n}_0, e_j \rangle_L)_{i,j=1}^m.$$

Note that the entries in Π_0^M are independent of L despite the fact that ∇_{e_i} and e_j are *a priori* dependent on L . Moreover, note that Π_0^M is not necessarily symmetric. In a similar fashion, we define the *vertical second fundamental form* by

$$(v_{ij,L}) = (\langle \nabla_{e_i} T_0, e_j \rangle_L)_{i,j=1}^m.$$

Note that $v_{ij,L}$ does depend on L .

Remark 3.8. While the second fundamental form Π_L^M is defined as acting on all of the tangent space, we will always tacitly indicate with Π_0^M the restriction of the horizontal second fundamental form to HM .

Lemma 3.9. *Given a smooth hypersurface $M \subset (\mathbb{G}, g_L)$ with adapted basis $\mathcal{F}_3^{\mathbb{G}}$, at non-characteristic points we have*

- (1) $\Pi_L^{M, \mathcal{F}_3^{\mathbb{G}}} |_{HM} = (h_{ij}^L)_{i,j=1}^m = (a_L h_{ij}^0 + b_L v_{ij,L})_{i,j=1}^m$;
- (2) *as $L \rightarrow \infty$, the restriction of $\Pi_L^{M, \mathcal{F}_3^{\mathbb{G}}}$ to the horizontal tangent space converges to the symmetrized horizontal second fundamental form, i.e.,*

$$\lim_{L \rightarrow \infty} h_{ij}^L = \frac{1}{2} (h_{ij}^0 + h_{ji}^0);$$

- (3) for any smooth function $u : \mathbb{G} \rightarrow \mathbb{R}$, $\lim_{L \rightarrow \infty} \text{Hess}_L^{\mathcal{F}_1^{\mathbb{G}}}(u) |_{H\mathbb{G}} = (D_0^2 u)^*$; i.e., the limit of the Riemannian Hessians restricted to the horizontal directions is the symmetrized horizontal Hessian.

Proof. (1) follows from the definitions of a_L, b_L, h_{ij}^0 and $v_{ij,L}$. To determine the existence and the value of the limit, we observe that

$$(3.3) \quad h_{ij}^L = a_L \frac{h_{ij}^0 + h_{ji}^0}{2} + b_L \frac{v_{ij,L} + v_{ji,L}}{2},$$

since $h_{ij}^L = h_{ji}^L$. By Lemma 2.11 we have $v_{ij,L} + v_{ji,L} = 0$ and the conclusion follows from (3.3) and from Lemma 3.5.

(3) is a trivial consequence of the definition of the Riemannian Hessian, the fact that it is symmetric and the definition of $D_0^2 u$. $\square$

Definition 3.10. Let $M \subset \mathbb{G}$ be a smooth hypersurface and denote by $(\Pi_0^M)^*$ its symmetrized horizontal second fundamental form. The (horizontal) principal curvatures $k_1, \dots, k_m$ of M at a point $x \in M$ are the eigenvalues of $(\Pi_0^M)^*(x)$. The (horizontal) mean curvature $\mathcal{H}_0^M$ of M at x is the trace of $(\Pi_0^M)^*(x)$. Finally, the horizontal Gauss curvature $\mathcal{G}_0^M$ of M at x is $\det(\Pi_0^M)^*(x)$. We will also denote by $\mathcal{H}_L^M$ and $\mathcal{G}_L^M$ the mean curvature and the Gauss curvature of M with respect to the metric g_L .

In view of Lemma 3.9(2) we have

Lemma 3.11. If $M \subset \mathbb{G}$ is a smooth hypersurface, then

$$\lim_{L \rightarrow \infty} \mathcal{H}_L^M = \mathcal{H}_0^M$$

at non-characteristic points.

Remark 3.12. Note that this convergence result is false for the Gauss curvature. Even in the first Heisenberg group, regardless of the choice of the approximating metric g_L , the Gauss curvature is always of the order of L at all non-characteristic points. See [7, Section 3.3].

On the other hand, if M is the zero level set of a smooth function $u : \mathbb{G} \rightarrow \mathbb{R}$, then $\mathcal{H}_L^M = \text{div}_{g_L}(\mathbf{n}_L)$ and a direct computation yields

$$(3.4) \quad |\nabla_L u| \mathcal{H}_L^M = \sum_{i,j=1}^{m+n+2} \left(\delta_{ij} - \frac{Z_i u Z_j u}{|\nabla_L u|^2} \right) Z_i Z_j u,$$

where we have denoted by $Z_1, \dots, Z_{m+n+2}$ any relabelling of the frame $\mathcal{F}_1^{(\mathbb{G},g)}$. As a direct consequence of (3.4) and of Definition 2.8 and (2.3) we recover the familiar expression

$$\mathcal{H}_0^M = \frac{1}{|\nabla_0 u|} \sum_{i,j=1}^{m+1} \left(\delta_{ij} - \frac{X_i u X_j u}{|\nabla_0 u|^2} \right) X_i X_j u$$

on $M \setminus \Sigma$. In fact we have the following convergence result:

$$(3.5) \quad \lim_{L \rightarrow \infty} |\nabla_L u| \mathcal{H}_L^M = \begin{cases} |\nabla_0 u| \mathcal{H}_0^M & \text{on } M \setminus \Sigma, \\ \sum_{i=1}^{m+1} X_i^2 u & \text{on } \Sigma. \end{cases}$$

This observation is due to Citti.

We conclude this section with an explicit relation between the symmetrized horizontal Hessian and the second fundamental form.

Proposition 3.13. *Let $u : \mathbb{G} \rightarrow \mathbb{R}$ be a smooth function and $M = \{u = 0\}$. Let $A = [\mathcal{F}_3^{\mathbb{G}} \rightarrow \mathcal{F}_1^{\mathbb{G}}]$ be the change of basis matrix from $\mathcal{F}_3^{\mathbb{G}}$ to $\mathcal{F}_1^{\mathbb{G}}$. We have the following identity of bilinear forms:*

$$|\nabla_0 u|(\Pi_0^M)^* = (A^t(D_0^2 u)^* A)|_{HM}$$

at non-characteristic points. More explicitly, for $V = \sum_{i=1}^m v_i e_i$ and $W = \sum_{i=1}^m w_i e_i$,

$$\sum_{i,j=1}^m |\nabla_0 u|(\Pi_0^M)^*_{ij} v_i w_j = \sum_{i,j=1}^m (A^t(D_0^2 u)^* A)_{ij} v_i w_j.$$

Proof. As both the Hessian and the second fundamental form are bilinear, the result follows immediately from Lemma 3.3 and Lemma 3.9 (2), (3). $\square$

Corollary 3.14. *If $u : \mathbb{G} \rightarrow \mathbb{R}$ is a smooth convex function and $M = \{x \in \mathbb{G} : u(x) = 0\}$ a smooth hypersurface, then $(\Pi_0^M)^*$ is positive semi-definite at any non-characteristic point.*

Proof. Since a smooth convex function satisfies $(D_0^2 u)^* \geq 0$, the conclusion follows directly from Proposition 3.13. $\square$

Example 3.15. The converse of Corollary 3.14 is false. Consider the first Heisenberg group $\mathbb{H}$, identified with $\mathbb{R}^3$ using coordinates (x_1, x_2, x_3) . Let $u(x_1, x_2, x_3) = x_3 - \frac{x_1 x_2}{2}$ and $M = \{u = 0\}$. The symmetrized horizontal Hessian of u ,

$$(D_0^2 u)^* = \begin{pmatrix} 0 & -\frac{1}{2} \\ -\frac{1}{2} & 0 \end{pmatrix},$$

has eigenvalues $\{\pm \frac{1}{2}\}$ and hence is not positive semi-definite. However, in this case, Π_0^M and $\Pi_L^{M, \mathcal{F}_3^{\mathbb{G}}}|_{HM}$ are 1×1 matrices whose only entry is zero. To see this, we will create the basis $\mathcal{F}_3^{\mathbb{H}}$ for M . Using the notation of Example 2.9, we compute $X_1 u = -x_2$, $X_2 u = 0$, and $\tilde{Y}_1 u = \frac{1}{\sqrt{L}}$. Then we have

$$\mathbf{n}_0 = \text{sign}(-x_2)X_1,$$

and so

$$\mathbf{n}_L = a_L \mathbf{n}_0 + b_L T_0,$$

where $T_0 = \tilde{Y}_1$, $a_L = |x_2|/\sqrt{1/L + x_2^2}$ and $b_L = 1/\sqrt{1 + Lx_2^2}$. Finally, $e_0 = b_L \mathbf{n}_0 - a_L T_0$ and $e_1 = X_2$. Thus

$$\Pi_L^{M, \mathcal{F}_3^{\mathbb{G}}}|_{HM} = (\langle \nabla_{e_1} \mathbf{n}_L, e_1 \rangle_L) = (-\langle \nabla_{e_1} e_1, \mathbf{n}_L \rangle_L) = (-\langle \nabla_{X_2} X_2, \mathbf{n}_L \rangle_L) = (0).$$

We note that $(D_0^2 u)_{1,1}^* = 0$, verifying Proposition 3.13 in this case.

4. GRAPHS IN $\mathbb{G} \times \mathbb{R}$

Let $\mathbb{G}$ be a Carnot group equipped with a family of approximating metrics g_L as in the previous section. Consider the Carnot group $\mathbb{G} \times \mathbb{R}$ with coordinates (x, s) , $x \in \mathbb{G}$, $s \in \mathbb{R}$. On the level of the Lie algebra, this corresponds to adding a single vector field, $S = \frac{\partial}{\partial s}$, to the first layer of the grading. For $u : \mathbb{G} \rightarrow \mathbb{R}$, let

$$(4.1) \quad \mathcal{G}(u) = \{(x, s) \in \mathbb{G} \times \mathbb{R} : u(x) = s\}.$$

Such graph hypersurfaces have no characteristic points; hence we will be able to use the results from the previous section without restriction.

In the first part of this section, we consider only smooth functions u . Later, we extend our main result to functions $u \in \Gamma^2(\mathbb{G})$.

For $(\mathbb{G} \times \mathbb{R}, g_L \oplus ds^2)$ we construct the three bases introduced in the previous section. We write⁵

$$\mathcal{F}_1^{\mathbb{G} \times \mathbb{R}} = \mathcal{F}_1^{\mathbb{G}} \cup \{S\}$$

and

$$\mathcal{F}_2^{\mathbb{G} \times \mathbb{R}} = \{\zeta_1, \dots, \zeta_{m+n+2}, \nu_L\},$$

where $\zeta_1, \dots, \zeta_{m+n+2}$ is a basis for $T\mathcal{G}(u)$ and ν_L is the unit normal in the $g_L \oplus ds^2$ metric. Here we are using a notational convention which will persist for the balance of the paper. Given an object defined for both $\mathbb{G}$ and $\mathbb{G} \times \mathbb{R}$, we use roman letters to denote the object for $\mathbb{G}$ and Greek letters to denote the corresponding object for $\mathbb{G} \times \mathbb{R}$.

Lemma 3.3 applied to this case yields

$$(4.2) \quad \sqrt{1 + |\nabla_L u|_L^2} \Pi_L^{\mathcal{G}(u)} = (\text{Hess}_L(u(x) - s))|_{T\mathcal{G}(u)}.$$

As before, we construct the basis $\mathcal{F}_3$ by introducing

$$\begin{aligned} \nu_L &= \frac{|\nabla_L u|_L \mathbf{n}_L - S}{\sqrt{1 + |\nabla_L u|_L^2}}, \\ \nu_0 &= \frac{|\nabla_0 u| \mathbf{n}_0 - S}{\sqrt{1 + |\nabla_0 u|^2}}, \\ \tau_0 &= \frac{\tilde{Y}_1 u \tilde{Y}_1 + \dots + \tilde{Y}_{n+1} u \tilde{Y}_{n+1}}{|\tilde{Y}_1 u \tilde{Y}_1 + \dots + \tilde{Y}_{n+1} u \tilde{Y}_{n+1}|_L}. \end{aligned}$$

Then we define $\alpha_L = \langle \nu_L, \nu_0 \rangle_L$ and $\beta_L = \langle \nu_L, \tau_0 \rangle_L$ so that $\nu_L = \alpha_L \nu_0 + \beta_L \tau_0$. From this we define $\epsilon_0 = \beta_L \nu_0 - \alpha_L \tau_0$ and the full frame for $\mathbb{G} \times \mathbb{R}$:

$$\mathcal{F}_3^{\mathbb{G} \times \mathbb{R}} = \{\epsilon_0, \epsilon_1, \dots, \epsilon_{m+1}, \tau_1, \dots, \tau_n, \nu_L\},$$

where $\{\epsilon_1, \dots, \epsilon_{m+1}\}$ is an orthonormal basis for $H\mathcal{G}(u)$ and $\{\tau_1, \dots, \tau_n\}$ is an orthonormal basis for $V\mathcal{G}(u)$. Applying Lemma 3.3 gives for $i, j = 1, \dots, m+1$,

$$(4.3) \quad \begin{aligned} \left(A^t \text{Hess}_L^{\mathcal{F}_1^{\mathbb{G} \times \mathbb{R}}} (u(x) - s) A \right)_{ij} &= \left(\text{Hess}_L^{\mathcal{F}_3^{\mathbb{G} \times \mathbb{R}}} (u(x) - s) \right)_{ij} \\ &= \sqrt{1 + |\nabla_L u|_L^2} \left(\Pi_L^{\mathcal{G}(u), \mathcal{F}_3^{\mathbb{G} \times \mathbb{R}}} \right)_{ij}, \end{aligned}$$

where $A = [\mathcal{F}_3^{\mathbb{G} \times \mathbb{R}} \rightarrow \mathcal{F}_1^{\mathbb{G} \times \mathbb{R}}]$ is the (orthogonal) change of coordinates matrix from $\mathcal{F}_3^{\mathbb{G} \times \mathbb{R}}$ to $\mathcal{F}_1^{\mathbb{G} \times \mathbb{R}}$.

Lemma 4.1. *Let $(\mathbb{G}, g_L)$ be a Riemannian graded nilpotent Lie group and $u : \mathbb{G} \rightarrow \mathbb{R}$ a smooth function. Then,*

$$(4.4) \quad \text{Hess}_L^{\mathcal{F}_1^{\mathbb{G} \times \mathbb{R}}} (u(x) - s)|_{H(\mathbb{G} \times \mathbb{R})} = \begin{pmatrix} (D_0^2 u)^* & 0 \\ 0 & 0 \end{pmatrix}.$$

⁵By an abuse of notation we denote with the same symbol vectors in $\mathfrak{g}$ and their lifts in $\mathfrak{g} \times \mathbb{R}$.

Proof. Denote by Γ_{ik}^j the Christoffel symbols for g_L . By the definition of the Hessian one has

$$\begin{aligned} (\text{Hess}_L^{\mathcal{F}_1^{\mathbb{G} \times \mathbb{R}}}(u(x) - s))_{X_i, X_j} &= \langle \nabla_{X_i} \nabla_L(u(x) - s), X_j \rangle_L \\ &= \langle \nabla_{X_i} (\sum_k X_k u X_k + \sum_l \tilde{Y}_l u \tilde{Y}_l - S), X_j \rangle_L \\ &= \sum_k X_i X_k u \langle X_k, X_j \rangle_L + \sum_k X_k u \langle \nabla_{X_i} X_k, X_j \rangle_L + \sum_l \tilde{Y}_l u \langle \nabla_{X_i} \tilde{Y}_l, X_j \rangle_L \\ &= X_i X_j u + \sum_k X_k u \Gamma_{ik}^j + \sum_l \tilde{Y}_l \Gamma_{i, l+m+1}^j. \end{aligned}$$

To simplify the latter, we note that since $[X_i, X_j] \notin H(\mathbb{G} \times \mathbb{R})$, $\Gamma_{i,k}^j = 0$ for $1 < i, j, k < m+1$. The symmetry of the (Riemannian) Hessian yields

$$(\text{Hess}_L^{\mathcal{F}_1^{\mathbb{G} \times \mathbb{R}}}(u(x) - s))_{X_i, X_j} = (\text{Hess}_L^{\mathcal{F}_1^{\mathbb{G} \times \mathbb{R}}}(u(x) - s))_{X_j, X_i},$$

which, coupled with the anti-symmetry of the Γ_{ik}^j in i and j [12, Corollary 3.3, p. 54], yields

$$\begin{aligned} X_i X_j u + \sum_l \tilde{Y}_l \Gamma_{i, l+m+1}^j &= \frac{1}{2} (X_i X_j u + X_j X_i u) + \sum_l \tilde{Y}_l u (\Gamma_{i, l+m+1}^j + \Gamma_{j, l+m+1}^i) \\ &= \frac{1}{2} (X_i X_j u + X_j X_i u). \end{aligned}$$

This shows that the upper left-hand block of the Hessian, written with respect to the basis $\mathcal{F}_1^{\mathbb{G} \times \mathbb{R}}$, is equal to the symmetrized horizontal Hessian. Moreover, the Kozul formula implies that $\nabla_S U = 0 = \nabla_U S$ for $U \in \{X_1, \dots, X_{m+1}, \tilde{Y}_1, \dots, \tilde{Y}_{n+1}\}$, showing that the remaining entries are zero. $\square$

Remark 4.2. Note that, in contrast with the Euclidean case, the derivation of (4.4) for general Carnot groups relies on significant cancellation properties stemming from the underlying symmetry of the Christoffel symbols.

If M is a symmetric matrix we will write $\lambda \leq M \leq \Lambda$ if $\lambda \leq \langle MW, W \rangle \leq \Lambda$ for all unit vectors W . We emphasize that the quantities Λ and λ denote *functions* on $\mathbb{G}$. We can now prove our main theorem.

Theorem 4.3. *Let $\mathbb{G}$ be a Carnot group equipped with a family of Riemannian approximating metrics, $\{g_L\}$, and let $u : \mathbb{G} \rightarrow \mathbb{R}$ be a smooth function.*

(a) *If $\Lambda \geq (D_0^2 u)^* \geq \lambda \geq 0$, then*

$$\frac{\lambda}{(1 + |\nabla_L u|_L^2)^{3/2}} \leq \Pi_L^{\mathcal{G}(u)}|_{H^{\mathcal{G}(u)}} \leq \frac{\Lambda}{(1 + |\nabla_L u|_L^2)^{1/2}},$$

for all $L > 0$, and

$$\frac{\lambda}{(1 + |\nabla_0 u|^2)^{3/2}} \leq (\Pi_0^{\mathcal{G}(u)})^* \leq \frac{\Lambda}{(1 + |\nabla_0 u|^2)^{1/2}}.$$

(b)(i) *If $\lambda \leq \Pi_L^{\mathcal{G}(u)}|_{H^{\mathcal{G}(u)}} \leq \Lambda$, then*

$$\lambda(1 + |\nabla_L u|_L^2)^{1/2} \leq (D_0^2 u)^* \leq \Lambda(1 + |\nabla_L u|_L^2)^{3/2}.$$

(b)(ii) *If $\lambda \leq (\Pi_0^{\mathcal{G}(u)})^* \leq \Lambda$, then*

$$\lambda(1 + |\nabla_0 u|^2)^{1/2} \leq (D_0^2 u)^* \leq \Lambda(1 + |\nabla_0 u|^2)^{3/2}.$$

(c) The horizontal Gauss curvature of $\mathcal{G}(u)$ is given by

$$\det((\Pi_0^{\mathcal{G}(u)})^*) = \left(\sqrt{1 + |\nabla_0 u|^2} \right)^{-(m+3)} \det[(D_0^2 u)^*].$$

Here $\mathcal{G}(u)$ denotes the graph of u in $\mathbb{G} \times \mathbb{R}$.

In particular,

Corollary 4.4. *A smooth function $u : \mathbb{G} \rightarrow \mathbb{R}$ is convex if and only if $(\Pi_0^{\mathcal{G}(u)})^*$ is positive semi-definite.*

Remark 4.5. Compare Theorem 4.3(c) with Definition 10.3 in [9], where the horizontal Gauss curvature was first introduced in the literature. Although our definition differs from the one in [9], Theorem 4.3 shows that they are in fact equivalent.

For the proof of Theorem 4.3, we begin with a simple linear algebraic lemma.

Lemma 4.6. *Let Π be a codimension-one linear subspace of $\mathbb{R}^{d+1}$ transverse to the x_{d+1} -axis. Let $\theta \in [0, \pi/2)$ be the angle between the normal to Π and the x_{d+1} -axis. Let M be an $n \times n$ symmetric matrix and denote by $\tilde{M}$ the restriction to Π of the bilinear form in $\mathbb{R}^{d+1}$ associated to the matrix*

$$\begin{pmatrix} M & 0 \\ 0 & 0 \end{pmatrix}.$$

Then (i) $\lambda \leq M \leq \Lambda$ implies $\lambda \cos^2 \theta \leq \tilde{M} \leq \Lambda$, and (ii) $\lambda \leq \tilde{M} \leq \Lambda$ implies $\lambda \leq M \leq \Lambda \sec^2 \theta$.

Furthermore, if we define the determinant of a bilinear form to be the determinant of a matrix representation (in any orthonormal frame), then

$$(4.5) \quad \det(\tilde{M}) = \cos^2 \theta \det(M).$$

Proof. Since M is symmetric and the conclusions are invariant under conjugation with an orthonormal matrix in $\mathbb{R}^d$ (i.e., a rotation around the x_{d+1} -axis), we can assume without loss of generality that the unit normal $\vec{v}$ to Π is given by the vector $(\sin \theta U_1, \cos \theta)$, where U_1 is the first vector in the canonical orthonormal basis $\{U_1, \dots, U_d\}$ of $\mathbb{R}^d$. Set $\tilde{U}_1 = (-\cos \theta U_1, \sin \theta)$, and $\tilde{U}_i = (U_i, 0) \in \mathbb{R}^{d+1}$, $i = 2, \dots, d$. Then $\{\tilde{U}_1, \dots, \tilde{U}_d\}$ is an orthonormal frame for Π . If $w = (w', w_{d+1}) \in \Pi$, then $|w_{d+1}| \leq \tan \theta |w'|$ and

$$|w| \leq \sec \theta |w'|.$$

Indeed, $0 = \langle w, \vec{v} \rangle = \sin \theta \langle w', U_1 \rangle + w_{d+1} \cos \theta$.

If $\lambda \leq M \leq \Lambda$ and $w = (w', w_{d+1}) \in \Pi$, then

$$\lambda \cos^2 \theta |w|^2 \leq \lambda |w'|^2 \leq \langle Mw', w' \rangle = \langle \tilde{M}w, w \rangle \leq \Lambda |w'|^2 \leq \Lambda |w|^2.$$

On the other hand, if $\lambda \leq \tilde{M} \leq \Lambda$ and $u \in \mathbb{R}^d$, choose w_{d+1} so that $w = (u, w_{d+1}) \in \Pi$. (This is possible by the transversality assumption.) Then

$$\lambda |u|^2 \leq \lambda |w|^2 \leq \langle \tilde{M}w, w \rangle = \langle Mu, u \rangle \leq \Lambda |w|^2 \leq \Lambda \sec^2 \theta |u|^2.$$

In order to prove (4.5) we observe that $\det \tilde{M}$ can be computed by evaluating the determinant of the matrix $(\langle \tilde{M} \tilde{U}_i, \tilde{U}_j \rangle)_{ij}$. The latter coincides with $\langle MU_i, U_j \rangle$ with the first row and the first column both multiplied by $-\cos \theta$. Consequently $\det \tilde{M} = \cos^2 \theta \det(M)$. This completes the proof of the lemma. $\square$

Proof of Theorem 4.3. By (4.3), the estimates

$$\Lambda \geq \sqrt{1 + |\nabla_L u|_L^2} \Pi_L^{\mathcal{G}(u), \mathcal{F}_3^{\mathbb{G} \times \mathbb{R}}} |_{H\mathcal{G}(u)} \geq \lambda$$

hold if and only if the estimates

$$(4.6) \quad \Lambda \geq \left(A^T \text{Hess}_L^{\mathcal{F}_1^{\mathbb{G} \times \mathbb{R}}} (u(x) - s) A \right)_{i,j=1,\dots,m+1} \geq \lambda$$

hold. Here A is as in (4.3).

By Lemma 4.1 we see that (4.6) holds if and only if

$$(4.7) \quad \Lambda \geq \left(A^T \begin{pmatrix} (D_0^2 u)^* & 0 \\ 0 & 0 \end{pmatrix} A \right) |_{H\mathcal{G}(u)} \geq \lambda.$$

To finish the proof of the first part of (a) and part (b)(i), we apply Lemma 4.6 with $\Pi = (H\mathcal{G}(u))$ (represented in the basis $\mathcal{F}_1^{\mathbb{G} \times \mathbb{R}}$), $M = (D_0^2 u)^*$ and $d = m + 1$. Note that the vectors $X_i + (X_i u)S$, $i = 1, \dots, m + 1$, form a basis for Π , and the cosine of θ is precisely the component of the unit normal ν_L in the direction $-x_{d+1}$, i.e.,

$$\cos \theta = \frac{1}{\sqrt{1 + |\nabla_L u|_L^2}}.$$

The results in the first part of (a) and (b)(i) follow from (4.7) and Lemma 4.6. The second part of (a) and (b)(ii) now follow by passing to the limit as $L \rightarrow \infty$. Finally, (c) follows from (4.3) and (4.5). $\square$

Combining Corollary 4.4 and Remark 2.7 gives the following additional corollary.

Corollary 4.7. *Let $U = \{(x, s) : u(x) \leq s\}$ be the epigraph of a smooth function $u : \mathbb{G} \rightarrow \mathbb{R}$. Then U is convex (as a set, in the sense of [9]) if and only if the symmetrized horizontal second fundamental form of its boundary $(\Pi_0^{\partial U})^*$ is positive semi-definite.*

Next we reduce the smoothness assumptions on the function u and its level sets M from C^∞ to Γ^2 . Let $u \in \Gamma^2(\mathbb{G})$ and let f be a standard mollifier in $\mathbb{G}$ (see e.g. [13]). For $\epsilon > 0$ set $f_\epsilon(x) = \epsilon^{-Q} f(\delta_{\epsilon^{-1}}(x))$, where $Q = \sum_{i=1}^k i \dim(\mathcal{V}_i)$ is the homogeneous dimension of $\mathbb{G}$ and $\delta_s : \mathbb{G} \rightarrow \mathbb{G}$, $s > 0$, is the anisotropic dilation

$$\delta_s \left(\exp \left(\sum_{i=1}^{m+1} a_i X_i + \sum_{j=1}^{n+1} b_j Y_j \right) \right) = \exp \left(\sum_{i=1}^{m+1} s a_i X_i + \sum_{j=1}^{n+1} s^{d(j)} b_j Y_j \right).$$

Let

$$u_\epsilon(x) = f_\epsilon * u(x) = \int_{\mathbb{G}} f_\epsilon(y) u(y^{-1}x) dy$$

denote the group convolution. Clearly u_ϵ is smooth and $u_\epsilon \rightarrow u$, $X_i u_\epsilon = (X_i u) * f_\epsilon \rightarrow X_i u$, and $X_i X_j u_\epsilon = (X_i X_j u) * f_\epsilon \rightarrow X_i X_j u$ as $\epsilon \rightarrow 0$, uniformly on compact subsets of $\mathbb{G}$.

Note that while at the level of the g_L metrics it is not possible to compute the full Hessian of a function $u \in \Gamma^2(\mathbb{G})$ or the second fundamental form of its graph (as there is no *a priori* differentiability along higher layers of the stratification), both the horizontal Hessian of u and the horizontal second fundamental form of $\mathcal{G}(u)$ are meaningful for $u \in \Gamma^2(\mathbb{G})$ (at least, off the characteristic set). Applying the previous results to u_ϵ and its graph for $L = 0$, we finally obtain the $\Gamma^2(\mathbb{G})$ (and $L = 0$) versions of Theorem 4.3 and Corollaries 4.4 and 4.7 as stated in Theorem 1.1.

Theorem 4.8. *Let $\mathbb{G}$ be a Carnot group.*

(i) *If $u \in \Gamma^2(\mathbb{G})$ and $\mathcal{G}(u) \subset \mathbb{G} \times \mathbb{R}$ denotes the graph of u , then*

$$\Lambda \geq (D_0^2 u)^* \geq \lambda \quad \text{for some functions } 0 \leq \lambda \leq \Lambda < \infty$$

if and only if

$$\Lambda' \geq (\Pi_0^{\mathcal{G}(u)})^* \geq \lambda' \quad \text{for some functions } 0 \leq \lambda' \leq \Lambda' < \infty$$

at every point of $\mathbb{G}$.

(ii) *$u \in \Gamma^2(\mathbb{G})$ is convex if and only if $(\Pi_0^{\mathcal{G}(u)})^*$ is positive semi-definite.*

(iii) *Let $U = \{(x, s) : u(x) \leq s\}$ be the epigraph of $u \in \Gamma^2(\mathbb{G})$. Then U is convex (as a set, in the sense of [9]) if and only if the symmetrized horizontal second fundamental form of its boundary, $(\Pi_0^{\partial U})^*$, is positive semi-definite.*

The choices of Λ', λ' in terms of Λ, λ coincide with those in the statement of Theorem 4.3.

5. EXAMPLES AND FURTHER DISCUSSION

In this section, we illustrate with some explicit computations and examples in the Heisenberg group. First, we give an explicit choice of the frame $\mathcal{F}_3^{\mathbb{H} \times \mathbb{R}}$ relative to which the horizontal component of the second fundamental form has a particularly nice formulation in terms of the symmetrized horizontal Hessian. Then, we present an explicit example showing the possibility of a difference between the horizontal second fundamental form and the limit of the Riemannian second fundamental forms.

Example 5.1. We use the standard vector fields $\{X_1, X_2, Y_1, S\}$ on the product space $\mathbb{H} \times \mathbb{R} = \{(x_1, x_2, x_3, s)\}$, as in Example 2.9. Here $S = \partial/\partial s$ is the vector field governing motion in the s -direction. Let $u \in C^\infty(\mathbb{H})$. We construct an adapted frame $\mathcal{F}_3^{\mathbb{H} \times \mathbb{R}} = \{\epsilon_0, \epsilon_1, \epsilon_2, \nu_L\}$ for the graph $\mathcal{G}(u)$ following the procedure described in section 4: setting $p = X_1 u$, $q = X_2 u$, $r = \tilde{Y}_1 u$,

$$l = |\nabla_0 u| = \sqrt{p^2 + q^2}, \quad m = \sqrt{1 + p^2 + q^2},$$

and

$$l_L = |\nabla_L u| = \sqrt{p^2 + q^2 + r^2}, \quad m_L = \sqrt{1 + p^2 + q^2 + r^2},$$

we choose tangent vectors

$$\epsilon_0 = \frac{pr}{mm_L} X_1 + \frac{qr}{mm_L} X_2 - \frac{m}{m_L} \tilde{Y}_1 - \frac{r}{mm_L} S, \quad \epsilon_1 = \bar{q} X_1 - \bar{p} X_2,$$

and

$$\epsilon_2 = \frac{\bar{p}}{m} X_1 + \frac{\bar{q}}{m} X_2 + \frac{l}{m} S,$$

orthogonal to the normal vector

$$\nu_L = \frac{p}{m_L} X_1 + \frac{q}{m_L} X_2 + \frac{r}{m_L} \tilde{Y}_1 - \frac{1}{m_L} S.$$

Here we use the convenient shorthand $\bar{p} = p/l$ and $\bar{q} = q/l$.

Then $A = [\mathcal{F}_3^{\mathbb{H} \times \mathbb{R}} \rightarrow \mathcal{F}_1^{\mathbb{H} \times \mathbb{R}}]$ is the matrix whose transpose has entries which are precisely the $X_1, X_2, \tilde{Y}_1$, and S coefficients of $\epsilon_0, \epsilon_1, \epsilon_2, \nu_L$. In the basis $\mathcal{F}_1^{\mathbb{H} \times \mathbb{R}}$, the matrix for $\text{Hess}_L^{\mathcal{F}_1^{\mathbb{H} \times \mathbb{R}}}(u(x) - s)$ has principal 2×2 minor $H^* := (D_0^2 u)^*$ and final

column and row consisting entirely of zeros. A simple computation shows that the horizontal second fundamental form, which is the central 2×2 minor of

$$\frac{1}{m} A^t \cdot \text{Hess}_L^{\mathcal{F}_1^{\mathbb{H} \times \mathbb{R}}}(u(x) - s) \cdot A,$$

takes the form

$$(5.1) \quad \Pi_0^{\mathcal{G}(u)} \Big|_{H\mathcal{G}(u)} = \begin{pmatrix} \frac{1}{m} \langle H^* v^\perp, v^\perp \rangle & \frac{1}{m^2} \langle H^* v, v^\perp \rangle \\ \frac{1}{m^2} \langle H^* v, v^\perp \rangle & \frac{1}{m^3} \langle H^* v, v \rangle \end{pmatrix},$$

where $v = (\bar{p}, \bar{q})$ and $v^\perp = (\bar{q}, -\bar{p})$. Then

$$\det \Pi_0^{\mathcal{G}(u)} \Big|_{H\mathcal{G}(u)} = \frac{\det(D_0^2 u)^*}{(1 + |\nabla_0 u|^2)^2}$$

is the horizontal Gauss curvature of $\mathcal{G}(u)$ (see Theorem 4.3 in the previous section), and

$$\text{Trace } \Pi_0^{\mathcal{G}(u)} \Big|_{H\mathcal{G}(u)} = \frac{\text{Trace}(D_0^2 u)^* + |\nabla_0 u|^3 \mathcal{H}_0^M}{(1 + |\nabla_0 u|^2)^{3/2}} = \mathcal{H}_0^{\mathcal{G}(u)}$$

is the horizontal mean curvature of $\mathcal{G}(u)$; see Definition 3.10. Here we wrote M for the level set of u .

Example 5.2. In this example, we show that the limit of the Riemannian second fundamental forms may differ from the horizontal second fundamental form. With notation as in the previous example, we consider the graph $\mathcal{G}(u)$ of $u(x_1, x_2, x_3) = \frac{x_1 x_2}{2} - x_3$ in $\mathbb{H} \times \mathbb{R}$. Then $X_1(u - s) = y$, $X_2(u - s) = 0$, $\tilde{Y}_1(u - s) = -1/\sqrt{L}$, and $S(u - s) = -1$. The Riemannian unit normal to $\mathcal{G}(u)$ is

$$\nu_L = \frac{x_2}{\sqrt{1 + \frac{1}{L} + x_2^2}} X_1 - \frac{1}{\sqrt{L} \sqrt{1 + \frac{1}{L} + x_2^2}} \tilde{Y}_1 - \frac{1}{\sqrt{1 + \frac{1}{L} + x_2^2}} S,$$

and the unit horizontal normal is

$$\nu_0 = \frac{y}{\sqrt{1 + x_2^2}} X_1 - \frac{1}{\sqrt{1 + x_2^2}} S.$$

We construct ϵ_0 as described in section 4 and choose two additional horizontal tangent vector fields $\epsilon_1 = X_2$ and

$$\epsilon_2 = \frac{1}{\sqrt{1 + x_2^2}} X_1 + \frac{x_2}{\sqrt{1 + x_2^2}} S,$$

tangent to $\mathcal{G}(u)$, and write $\mathcal{F} = \{\epsilon_0, \epsilon_1, \epsilon_2, \nu_L\}$ and $\mathcal{F}_0 = \{\epsilon_0, \epsilon_1, \epsilon_2, \nu_0\}$. Next, we compute the horizontal component of the Riemannian second fundamental form, as well as the entire horizontal second fundamental form, using the observations

$$\nabla_{\epsilon_1} \epsilon_2 = \left(\frac{1}{\sqrt{1 + x_2^2}} \right)_{x_2} X_1 + \left(\frac{x_2}{\sqrt{1 + x_2^2}} \right)_{x_2} X_4 - \frac{\sqrt{L}}{2\sqrt{1 + x_2^2}} \tilde{Y}_1$$

and

$$\nabla_{\epsilon_2} \epsilon_1 = \frac{\sqrt{L}}{2\sqrt{1 + x_2^2}} \tilde{Y}_1.$$

Taking the inner product with ν_L and ν_0 , respectively, gives

$$\Pi_L^{\mathcal{G}(u), \mathcal{F}} \Big|_{H\mathcal{G}(u)} = \begin{pmatrix} 0 & -\frac{1}{2\sqrt{1 + \frac{1}{L} + x_2^2}} \\ -\frac{1}{2\sqrt{1 + \frac{1}{L} + x_2^2}} & 0 \end{pmatrix}$$

and

$$\Pi_0^{\mathcal{G}(u), \mathcal{F}_0} \Big|_{H\mathcal{G}(u)} = \begin{pmatrix} 0 & -\frac{1}{\sqrt{1+x_2^2}} \\ 0 & 0 \end{pmatrix}.$$

Observe that $\lim_{L \rightarrow \infty} \Pi_L = \Pi_0^*$.

REFERENCES

- [1] ARCOZZI, N., AND FERRARI, F. Metric normal and distance function in the Heisenberg group. *Math. Z.* 256, 3 (2007), 661–684. MR2299576 (2008e:49043)
- [2] BALOGH, Z. Size of characteristic sets and functions with prescribed gradient. *J. Reine Angew. Math.* 564 (2003), 63–83. MR2021034 (2005d:43007)
- [3] BALOGH, Z., AND RICKLY, M. Regularity of convex functions on Heisenberg groups. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* 2, 4 (2003), 847–868. MR2040646 (2006d:43006)
- [4] CALIN, O., AND MANGIONE, V. *Variational calculus on sub-Riemannian manifolds*. Balkan J. Geom. Appl. 8 (2003), 21–32. MR2030318 (2004m:53051)
- [5] CALIN, O., AND MANGIONE, V. Geodesics with constraints on Heisenberg manifolds. *Results Math.* 1–2, 44 (2003), 44–53. MR2011905 (2004j:37123)
- [6] CAPOGNA, L., AND CITTI, G. Generalized mean curvature flow in Carnot groups. *Comm. Partial Differential Equations* 34, 7–9 (2009), 937–956. MR2560306
- [7] CAPOGNA, L., DANIELLI, D., PAULS, S. D., AND TYSON, J. T. *An introduction to the Heisenberg group and the sub-Riemannian isoperimetric problem*, vol. 259 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2007. MR2312336
- [8] DANIELLI, D., GAROFALO, N., AND NHIEU, D.-M. Minimal surfaces, surfaces of constant mean curvature and isoperimetry in Carnot groups. preprint, 2001.
- [9] DANIELLI, D., GAROFALO, N., AND NHIEU, D.-M. Notions of convexity in Carnot groups. *Comm. Anal. Geom.* 11, 2 (2003), 263–341. MR2014879 (2004m:22014)
- [10] DANIELLI, D., GAROFALO, N., AND NHIEU, D. M. Sub-Riemannian calculus on hypersurfaces in Carnot groups. *Adv. Math.* 215, 1 (2007), 292–378. MR2354992
- [11] DERRIDI, M. Sur un théorème de traces. *Ann. Inst. Fourier (Grenoble)* 22, 2 (1972), 73–83. MR0343011 (49:7755)
- [12] DO CARMO, M. P. *Riemannian geometry*. Mathematics: Theory & Applications. Birkhäuser Boston Inc., Boston, MA, 1992. Translated from the second Portuguese edition by Francis Flaherty. MR1138207 (92i:53001)
- [13] FOLLAND, G. B. Subelliptic estimates and function spaces on nilpotent Lie groups. *Ark. Mat.* 2, 13 (1975), 161–207. MR0494315 (58:13215)
- [14] GAROFALO, N., AND TOURNIER, F. New properties of convex functions in the Heisenberg group. *Trans. Amer. Math. Soc.* 358, 5 (2006), 2011–2055 (electronic). MR2197446 (2006k:26017)
- [15] GROMOV, M. Carnot-Carathéodory spaces seen from within. In *Sub-Riemannian Geometry*, vol. 144 of *Progress in Mathematics*. Birkhäuser, Basel, 1996, pp. 79–323. MR1421823 (2000f:53034)
- [16] GUTIÉRREZ, C. E., AND MONTANARI, A. Maximum and comparison principles for convex functions on the Heisenberg group. *Comm. Partial Differential Equations* 29, 9–10 (2004), 1305–1334. MR2103838 (2005h:35024)
- [17] GUTIÉRREZ, C. E., AND MONTANARI, A. On the second order derivatives of convex functions on the Heisenberg group. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* 3, 2 (2004), 349–366. MR2075987 (2005f:26038)
- [18] HLADKY, R. K., AND PAULS, S. D. Constant mean curvature surfaces in sub-Riemannian spaces. *J. Diff. Geom.* 79 (2008), 111–139. MR2401420
- [19] HUISKEN, G. Flow by mean curvature of convex surfaces into spheres. *J. Diff. Geom.* 1, 20 (1984), 237–266. MR772132 (86j:53097)
- [20] JUUTINEN, P., LU, G., MANFREDI, J. J., AND STROFFOLINI, B. Convex functions on Carnot groups. *Rev. Mat. Iberoam.* 23, 1 (2007), 191–200. MR2351130
- [21] KORÁNYI, A. *Geometric aspects of analysis on the Heisenberg group*. Ist. Naz. Alta Mat. Francesco Severi, 1983, pp. 209–258. MR748865 (85h:32055)

- [22] KORÁNYI, A., AND REIMANN, H. M. Horizontal normal vectors and conformal capacity of spherical rings in the Heisenberg group. *Bull. Sci. Math. (2)* 111, 1 (1987), 3–21. MR886958 (88e:22013)
- [23] LU, G., MANFREDI, J. J., AND STROFFOLINI, B. Convex functions on the Heisenberg group. *Calc. Var. Partial Differential Equations* 19, 1 (2004), 1–22. MR2027845 (2004m:35088)
- [24] MAGNANI, V. Lipschitz continuity, Aleksandrov theorem, and characterizations for H -convex functions. *Math. Ann.* 334 (2006), 199–233. MR2208954 (2007c:49005)
- [25] MONTGOMERY, R. *A tour of sub-Riemannian geometries, their geodesics and applications*. No. 91 in Mathematical Surveys and Monographs. American Mathematical Society, 2002. MR1867362 (2002m:53045)
- [26] PANSU, P. Une inégalité isopérimétrique sur le groupe de Heisenberg. *C. R. Acad. Sci. Paris Sér. I Math.* 295, 2 (1982), 127–130. MR676380 (85b:53044)
- [27] ROTHCHILD, L. P., AND STEIN, E. M. Hypoelliptic differential operators and nilpotent groups. *Acta Math.* 137, 3-4 (1976), 247–320. MR0436223 (55:9171)
- [28] WANG, C. Y. Viscosity convex functions on Carnot groups. *Proc. Amer. Math. Soc.* 133, 4 (2005), 1247–1253 (electronic). MR2117228 (2006a:58026)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ARKANSAS, FAYETTEVILLE, ARKANSAS 72701
E-mail address: `lcapogna@uark.edu`

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, HANOVER, NEW HAMPSHIRE 03755
E-mail address: `scott.pauls@dartmouth.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, URBANA-CHAMPAIGN, ILLINOIS 61801
E-mail address: `tyson@math.uiuc.edu`