1-POINT GROMOV-WITTEN INVARIANTS OF THE MODULI SPACES OF SHEAVES OVER THE PROJECTIVE PLANE

WEI-PING LI AND ZHENBO QIN

ABSTRACT. The Gieseker-Uhlenbeck morphism maps the Gieseker moduli space of stable rank-2 sheaves on a smooth projective surface to the Uhlenbeck compactification and is a generalization of the Hilbert-Chow morphism for Hilbert schemes of points. When the surface is the complex projective plane, we determine all the 1-point genus-0 Gromov-Witten invariants extremal with respect to the Gieseker-Uhlenbeck morphism. The main idea is to understand the virtual fundamental class of the moduli space of stable maps by studying the obstruction sheaf and using a meromorphic 2-form on the Gieseker moduli space.

1. INTRODUCTION

Recently there has been intensive interest in studying the quantum cohomology and Gromov-Witten theory of Hilbert schemes of points on algebraic surfaces. Two main reasons are the connections with the Donaldson-Thomas theory of 3-folds and with Ruan's Cohomological Crepant Resolution Conjecture. Roughly speaking, the Crepant Resolution Conjecture asserts that the quantum cohomology of an orbifold Z coincides with the quantum cohomology of a crepant resolution Y of Z after analytic continuation and specialization of quantum parameters. For an algebraic surface X , let $X^{[n]}$ be the Hilbert scheme of n points on X and $\text{Sym}^n(X)$ be the n -th symmetric product of X . It is well known that $X^{[n]}$ is smooth of dimension $2n$ and the Hilbert-Chow morphism $\Phi : X^{[n]} \rightarrow \text{Sym}^n(X)$ is a crepant resolution of the global orbifold $\text{Sym}^n(X)$.

A natural generalization of the Hilbert-Chow morphism Φ is the *Gieseker-Uhlenbeck morphism* Ψ from the moduli space of Gieseker semistable rank-2 torsion-free sheaves on X to the Uhlenbeck compactification space. This morphism was constructed in [LJ1, Mor], and was shown to be crepant [LJ2, Q-Z] when the Gieseker moduli space is smooth. For the projective plane $X = \mathbb{P}^2$, the moduli space $\overline{\mathfrak{M}}(n)$ of Gieseker semistable sheaves V on X with $c_1(V) = -1$ and $c_2(V) = n$ is a smooth irreducible projective variety of dimension $4n - 4$ when $n \geq 1$. In [Q-Z], it is proved that there is exactly one primitive integral class $\mathfrak{f} \in H_2(\overline{\mathfrak{M}}(n); \mathbb{Z})$ contracted by the Gieseker-Uhlenbeck morphism $\Psi : \overline{\mathfrak{M}}(n) \rightarrow \overline{\mathfrak{U}}(n)$.

Received by the editors February 6, 2009 and, in revised form, June 1, 2009.

2000 *Mathematics Subject Classification.* Primary 14D20, 14N35.

Key words and phrases. Gieseker moduli spaces, Gromov-Witten invariants.

The first author was partially supported by the grants GRF601905 and GRF601808.

The second author was partially supported by an NSF grant.

©2010 American Mathematical Society
 Reverts to public domain 28 years from publication

The goal of this paper is to determine all the 1-point genus-0 Gromov-Witten invariants $\langle \alpha \rangle_{0,d\mathfrak{f}}$, $\alpha \in H^{8n-12}(\overline{\mathfrak{M}}(n); \mathbb{C})$ extremal with respect to Ψ for $n \geq 3$. When $n \geq 3$, the homology group $H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$ is of rank 6, and a basis is given by $\{\Xi_1, \dots, \Xi_6\}$ (see §4 for details). The Poincaré duals $\text{PD}(\Xi_1), \dots, \text{PD}(\Xi_6)$ form a basis of $H^{8n-12}(\overline{\mathfrak{M}}(n); \mathbb{C})$.

Theorem 1.1. *Let $d \geq 1$ and $n \geq 3$. The Gromov-Witten invariants $\langle \alpha \rangle_{0,d\mathfrak{f}}$ for the classes $\alpha = \text{PD}(\Xi_1), \dots, \text{PD}(\Xi_6) \in H^{8n-12}(\overline{\mathfrak{M}}(n); \mathbb{C})$ are respectively equal to*

$$-6/d^2, \quad 12/d^2, \quad 0, \quad -6/d^2, \quad 0, \quad 0.$$

When $n = 1$, the moduli space $\overline{\mathfrak{M}}(n)$ is a point. When $n = 2$, the fourth Betti number b_4 of the moduli space $\overline{\mathfrak{M}}(n)$ is equal to 3, which is different from the case $n \geq 3$. The result for $n = 2$ will appear elsewhere via a different method (see Remark 3.2).

An interesting observation is that the 1-point genus-0 Gromov-Witten invariants $\langle \alpha \rangle_{0,d\mathfrak{f}}$ are independent of the second Chern class n .

Conjecture 1.2. *Let $d \geq 1$ and $n \geq 3$. Then the extremal genus-0 Gromov-Witten invariants $\langle \alpha_1, \dots, \alpha_k \rangle_{0,d\mathfrak{f}}$ of the moduli space $\overline{\mathfrak{M}}(n)$ are independent of n .*

There are two main ideas in our proof of Theorem 1.1. The first one is to determine the restriction of the obstruction sheaf of the Gromov-Witten theory for $\overline{\mathfrak{M}}(n)$ to certain open subsets of the moduli space $\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f})$ of stable maps. This enables us to determine the 1-point invariants $\langle \alpha \rangle_{0,d\mathfrak{f}}$ for the first four cohomology classes $\alpha = \text{PD}(\Xi_1), \dots, \text{PD}(\Xi_4)$.

The second one is to study the support of the virtual fundamental class

$$[\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f})]^{\text{vir}} \in A_{4n-6}(\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f}))$$

using the techniques developed in [K-L, L-L]. By introducing a suitable meromorphic 2-form Θ on the Gieseker moduli space $\overline{\mathfrak{M}}(n)$, we show that

$$(1.1) \quad \text{ev}_1 \left(\text{Supp}([\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f})]^{\text{vir}}) \right) \subset \mathfrak{T}_{C_0}(n) \coprod \mathfrak{U}_{C_0}(n),$$

where $\text{ev}_1 : \overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f}) \rightarrow \overline{\mathfrak{M}}(n)$ is the evaluation map, and $\mathfrak{T}_{C_0}(n)$ (respectively, $\mathfrak{U}_{C_0}(n)$) is the subset of $\overline{\mathfrak{M}}(n)$ consisting of all the nonlocally free sheaves V such that $V|_{C_0}$ contains torsion (respectively, $V|_{C_0}$ is torsion-free and unstable). This allows us to show that $\langle \alpha \rangle_{0,d\mathfrak{f}} = 0$ for $\alpha = \text{PD}(\Xi_5), \text{PD}(\Xi_6)$.

This paper is organized as follows. In §2, the Gromov-Witten theory is reviewed. In §3, we recall some properties of the Gieseker moduli space $\overline{\mathfrak{M}}(n)$ and the Gieseker-Uhlenbeck morphism Ψ . We study the boundary divisor of $\overline{\mathfrak{M}}(n)$ consisting of nonlocally free sheaves in $\overline{\mathfrak{M}}(n)$. In §4, the basis $\{\Xi_1, \dots, \Xi_6\}$ for $H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$ is constructed. In §5, we analyze the obstruction sheaf of the Gromov-Witten theory for $\overline{\mathfrak{M}}(n)$. In §6, (1.1) is proved. In §7, we verify Theorem 1.1.

2. STABLE MAPS AND GROMOV-WITTEN INVARIANTS

Let Y be a smooth projective variety. A k -pointed *stable map* to Y consists of a complete nodal curve D with k distinct ordered smooth points $p_1, \dots, p_k$ and a morphism $\mu : D \rightarrow Y$ such that the data $(\mu, D, p_1, \dots, p_k)$ has only finitely many automorphisms. In this case, the stable map is denoted by $[\mu : (D; p_1, \dots, p_k) \rightarrow Y]$. For a fixed homology class $\beta \in H_2(Y, \mathbb{Z})$, let $\overline{\mathfrak{M}}_{g,k}(Y, \beta)$ be the coarse moduli space

parameterizing all the stable maps $[\mu : (D; p_1, \dots, p_k) \rightarrow Y]$ such that $\mu_*[D] = \beta$ and the arithmetic genus of D is g . Then, we have the evaluation map:

$$(2.1) \quad ev_k : \overline{\mathfrak{M}}_{g,k}(Y, \beta) \rightarrow Y^k$$

defined by $ev_k([\mu : (D; p_1, \dots, p_k) \rightarrow Y]) = (\mu(p_1), \dots, \mu(p_k))$. It is known [F-P, LT1, LT2, Beh, B-F] that the coarse moduli space $\overline{\mathfrak{M}}_{g,k}(Y, \beta)$ is projective and has a virtual fundamental class $[\overline{\mathfrak{M}}_{g,k}(Y, \beta)]^{\text{vir}} \in A_{\mathfrak{d}}(\overline{\mathfrak{M}}_{g,k}(Y, \beta))$, where

$$(2.2) \quad \mathfrak{d} = -(K_Y \cdot \beta) + (\dim(Y) - 3)(1 - g) + k$$

is the expected complex dimension of $\overline{\mathfrak{M}}_{g,k}(Y, \beta)$, and $A_{\mathfrak{d}}(\overline{\mathfrak{M}}_{g,k}(Y, \beta))$ is the Chow group of $\mathfrak{d}$ -dimensional cycles in the moduli space $\overline{\mathfrak{M}}_{g,k}(Y, \beta)$.

The Gromov-Witten invariants are defined by using the virtual fundamental class $[\overline{\mathfrak{M}}_{g,k}(Y, \beta)]^{\text{vir}}$. Recall that an element $\alpha \in H^*(Y, \mathbb{C}) \stackrel{\text{def}}{=} \bigoplus_{j=0}^{2 \dim_{\mathbb{C}}(Y)} H^j(Y, \mathbb{C})$ is *homogeneous* if $\alpha \in H^j(Y, \mathbb{C})$ for some j ; in this case, we take $|\alpha| = j$. Let $\alpha_1, \dots, \alpha_k \in H^*(Y, \mathbb{C})$ such that every α_i is homogeneous and $\sum_{i=1}^k |\alpha_i| = 2\mathfrak{d}$. Then, we have the k -point Gromov-Witten invariant defined by

$$(2.3) \quad \langle \alpha_1, \dots, \alpha_k \rangle_{g,\beta} = \int_{[\overline{\mathfrak{M}}_{g,k}(Y, \beta)]^{\text{vir}}} ev_k^*(\alpha_1 \otimes \dots \otimes \alpha_k).$$

Next, we summarize certain properties concerning the virtual fundamental class. To begin with, we recall that *the excess dimension* is the difference between the dimension of $\overline{\mathfrak{M}}_{g,k}(Y, \beta)$ and the expected dimension $\mathfrak{d}$ in (2.2). For $0 \leq i < k$, use

$$(2.4) \quad f_{k,i} : \overline{\mathfrak{M}}_{g,k}(Y, \beta) \rightarrow \overline{\mathfrak{M}}_{g,i}(Y, \beta)$$

to stand for the forgetful map obtained by forgetting the last $k - i$ marked points and contracting all the unstable components. It is known that $f_{k,i}$ is flat when $\beta \neq 0$ and $0 \leq i < k$. The following can be found in [LT1, Beh, Get, C-K].

Proposition 2.1. *Let $\beta \in H_2(Y, \mathbb{Z})$ and $\beta \neq 0$. Let e be the excess dimension of $\overline{\mathfrak{M}}_{g,k}(Y, \beta)$, and $\mathfrak{M} \subset \overline{\mathfrak{M}}_{g,k}(Y, \beta)$ be a closed subscheme. Then,*

- (i) $[\overline{\mathfrak{M}}_{g,k}(Y, \beta)]^{\text{vir}} = (f_{k,0})^*[\overline{\mathfrak{M}}_{g,0}(Y, \beta)]^{\text{vir}}$;
- (ii) $[\overline{\mathfrak{M}}_{g,k}(Y, \beta)]^{\text{vir}} = c_e(R^1(f_{k+1,k})_*(ev_{k+1})^*T_Y)$ if $R^1(f_{k+1,k})_*(ev_{k+1})^*T_Y$ is a rank- e locally free sheaf over the moduli space $\overline{\mathfrak{M}}_{g,k}(Y, \beta)$;
- (iii) $[\overline{\mathfrak{M}}_{g,k}(Y, \beta)]^{\text{vir}}|_{\mathfrak{M}} = c_e((R^1(f_{k+1,k})_*(ev_{k+1})^*T_Y)|_{\mathfrak{M}})$ if there exists an open subset $\mathfrak{D}$ of $\overline{\mathfrak{M}}_{g,k}(Y, \beta)$ such that $\mathfrak{M} \subset \mathfrak{D}$ (i.e., $\mathfrak{D}$ is an open neighborhood of $\mathfrak{M}$) and the restriction $(R^1(f_{k+1,k})_*(ev_{k+1})^*T_Y)|_{\mathfrak{D}}$ is a rank- e locally free sheaf over $\mathfrak{D}$.

3. THE MODULI SPACE OF STABLE RANK-2 SHEAVES ON $\mathbb{P}^2$

3.1. Some basic facts of the moduli space. Throughout the rest of this paper, let $X = \mathbb{P}^2$ be the projective plane, and let ℓ be a line in X . For an integer n , let $\overline{\mathfrak{M}}(n)$ be the moduli space parametrizing all Gieseker-semistable rank-2 sheaves V over X with $c_1(V) = -\ell, c_2(V) = n$. Note that every such sheaf V is actually slope-stable and hence is Gieseker-stable. It is well known that, when $n \geq 1$, $\overline{\mathfrak{M}}(n)$ is nonempty, smooth, irreducible and rational with the expected dimension $4n - 4$; in addition, a universal sheaf over $\overline{\mathfrak{M}}(n) \times X$ exists. By Theorem 1 in [Mar], the cohomology groups $H^i(\overline{\mathfrak{M}}(n); \mathbb{Z})$ are torsion-free for all i and vanish for odd i . The

same is true of the homology groups $H_i(\overline{\mathfrak{M}}(n); \mathbb{Z})$. Let $b_i(\overline{\mathfrak{M}}(n))$ be the i -th Betti number of $\overline{\mathfrak{M}}(n)$, and put

$$p(\overline{\mathfrak{M}}(n); q) = \sum_{i=0}^{8n-8} b_i(\overline{\mathfrak{M}}(n)) q^i.$$

By Theorem 0.1 of [Yos], $p(\overline{\mathfrak{M}}(n); q)$ equals the coefficient of t^n of the series

$$(3.1) \quad \frac{1}{(q^2 - 1) \cdot \sum_{n \in \mathbb{Z}} q^{2n(2n-1)} t^{n^2}} \cdot \sum_{b \geq 0} \left(\frac{q^{2(b+1)(2b+1)}}{1 - q^{8(b+1)} t^{2b+1}} - \frac{q^{2b(2b+5)}}{1 - q^{8b} t^{2b+1}} \right) t^{(b+1)^2} \\ \cdot \prod_{d \geq 1} \frac{1}{(1 - q^{4d-2} t^d)^2 (1 - q^{4d} t^d)^2 (1 - q^{4d+2} t^d)^2}.$$

In the rest of the subsection, we review Strømme's work in [Str] and give a basis of $H^4(\overline{\mathfrak{M}}(n), \mathbb{Z})$ in terms of the classes from [Str]. A basis of $H_4(\overline{\mathfrak{M}}(n), \mathbb{Z})$ with geometric flavors will be constructed in §4.

Fix $n \geq 2$. Let $\mathcal{E}$ be a universal sheaf over $\overline{\mathfrak{M}}(n) \times X$, and let π_1 and π_2 be the two natural projections on $\overline{\mathfrak{M}}(n) \times X$. For $0 \leq k \leq 2$, define

$$\mathcal{A}_k = R^1 \pi_{1*}(\mathcal{E} \otimes \pi_2^* \mathcal{O}_X(-k\ell)).$$

For $V \in \overline{\mathfrak{M}}(n)$, denote $V \otimes \mathcal{O}_X(k\ell)$ by $V \otimes \mathcal{O}_X(k)$ or $V(k)$. Then,

$$(3.2) \quad h^0(X, V(-k)) = h^2(X, V(-k)) = 0 \quad \text{for } 0 \leq k \leq 2,$$

$$(3.3) \quad h^1(X, V) = h^1(X, V(-2)) = n - 1, \quad h^1(X, V(-1)) = n$$

by Proposition 1.5 in [Str]. It follows that the three sheaves $\mathcal{A}_0, \mathcal{A}_1, \mathcal{A}_2$ over $\overline{\mathfrak{M}}(n)$ are locally free of rank $n - 1, n, n - 1$ respectively.

Definition 3.1. Let $0 \leq k \leq 2$ and r_k be the rank of $\mathcal{A}_k$. Define

$$\begin{aligned} \epsilon &= c_1(\mathcal{A}_0) - c_1(\mathcal{A}_2), \\ \delta &= n \cdot c_1(\mathcal{A}_0) - (n - 1) \cdot c_1(\mathcal{A}_1), \\ \tau_k &= 2r_k \cdot c_2(\mathcal{A}_k) - (r_k - 1) \cdot c_1(\mathcal{A}_k)^2. \end{aligned}$$

Note that these classes ϵ, δ, τ_k are independent of the choices of the universal sheaf $\mathcal{E}$ over $\overline{\mathfrak{M}}(n) \times X$. Let $K_{\overline{\mathfrak{M}}(n)}$ be the canonical class of $\overline{\mathfrak{M}}(n)$, $\mathfrak{M}(n)$ be the open subset of $\overline{\mathfrak{M}}(n)$ parametrizing stable bundles, and

$$(3.4) \quad \mathfrak{B} = \overline{\mathfrak{M}}(n) - \mathfrak{M}(n).$$

By the theorem in [Str], $\text{Pic}(\overline{\mathfrak{M}}(n))$ is freely generated by ϵ and δ , and

$$(3.5) \quad K_{\overline{\mathfrak{M}}(n)} = -3\epsilon, \quad \mathfrak{B} = n\epsilon - 2\delta.$$

Also, $a\epsilon + b\delta$ is ample if and only if $a, b > 0$, and $\mathfrak{B}$ is irreducible and reduced.

It is known from [Bea, E-S, Mar] that the cohomology ring $H^*(\overline{\mathfrak{M}}(n); \mathbb{Z})$ is generated by the Chern classes of the bundles $\mathcal{A}_0, \mathcal{A}_1, \mathcal{A}_2$. It follows that $H^4(\overline{\mathfrak{M}}(n); \mathbb{Z})$ is the $\mathbb{Z}$ -linear span of the six integral classes:

$$(3.6) \quad \epsilon^2, \epsilon\delta, \delta^2, \tau_0, \tau_1, \tau_2.$$

By (3.1), the rank of $H^4(\overline{\mathfrak{M}}(n); \mathbb{Z})$ is 3 when $n = 2$, and is 6 when $n \geq 3$. Therefore, if $n \geq 3$, then a linear basis of $H^4(\overline{\mathfrak{M}}(n); \mathbb{Z})$ is given by the six classes in (3.6).

Remark 3.2. When $n = 2$, the rank of $H_4(\overline{\mathfrak{M}}(n); \mathbb{Z})$ is different from that of the case $n \geq 3$. Therefore the construction of a basis of $H_4(\overline{\mathfrak{M}}(n); \mathbb{Z})$ in §4 for $n \geq 3$ needs to be modified. However, there is a method to describe the moduli space $\overline{\mathfrak{M}}(2)$ using the moduli spaces of stable sheaves on the Hirzebruch surface $\mathbb{F}_1$ and chamber structures, which enable us to compute all the Gromov-Witten invariants of $\overline{\mathfrak{M}}(2)$ (instead of a special kind considered in this paper for $n \geq 3$). Since the method is different, the result for $n = 2$ will appear elsewhere.

3.2. The boundary and the Uhlenbeck compactification.

The quasi-projective variety $\mathfrak{M}(n)$ has a Uhlenbeck compactification

$$(3.7) \quad \overline{\mathfrak{U}}(n) = \coprod_{0 \leq i \leq n-1} \mathfrak{M}(n-i) \times \mathrm{Sym}^i(X)$$

according to [Uhl, LJ1, Mor]. Moreover, there exists a birational morphism, called the *Gieseker-Uhlenbeck morphism*,

$$(3.8) \quad \Psi : \overline{\mathfrak{M}}(n) \rightarrow \overline{\mathfrak{U}}(n)$$

sending $V \in \overline{\mathfrak{M}}(n)$ to the pair (V^{**}, η) , where V^{**} is the double-dual of V and

$$\eta = \sum_{x \in X} h^0(X, (V^{**}/V)_x) x.$$

Then the boundary divisor $\mathfrak{B}$ in (3.4) is contracted by Ψ to the codimension-2 subset $\coprod_{1 \leq i \leq n-1} \mathfrak{M}(n-i) \times \mathrm{Sym}^i(X)$ in $\overline{\mathfrak{U}}(n)$.

Let $\mathfrak{B}_* = \Psi^{-1}(\mathfrak{M}(n-1) \times X) \subset \overline{\mathfrak{M}}(n)$ which parametrizes all $V \in \overline{\mathfrak{M}}(n)$ sitting in exact sequences $0 \rightarrow V \rightarrow V_1 \rightarrow \mathcal{O}_x \rightarrow 0$ for some bundle $V_1 \in \mathfrak{M}(n-1)$ and some point $x \in X$. It is an open dense subset of the boundary divisor $\mathfrak{B}$.

To construct a universal sheaf over $\mathfrak{B}_* \times X$, let $\mathcal{E}_{n-1}^0$ be a universal sheaf over $\mathfrak{M}(n-1) \times X$. By [Q-Z], $\mathfrak{B}_* \cong \mathbb{P}(\mathcal{E}_{n-1}^0)$. For simplicity, write $\mathfrak{B}_* = \mathbb{P}(\mathcal{E}_{n-1}^0)$. Let

$$(3.9) \quad \pi : \mathfrak{B}_* = \mathbb{P}(\mathcal{E}_{n-1}^0) \rightarrow \mathfrak{M}(n-1) \times X$$

be the natural projection. Let Δ_X be the diagonal of $X \times X$. Consider the obvious isomorphism $\alpha : \mathfrak{M}(n-1) \times \Delta_X \rightarrow \mathfrak{M}(n-1) \times X$. Then, we have the isomorphisms:

$$(3.10) \quad \begin{aligned} (\pi \times \mathrm{Id}_X)^{-1}(\mathfrak{M}(n-1) \times \Delta_X) &\cong \mathbb{P}(\alpha^* \mathcal{E}_{n-1}^0), \\ \tilde{\pi}^* \mathcal{E}_{n-1}^0|_{(\pi \times \mathrm{Id}_X)^{-1}(\mathfrak{M}(n-1) \times \Delta_X)} &\cong \tilde{\alpha}^* \mathcal{E}_{n-1}^0, \end{aligned}$$

where $\tilde{\pi} : \mathfrak{B}_* \times X \rightarrow \mathfrak{M}(n-1) \times X$ is the composition of $\pi \times \mathrm{Id}_X$ and the map $\mathfrak{M}(n-1) \times X \times X \rightarrow \mathfrak{M}(n-1) \times X$ which denotes the projection to the product of the first and third factors, and $\tilde{\alpha} : \mathbb{P}(\alpha^* \mathcal{E}_{n-1}^0) \rightarrow \mathfrak{M}(n-1) \times X$ is the composition of the natural projection $\mathbb{P}(\alpha^* \mathcal{E}_{n-1}^0) \rightarrow \mathfrak{M}(n-1) \times \Delta_X$ and α . A universal sheaf $\mathcal{E}'$ over $\mathfrak{B}_* \times X$ sits in the exact sequence

$$(3.11) \quad 0 \rightarrow \mathcal{E}' \rightarrow \tilde{\pi}^* \mathcal{E}_{n-1}^0 \rightarrow \mathcal{O}_{(\pi \times \mathrm{Id}_X)^{-1}(\mathfrak{M}(n-1) \times \Delta_X)}(1) \rightarrow 0.$$

The following lemma will be used in later sections.

Lemma 3.3. *Let $\tilde{\pi}_i$ be the i -th projection on $\mathfrak{M}(n-1) \times X$, and let*

$$\mathfrak{D}_{n-1} = 2c_1(\mathcal{A}_{n-1,1}^0) - c_1(\mathcal{A}_{n-1,2}^0) - c_1(\mathcal{A}_{n-1,0}^0),$$

where $\mathcal{A}_{n-1,k}^0 = R^1 \tilde{\pi}_{1}(\mathcal{E}_{n-1}^0 \otimes \tilde{\pi}_2^* \mathcal{O}_X(-k\ell))$. Then, as divisors in $\mathfrak{B}_*$,*

$$(3.12) \quad \mathfrak{B}_*|_{\mathfrak{B}_*} = -2c_1(\mathcal{O}_{\mathfrak{B}_*}(1)) + (\tilde{\pi}_1 \circ \pi)^* \mathfrak{D}_{n-1} + 2(\tilde{\pi}_2 \circ \pi)^* \ell.$$

Proof. Since $\mathfrak{B}_*$ is open and dense in $\mathfrak{B}$, we see from (3.5) and Definition 3.1 that

$$(3.13) \quad \mathfrak{B}_*|_{\mathfrak{B}_*} = \mathfrak{B}|_{\mathfrak{B}_*} = (2(n-1)c_1(\mathcal{A}_1) - nc_1(\mathcal{A}_2) - nc_1(\mathcal{A}_0))|_{\mathfrak{B}_*}.$$

Next, let $0 \leq k \leq 2$, and let π_i be the i -th projection on $\mathfrak{B}_* \times X$. Then the restriction of π_1 to the subset $(\pi \times \text{Id}_X)^{-1}(\mathfrak{M}(n-1) \times \Delta_X)$ is an isomorphism from $(\pi \times \text{Id}_X)^{-1}(\mathfrak{M}(n-1) \times \Delta_X)$ to $\mathfrak{B}_*$. Thus tensoring (3.11) by $\pi_2^* \mathcal{O}_X(-k\ell)$ and then applying the functor π_{1*} , we get the exact sequence

$$\begin{aligned} 0 \rightarrow \mathcal{O}_{\mathfrak{B}_*}(1) \otimes (\tilde{\pi}_2 \circ \pi)^* \mathcal{O}_X(-k\ell) &\rightarrow R^1 \pi_{1*}(\mathcal{E}' \otimes \pi_2^* \mathcal{O}_X(-k\ell)) \\ &\rightarrow R^1 \pi_{1*}(\tilde{\pi}^* \mathcal{E}_{n-1}^0 \otimes \pi_2^* \mathcal{O}_X(-k\ell)) \rightarrow 0. \end{aligned}$$

Note that $\pi_2 = \tilde{\pi}_2 \circ \tilde{\pi}$. Also, $R^1 \pi_{1*} \tilde{\pi}^* \cong (\tilde{\pi}_1 \circ \pi)^* R^1 \tilde{\pi}_{1*}$ via the trivial base change:

$$\begin{array}{ccc} \mathfrak{B}_* \times X & \xrightarrow{\pi_1} & \mathfrak{B}_* \\ \downarrow \tilde{\pi} & & \downarrow \tilde{\pi}_1 \circ \pi \\ \mathfrak{M}(n-1) \times X & \xrightarrow{\tilde{\pi}_1} & \mathfrak{M}(n-1). \end{array}$$

Therefore, rewriting the 3rd term in the above exact sequence, we obtain

$$\begin{aligned} 0 \rightarrow \mathcal{O}_{\mathfrak{B}_*}(1) \otimes (\tilde{\pi}_2 \circ \pi)^* \mathcal{O}_X(-k\ell) &\rightarrow R^1 \pi_{1*}(\mathcal{E}' \otimes \pi_2^* \mathcal{O}_X(-k\ell)) \\ &\rightarrow (\tilde{\pi}_1 \circ \pi)^* \mathcal{A}_{n-1,k}^0 \rightarrow 0. \end{aligned}$$

So the first Chern class $c_1(R^1 \pi_{1*}(\mathcal{E}' \otimes \pi_2^* \mathcal{O}_X(-k\ell)))$ equals

$$(3.14) \quad c_1(\mathcal{O}_{\mathfrak{B}_*}(1)) - k(\tilde{\pi}_2 \circ \pi)^* \ell + (\tilde{\pi}_1 \circ \pi)^* c_1(\mathcal{A}_{n-1,k}^0),$$

and the second Chern class $c_2(R^1 \pi_{1*}(\mathcal{E}' \otimes \pi_2^* \mathcal{O}_X(-k\ell)))$ is equal to

$$(3.15) \quad (c_1(\mathcal{O}_{\mathfrak{B}_*}(1)) - k(\tilde{\pi}_2 \circ \pi)^* \ell) \cdot (\tilde{\pi}_1 \circ \pi)^* c_1(\mathcal{A}_{n-1,k}^0) + (\tilde{\pi}_1 \circ \pi)^* c_2(\mathcal{A}_{n-1,k}^0).$$

Finally, we conclude from (3.13) and (3.14) that

$$\mathfrak{B}_*|_{\mathfrak{B}_*} = -2c_1(\mathcal{O}_{\mathfrak{B}_*}(1)) + 2(\tilde{\pi}_2 \circ \pi)^* \ell + (\tilde{\pi}_1 \circ \pi)^* \mathfrak{D},$$

where

$$\mathfrak{D} = 2(n-1)c_1(\mathcal{A}_{n-1,1}^0) - nc_1(\mathcal{A}_{n-1,2}^0) - nc_1(\mathcal{A}_{n-1,0}^0).$$

Let $\epsilon_1, \delta_1 \in \text{Pic}(\overline{\mathfrak{M}}(n-1))$ be the counterparts of $\epsilon, \delta \in \text{Pic}(\overline{\mathfrak{M}}(n))$. By (3.5) and Definition 3.1,

$$\begin{aligned} 0 &= (\overline{\mathfrak{M}}(n-1) - \mathfrak{M}(n-1))|_{\mathfrak{M}(n-1)} \\ &= [(n-1)\epsilon_1 - 2\delta_1]|_{\mathfrak{M}(n-1)} \\ &= 2(n-2)c_1(\mathcal{A}_{n-1,1}^0) - (n-1)c_1(\mathcal{A}_{n-1,2}^0) - (n-1)c_1(\mathcal{A}_{n-1,0}^0). \end{aligned}$$

So $\mathfrak{B}_*|_{\mathfrak{B}_*} = -2c_1(\mathcal{O}_{\mathfrak{B}_*}(1)) + 2(\tilde{\pi}_2 \circ \pi)^* \ell + (\tilde{\pi}_1 \circ \pi)^* \mathfrak{D}_{n-1}$. $\square$

3.3. Curves in $\overline{\mathfrak{M}}(n)$.

We shall construct two curves in the Gieseker moduli space $\overline{\mathfrak{M}}(n)$ which freely generate the homology group $H_2(\overline{\mathfrak{M}}(n), \mathbb{Z})$. One such curve is a fiber $\mathfrak{f}$ of the morphism π from (3.9). The following is Lemma 3.2 in [Q-Z].

Lemma 3.4. *Let $N_{\mathfrak{f} \subset \overline{\mathfrak{M}}(n)}$ be the normal bundle of $\mathfrak{f}$ in $\overline{\mathfrak{M}}(n)$. Then,*

- (i) $\mathfrak{f} \cdot K_{\overline{\mathfrak{M}}(n)} = 0$ and $\mathfrak{f} \cdot \mathfrak{B} = -2$;
- (ii) $N_{\mathfrak{f} \subset \overline{\mathfrak{M}}(n)} \cong \mathcal{O}_{\mathfrak{f}}^{\oplus(4n-6)} \oplus \mathcal{O}_{\mathfrak{f}}(-2)$;
- (iii) $T_{\overline{\mathfrak{M}}(n)}|_{\mathfrak{f}} \cong \mathcal{O}_{\mathfrak{f}}^{\oplus(4n-6)} \oplus \mathcal{O}_{\mathfrak{f}}(-2) \oplus \mathcal{O}_{\mathfrak{f}}(2)$. $\square$

Next, we shall construct the other curve. Let $n \geq 3$, and let ξ consist of n distinct points in general position in $X = \mathbb{P}^2$. If V sits in a nontrivial extension

$$(3.16) \quad 0 \rightarrow \mathcal{O}_X(-1) \rightarrow V \rightarrow I_\xi \rightarrow 0,$$

then V is stable and hence $V \in \overline{\mathfrak{M}}(n)$. Moreover, since $H^0(X, I_\xi \otimes \mathcal{O}_X(1)) = 0$, the injection $\mathcal{O}_X(-1) \rightarrow V$ is unique up to scalars. It follows that

$$(3.17) \quad \mathfrak{E}_n = \mathbb{P}(\text{Ext}^1(I_\xi, \mathcal{O}_X(-1))) \cong \mathbb{P}^{n-1}$$

can be regarded as the subset of $\overline{\mathfrak{M}}(n)$ parametrizing all the sheaves $V \in \overline{\mathfrak{M}}(n)$ sitting in nontrivial extensions (3.16). A universal sheaf $\mathcal{E}'$ over $\mathfrak{E}_n \times X$ sits in

$$(3.18) \quad 0 \rightarrow \pi_2^* \mathcal{O}_X(-\ell) \rightarrow \mathcal{E}' \rightarrow \pi_1^* \mathcal{O}_{\mathfrak{E}_n}(-1) \otimes \pi_2^* I_\xi \rightarrow 0.$$

Tensoring by $\pi_2^* \mathcal{O}_X(-k\ell)$ and applying π_{1*} lead to the exact sequence:

$$0 \rightarrow R^1 \pi_{1*}(\mathcal{E}' \otimes \pi_2^* \mathcal{O}_X(-k\ell)) \rightarrow \mathcal{O}_{\mathfrak{E}_n}(-1)^{\oplus h^1(X, I_\xi(-k))} \rightarrow \mathcal{O}_{\mathfrak{E}_n}^{\oplus h^2(X, \mathcal{O}_X(-(k+1)\ell))} \rightarrow 0,$$

where $0 \leq k \leq 2$. An easy computation gives rise to the following:

$$\begin{aligned} c_1(R^1 \pi_{1*} \mathcal{E}') &= -(n-1) \cdot c_1(\mathcal{O}_{\mathfrak{E}_n}(1)), \\ c_1(R^1 \pi_{1*}(\mathcal{E}' \otimes \pi_2^* \mathcal{O}_X(-k\ell))) &= -n \cdot c_1(\mathcal{O}_{\mathfrak{E}_n}(1)), \end{aligned}$$

where $k = 1, 2$. It follows immediately from Definition 3.1 that

$$(3.19) \quad \epsilon|_{\mathfrak{E}_n} = c_1(\mathcal{O}_{\mathfrak{E}_n}(1)), \quad \delta|_{\mathfrak{E}_n} = 0.$$

Lemma 3.5. *Let $n \geq 3$ and $\mathfrak{l}$ be a line in the projective space $\mathfrak{E}_n$.*

- (i) *The homology group $H_2(\overline{\mathfrak{M}}(n); \mathbb{Z})$ is freely generated by $\mathfrak{f}$ and $\mathfrak{l}$.*
- (ii) *The class $a\mathfrak{f} + b\mathfrak{l} \in H_2(\overline{\mathfrak{M}}(n); \mathbb{Z})$ is effective if and only if $a, b \geq 0$.*
- (iii) *If $C \subset \overline{\mathfrak{M}}(n)$ is an irreducible curve contracted by the Gieseker-Uhlenbeck morphism Ψ , then $C = d\mathfrak{f} \in H_2(\overline{\mathfrak{M}}(n); \mathbb{Z})$ for some positive integer d .*

Proof. (i) Since $\overline{\mathfrak{M}}(n)$ is rational, $H^i(\overline{\mathfrak{M}}(n), \mathcal{O}_{\overline{\mathfrak{M}}(n)}) = 0$ for all $i \geq 1$. Hence $H^2(\overline{\mathfrak{M}}(n); \mathbb{Z}) \cong \text{Pic}(\overline{\mathfrak{M}}(n))$, and $H^2(\overline{\mathfrak{M}}(n); \mathbb{Z})$ is freely generated by ϵ and δ . By (3.19), $\mathfrak{l} \cdot \epsilon = 1$ and $\mathfrak{l} \cdot \delta = 0$. By Definition 3.1 and (3.14), $\mathfrak{f} \cdot \epsilon = 0$ and $\mathfrak{f} \cdot \delta = 1$. Since $H_2(\overline{\mathfrak{M}}(n); \mathbb{Z})$ is torsion-free, $H_2(\overline{\mathfrak{M}}(n); \mathbb{Z})$ is freely generated by $\mathfrak{f}$ and $\mathfrak{l}$.

(ii) Since $\mathfrak{f}$ and $\mathfrak{l}$ are effective, $a\mathfrak{f} + b\mathfrak{l}$ is effective if $a, b \geq 0$. Conversely, if $a\mathfrak{f} + b\mathfrak{l}$ is effective, then $a, b \geq 0$ since the divisor $c\epsilon + d\delta$ is ample if and only if $c, d > 0$.

(iii) By (ii), $C = d\mathfrak{f} + b\mathfrak{l} \in H_2(\overline{\mathfrak{M}}(n); \mathbb{Z})$, where d and b are nonnegative integers not both zero. Let L be a very ample divisor on $\overline{\mathfrak{U}}(n)$. Then, $\mathfrak{f} \cdot \Psi^* L = 0$ and $\mathfrak{l} \cdot \Psi^* L \geq 0$. Since $\Psi^* L$ is a nonzero divisor and $H_2(\overline{\mathfrak{M}}(n); \mathbb{Z})$ is freely generated by $\mathfrak{f}$ and $\mathfrak{l}$, we must have $\mathfrak{l} \cdot \Psi^* L > 0$. Thus, $C \cdot \Psi^* L = 0$ forces $b = 0$. $\square$

4. A BASIS OF $H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$

In this section, we assume $n \geq 3$. Then the integral homology group $H_4(\overline{\mathfrak{M}}(n); \mathbb{Z})$ is free of rank 6. In the following, we construct a basis $\{\Xi_1, \dots, \Xi_6\}$ for $H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$. This construction makes use of a result due to Hirschowitz and Hulek.

We review the results in [H-H], where complete rational curves were found in $\overline{\mathfrak{M}}(n)$. Let $n \geq 2$ and $\Gamma = \mathbb{P}^1$. Fix lines $\ell_1, \dots, \ell_n \subset X = \mathbb{P}^2$ in general position. For $1 \leq i \leq n$, let $\phi_i : \Gamma \rightarrow \ell_i$ be an isomorphism, and define $Y_i \subset \Gamma \times X$ to be the graph of ϕ_i . For generic choices of $\phi_1, \dots, \phi_n$, it was proved in [H-H] that $Y_1, \dots, Y_n$

are disjoint. Moreover, if $N_{Y/\Gamma \times X}$ denotes the normal bundle of $Y \stackrel{\text{def}}{=} \coprod_{i=1}^n Y_i$ in $\Gamma \times X$, then $\det(N_{Y/\Gamma \times X}) \cong \mathcal{O}_{\Gamma \times X}(2, 1)|_Y$. Therefore, the element

$$1 \in H^0(Y, \mathcal{O}_Y) \cong \text{Ext}^1(\mathcal{O}_{\Gamma \times X}(2, 0) \otimes I_Y, \mathcal{O}_{\Gamma \times X}(0, -1))$$

defines a rank-2 bundle $\tilde{\mathcal{E}}_n$ over $\Gamma \times X$ sitting in an exact sequence

$$(4.1) \quad 0 \rightarrow \mathcal{O}_{\Gamma \times X}(0, -1) \rightarrow \tilde{\mathcal{E}}_n \rightarrow \mathcal{O}_{\Gamma \times X}(2, 0) \otimes I_Y \rightarrow 0$$

such that $\tilde{\mathcal{E}}_n|_{\{p\} \times X} \in \mathfrak{M}(n)$ for all $p \in \Gamma$, and $\tilde{\mathcal{E}}_n$ induces a nonconstant morphism

$$\iota : \Gamma \rightarrow \mathfrak{M}(n).$$

Let $\tilde{\pi}_1$ and $\tilde{\pi}_2$ be the natural projections on $\Gamma \times X$. Let

$$(4.2) \quad \tilde{\mathcal{A}}_{n,k} = R^1 \tilde{\pi}_{1*}(\tilde{\mathcal{E}}_n \otimes \tilde{\pi}_2^* \mathcal{O}_X(-k\ell)).$$

By Lemma 3.5 in [H-H], the degrees of the bundles $\tilde{\mathcal{A}}_{n,k}$ are

$$(4.3) \quad a_{n,k} = \begin{cases} 2n-2 & \text{if } k=0, \\ n & \text{if } k=1, \\ 0 & \text{if } k=2. \end{cases}$$

Remark 4.1. (i) Let $n \geq 2$. By (3.3), the condition in Corollarie (6.9.9) of [Gro] is satisfied. Hence the base-change theorem of the first direct image holds for every projective morphism to $\overline{\mathfrak{M}}(n)$ and for the sheaf $\mathcal{E} \otimes \pi_2^* \mathcal{O}_X(-k\ell)$ with $0 \leq k \leq 2$. For a finite morphism φ from Y onto a subvariety of $\overline{\mathfrak{M}}(n)$, the intersection numbers of $\varphi(Y)$ with $\epsilon^2, \epsilon \cdot \delta, \delta^2, \tau_0, \tau_1, \tau_2$ can be computed on Y , via the projection formula, by pulling back the Chern classes of the first direct images of the sheaf $\mathcal{E} \otimes \pi_2^* \mathcal{O}_X(-k\ell)$ with $0 \leq k \leq 2$.

(ii) Let $n \geq 3$. Then $(n-1) \geq 2$. In Subsection 4.2, we will construct a surface $Y = \pi^{-1}(\Gamma \times \{y\})$ admitting a finite morphism onto a surface Ξ_3 of $\overline{\mathfrak{M}}(n)$. This finite morphism factors through the nonconstant morphism $\iota : \Gamma \rightarrow \mathfrak{M}(n-1)$. Applying (i), we may assume for convenience that $\Gamma \subset \mathfrak{M}(n-1)$ and consequently $Y = \Xi_3 \subset \overline{\mathfrak{M}}(n)$ (in fact, it can be proved that when $n \geq 5$, the morphism $\iota : \Gamma \rightarrow \mathfrak{M}(n-1)$ is injective). A similar discussion works for Ξ_4 .

4.1. The homology classes $\Xi_1, \Xi_2 \in H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$. Let $n \geq 3$, and assume $\Gamma \subset \mathfrak{M}(n-1)$ as pointed out in Remark 4.1 (ii). Let $\mathbb{P} = \mathbb{P}(\tilde{\mathcal{E}}_{n-1}) = \pi^{-1}(\Gamma \times X) \subset \mathfrak{B}_*$. Then $\mathbb{P}$ parametrizes all the sheaves $V \in \overline{\mathfrak{M}}(n)$ sitting in

$$0 \rightarrow V \rightarrow \tilde{\mathcal{E}}_{n-1}|_{\{p\} \times X} \rightarrow \mathcal{O}_x \rightarrow 0$$

for some $p \in \Gamma$ and $x \in X$. We still use π to denote the natural projection

$$\pi : \mathbb{P} \rightarrow \Gamma \times X.$$

By (3.11), a universal sheaf $\tilde{\mathcal{E}}$ over $\mathbb{P} \times X$ sits in the exact sequence

$$(4.4) \quad 0 \rightarrow \tilde{\mathcal{E}} \rightarrow \tilde{\pi}^* \tilde{\mathcal{E}}_{n-1} \rightarrow \mathcal{O}_{(\pi \times \text{Id}_X)^{-1}(\mathbb{P} \times \Delta_X)}(1) \rightarrow 0,$$

where $\tilde{\pi}$ is the composition of $\pi \times \text{Id}_X : \mathbb{P} \times X \rightarrow \Gamma \times X \times X$ and the projection $\Gamma \times X \times X \rightarrow \Gamma \times X$ to the first and third factors.

Let π_i (respectively, $\tilde{\pi}_i$) be the natural projections on $\mathbb{P} \times X$ (respectively, $\Gamma \times X$). By (3.14) and (3.15), $c_1(R^1 \pi_{1*}(\tilde{\mathcal{E}} \otimes \pi_2^* \mathcal{O}_X(-k\ell)))$ equals

$$(4.5) \quad c_1(\mathcal{O}_{\mathbb{P}}(1)) - k(\tilde{\pi}_2 \circ \pi)^* \ell + (\tilde{\pi}_1 \circ \pi)^* c_1(\tilde{\mathcal{A}}_{n-1,k}),$$

where $\tilde{\mathcal{A}}_{n-1,k}$ is defined in (4.2), and $c_2(R^1\pi_{1*}(\tilde{\mathcal{E}} \otimes \pi_2^* \mathcal{O}_X(-k\ell)))$ equals

$$(4.6) \quad (c_1(\mathcal{O}_{\mathbb{P}}(1)) - k(\tilde{\pi}_2 \circ \pi)^* \ell) \cdot (\tilde{\pi}_1 \circ \pi)^* c_1(\tilde{\mathcal{A}}_{n-1,k}).$$

In addition, we conclude from (4.1) that

$$(4.7) \quad \begin{aligned} c_1(\mathcal{O}_{\mathbb{P}}(1))^2 &= \pi^* c_1(\tilde{\mathcal{E}}_{n-1}) \cdot c_1(\mathcal{O}_{\mathbb{P}}(1)) - \pi^* c_2(\tilde{\mathcal{E}}_{n-1}) \\ &= \pi^*(2, -1) \cdot c_1(\mathcal{O}_{\mathbb{P}}(1)) - \pi^* \left[(2, 0) \cdot (0, -1) + \sum_{i=1}^{n-1} Y_i \right]. \end{aligned}$$

Fix a point $p \in \Gamma$ and let $V_1 = \tilde{\mathcal{E}}_{n-1}|_{\{p\} \times X}$. Consider $\mathbb{P}(V_1) \subset \mathbb{P} \subset \overline{\mathfrak{M}}(n)$. By Definition 3.1, (4.5), (4.6) and (4.7), we obtain

$$(4.8) \quad \begin{aligned} \epsilon|_{\mathbb{P}(V_1)} &= 2\pi^* \ell, \\ \delta|_{\mathbb{P}(V_1)} &= c_1(\mathcal{O}_{\mathbb{P}(V_1)}(1)) + (n-1)\pi^* \ell, \\ \tau_k|_{\mathbb{P}(V_1)} &= -(r_k - 1)(c_1(\mathcal{O}_{\mathbb{P}(V_1)}(1)) - k\pi^* \ell)^2, \\ c_1(\mathcal{O}_{\mathbb{P}(V_1)}(1))^2 &= -\pi^* \ell \cdot c_1(\mathcal{O}_{\mathbb{P}(V_1)}(1)) - (n-1)\pi^* x, \end{aligned}$$

where by abusing notation, π denotes the natural projection $\mathbb{P}(V_1) \rightarrow X$.

Define $\Xi_1 \in H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$ to be the surface $\pi^{-1}(\{p\} \times \ell) \subset \mathbb{P}(V_1) \subset \mathbb{P} \subset \overline{\mathfrak{M}}(n)$, where $\ell \subset X = \mathbb{P}^2$ is a fixed line. Then we have

$$(4.9) \quad \begin{aligned} \epsilon^2 \cdot \Xi_1 &= 0, \quad \epsilon \cdot \delta \cdot \Xi_1 = 2, \quad \delta^2 \cdot \Xi_1 = 2n - 3, \\ \tau_0 \cdot \Xi_1 &= n - 2, \quad \tau_1 \cdot \Xi_1 = 3n - 3, \quad \tau_2 \cdot \Xi_1 = 5n - 10. \end{aligned}$$

Next, define $\Xi_2 \in H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$ to be $c_1(\mathcal{O}_{\mathbb{P}(V_1)}(1))$ regarded as a 2-dimensional cycle in $\overline{\mathfrak{M}}(n)$ via the inclusion $\mathbb{P}(V_1) \subset \overline{\mathfrak{M}}(n)$. Then

$$(4.10) \quad \begin{aligned} \epsilon^2 \cdot \Xi_2 &= 4, \quad \epsilon \cdot \delta \cdot \Xi_2 = 2n - 4, \quad \delta^2 \cdot \Xi_2 = n^2 - 5n + 5, \\ \tau_0 \cdot \Xi_2 &= (n - 2)^2, \quad \tau_1 \cdot \Xi_2 = (n - 1)(n - 5), \quad \tau_2 \cdot \Xi_2 = (n - 2)(n - 10). \end{aligned}$$

4.2. The homology classes $\Xi_3, \Xi_4 \in H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$.

Fix a line $\ell \subset X = \mathbb{P}^2$. Let $W = \pi^{-1}(\Gamma \times \ell) \subset \mathbb{P} \subset \overline{\mathfrak{M}}(n)$, where $\Gamma, \mathbb{P}$ and π are from the previous subsection. By Definition 3.1, (4.3), (4.5) and (4.6),

$$\begin{aligned} \epsilon|_W &= \pi^*(2n - 4, 2), \\ \delta|_W &= c_1(\mathcal{O}_W(1)) + \pi^*(n^2 - 2n - 1, n - 1), \\ \tau_k|_W &= \pi^*(2a_{n-1,k} - 2(r_k - 1), (2k + 1)(r_k - 1)) \cdot c_1(\mathcal{O}_W(1)) \\ &\quad + ((r_k - 1)(n - 3) - 2ka_{n-1,k}) \pi^* x, \end{aligned}$$

where $x \in \Gamma \times \ell$ is a fixed point and by abusing notation, $\pi : W \rightarrow \Gamma \times \ell \cong \mathbb{P}^1 \times \mathbb{P}^1$ stands for the natural projection. In addition, by (4.7), we conclude that

$$(4.11) \quad c_1(\mathcal{O}_W(1))^2 = \pi^*(2, -1) c_1(\mathcal{O}_W(1)) - (n - 3) \pi^* x.$$

Define $\Xi_3 \in H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$ to be the surface $\pi^{-1}(\Gamma \times \{y\}) \subset W \subset \overline{\mathfrak{M}}(n)$, where $y \in \ell$ is a fixed point. A straightforward computation shows that

$$(4.12) \quad \begin{aligned} \epsilon^2 \cdot \Xi_3 &= 0, \quad \epsilon \cdot \delta \cdot \Xi_3 = 2n - 4, \quad \delta^2 \cdot \Xi_3 = 2n^2 - 4n, \\ \tau_0 \cdot \Xi_3 &= 2n - 4, \quad \tau_1 \cdot \Xi_3 = 0, \quad \tau_2 \cdot \Xi_3 = -2n + 4. \end{aligned}$$

Define $\Xi_4 \in H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$ to be $c_1(\mathcal{O}_W(1))$ regarded as a 2-dimensional cycle in $\overline{\mathfrak{M}}(n)$ via the inclusion $W \subset \overline{\mathfrak{M}}(n)$. Then we have

$$(4.13) \quad \begin{aligned} \epsilon^2 \cdot \Xi_4 &= 8(n-2), \quad \epsilon \cdot \delta \cdot \Xi_4 = 4n^2 - 12n + 10, \quad \delta^2 \cdot \Xi_4 = 2n^3 - 8n^2 + 9n - 1, \\ \tau_0 \cdot \Xi_4 &= n^2 - 5n + 6, \quad \tau_1 \cdot \Xi_4 = n^2 - 1, \quad \tau_2 \cdot \Xi_4 = (n-2)(n+9). \end{aligned}$$

4.3. The homology class $\Xi_5 \in H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$.

Since $n \geq 3$, the moduli space $\mathfrak{M}(n-2)$ is nonempty. Fix a vector bundle $V_2 \in \mathfrak{M}(n-2)$ and two distinct points $x_1, x_2 \in X$. Let $\Xi_5 \subset \overline{\mathfrak{M}}(n)$ parametrize all the sheaves $V \in \overline{\mathfrak{M}}(n)$ sitting in exact sequences:

$$0 \rightarrow V \rightarrow V_2 \rightarrow \mathcal{O}_{x_1} \oplus \mathcal{O}_{x_2} \rightarrow 0.$$

Then we have the isomorphisms $\Xi_5 \cong \mathbb{P}(V|_{x_1}) \times \mathbb{P}(V|_{x_2}) \cong \mathbb{P}^1 \times \mathbb{P}^1$. Moreover, a universal sheaf $\mathcal{E}'$ over $\Xi_5 \times X$ sits in the exact sequence:

$$0 \rightarrow \mathcal{E}' \rightarrow \pi_2^* V_2 \rightarrow \tilde{\pi}_1^* \mathcal{O}_{\mathbb{P}^1}(1)|_{\mathbb{P}^1 \times \mathbb{P}^1 \times \{x_1\}} \oplus \tilde{\pi}_2^* \mathcal{O}_{\mathbb{P}^1}(1)|_{\mathbb{P}^1 \times \mathbb{P}^1 \times \{x_2\}} \rightarrow 0,$$

where π_i and $\tilde{\pi}_i$ denote the i -th projection on $\Xi_5 \times X$ and $\mathbb{P}^1 \times \mathbb{P}^1 \times X$, respectively. Tensoring the above exact sequence by $\pi_2^* \mathcal{O}_X(-k\ell)$ and applying π_{1*} yield

$$0 \rightarrow \mathcal{O}_{\Xi_5}(1, 0) \oplus \mathcal{O}_{\Xi_5}(0, 1) \rightarrow R^1 \pi_{1*}(\mathcal{E}' \otimes \pi_2^* \mathcal{O}_X(-k\ell)) \rightarrow \mathcal{O}_{\Xi_5}^{\oplus h^1(X, V_2(-k))} \rightarrow 0,$$

where $0 \leq k \leq 2$. Therefore $c_1(R^1 \pi_{1*}(\mathcal{E}' \otimes \pi_2^* \mathcal{O}_X(-k\ell))) = c_1(\mathcal{O}_{\Xi_5}(1, 1))$ and $c_2(R^1 \pi_{1*}(\mathcal{E}' \otimes \pi_2^* \mathcal{O}_X(-k\ell))) = 1$ for $0 \leq k \leq 2$. It follows from Definition 3.1 that

$$\epsilon|_{\Xi_5} = 0, \quad \delta|_{\Xi_5} = c_1(\mathcal{O}_{\Xi_5}(1, 1)).$$

Regarding $\Xi_5 \in H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$, we obtain the intersection numbers on $\overline{\mathfrak{M}}(n)$:

$$(4.14) \quad \epsilon^2 \cdot \Xi_5 = \epsilon \cdot \delta \cdot \Xi_5 = 0, \quad \delta^2 \cdot \Xi_5 = 2, \quad \tau_0 \cdot \Xi_5 = \tau_1 \cdot \Xi_5 = \tau_2 \cdot \Xi_5 = 2.$$

4.4. The surface Ξ_6 in $\overline{\mathfrak{M}}(n)$.

Fix a vector bundle $V_2 \in \mathfrak{M}(n-2)$ and a point $x \in X$. Fix a trivialization of V_2 in an open neighborhood $\mathcal{O}_x$ of x . Let $X^{[2]}$ be the Hilbert scheme parametrizing the length-2 closed subschemes of X , and let

$$M_2(x) = \{\xi \in X^{[2]} \mid \text{Supp}(\xi) = \{x\}\} \cong \mathbb{P}^1.$$

For $\xi \in M_2(x)$, let $\iota_\xi : \mathcal{O}_X \rightarrow \mathcal{O}_\xi$ be the natural quotient morphism.

Let $\Xi_6 = \mathbb{P}^1 \times M_2(x)$ be the subset of $\overline{\mathfrak{M}}(n)$ parametrizing all the sheaves $V \in \overline{\mathfrak{M}}(n)$ sitting in extensions of the form

$$0 \longrightarrow V \longrightarrow V_2 \xrightarrow{(a,b)} \mathcal{O}_\xi \longrightarrow 0,$$

where $\xi \in M_2(x)$, and for $(a, b) \in \mathbb{P}^1$, the map $V_2 \xrightarrow{(a,b)} \mathcal{O}_\xi$ denotes the composition:

$$V_2 \longrightarrow V_2|_{\mathcal{O}_x} \cong \mathcal{O}_{\mathcal{O}_x}^{\oplus 2} \xrightarrow{(a\iota_\xi, b\iota_\xi)} \mathcal{O}_\xi.$$

Next, we construct a universal sheaf over $\Xi_6 \times X = \mathbb{P}^1 \times M_2(x) \times X$. Let $\pi_{i_1, \dots, i_m}$ denote the projection of $\mathbb{P}^1 \times M_2(x) \times X$ to the product of the i_1 -th, $\dots$, i_m -th factors. Over $\Xi_6 = \mathbb{P}^1 \times M_2(x)$, there is a tautological surjection $\mathcal{O}_{\Xi_6}^{\oplus 2} \rightarrow \mathcal{O}_{\Xi_6}(1, 0) \rightarrow 0$. This pulls back to a surjection over $\mathbb{P}^1 \times M_2(x) \times X$:

$$(4.15) \quad \mathcal{O}_{\Xi_6 \times X}^{\oplus 2} \rightarrow \pi_1^* \mathcal{O}_{\mathbb{P}^1}(1) \rightarrow 0.$$

Let $\mathcal{Z}_2(x)$ be the universal codimension-2 subscheme in $M_2(x) \times X$, and let $\mathcal{O}_{M_2(x) \times X} \rightarrow \mathcal{O}_{\mathcal{Z}_2(x)} \rightarrow 0$ be the natural surjection. Pulling back to $\mathbb{P}^1 \times M_2(x) \times X$ yields a surjection:

$$(4.16) \quad \mathcal{O}_{\Xi_6 \times X} \rightarrow \pi_{2,3}^* \mathcal{O}_{\mathcal{Z}_2(x)} \rightarrow 0.$$

Tensoring (4.15) and (4.16), we obtain a surjection

$$(4.17) \quad \mathcal{O}_{\Xi_6 \times X}^{\oplus 2} \rightarrow \pi_1^* \mathcal{O}_{\mathbb{P}^1}(1) \otimes \pi_{2,3}^* \mathcal{O}_{\mathcal{Z}_2(x)} \rightarrow 0.$$

Since $\mathcal{Z}_2(x)$ is supported on $M_2(x) \times \{x\}$, in view of the trivialization of V_2 near x , (4.17) induces a surjection $\pi_3^* V_2 \rightarrow \pi_1^* \mathcal{O}_{\mathbb{P}^1}(1) \otimes \pi_{2,3}^* \mathcal{O}_{\mathcal{Z}_2(x)} \rightarrow 0$. The kernel $\mathcal{E}'$ of the map $\pi_3^* V_2 \rightarrow \pi_1^* \mathcal{O}_{\mathbb{P}^1}(1) \otimes \pi_{2,3}^* \mathcal{O}_{\mathcal{Z}_2(x)}$ is a universal sheaf over $\Xi_6 \times X$:

$$0 \rightarrow \mathcal{E}' \rightarrow \pi_3^* V_2 \rightarrow \pi_1^* \mathcal{O}_{\mathbb{P}^1}(1) \otimes \pi_{2,3}^* \mathcal{O}_{\mathcal{Z}_2(x)} \rightarrow 0.$$

It follows that for $0 \leq k \leq 2$, we have the exact sequence over Ξ_6 :

$$\begin{aligned} 0 \rightarrow \tilde{\pi}_1^* \mathcal{O}_{\mathbb{P}^1}(1) \otimes \tilde{\pi}_2^* (\mathcal{O}_X^{[2]}|_{M_2(x)}) &\rightarrow R^1(\pi_{1,2})_*(\mathcal{E}' \otimes \pi_3^* \mathcal{O}_X(-k\ell)) \\ &\rightarrow \mathcal{O}_{\Xi_6}^{\oplus h^1(X, V_2(-k))} \rightarrow 0, \end{aligned}$$

where $\tilde{\pi}_1$ and $\tilde{\pi}_2$ are the two natural projections on $\Xi_6 = \mathbb{P}^1 \times M_2(x)$, and $\mathcal{O}_X^{[2]}$ is the tautological rank-2 bundle over the Hilbert scheme $X^{[2]}$ whose fiber at a point $\xi \in X^{[2]}$ is the space $H^0(X, \mathcal{O}_\xi)$. It is well known that $-2c_1(\mathcal{O}_X^{[2]}) = M_2(X)$, where $M_2(X) \subset X^{[2]}$ consists of all the elements $\xi \in X^{[2]}$ such that $|\text{Supp}(\xi)| = 1$. Since $M_2(X) \cdot M_2(x) = -2$, we conclude from the above exact sequence that

$$\begin{aligned} c_1(R^1(\pi_{1,2})_*(\mathcal{E}' \otimes \pi_3^* \mathcal{O}_X(-k\ell))) &= (2, 1) \in \text{Pic}(\Xi_6) \cong \text{Pic}(\mathbb{P}^1 \times \mathbb{P}^1), \\ c_2(R^1(\pi_{1,2})_*(\mathcal{E}' \otimes \pi_3^* \mathcal{O}_X(-k\ell))) &= 1. \end{aligned}$$

By Definition 3.1, we get the six intersection numbers on $\overline{\mathfrak{M}}(n)$:

$$(4.18) \quad \begin{aligned} \epsilon^2 \cdot \Xi_6 &= 0, & \epsilon \cdot \delta \cdot \Xi_6 &= 0, & \delta^2 \cdot \Xi_6 &= 4, \\ \tau_0 \cdot \Xi_6 &= -2n + 6, & \tau_1 \cdot \Xi_6 &= -2n + 4, & \tau_2 \cdot \Xi_6 &= -2n + 6. \end{aligned}$$

Now we can summarize the above in the following proposition.

Proposition 4.2. *Let $n \geq 3$. Then $\{\Xi_1, \dots, \Xi_6\}$ is a linear basis of $H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$.*

Proof. Note from (4.9), (4.1), (4.12), (4.13), (4.14) and (4.18) that the intersection matrix between the classes $\Xi_1, \dots, \Xi_6$ and the classes $\epsilon^2, \epsilon \cdot \delta, \delta^2, \tau_0, \tau_1, \tau_2$ has a nonzero determinant. Since $H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$ has dimension 6, our result follows. $\square$

5. THE RESTRICTION OF THE OBSTRUCTION SHEAF ON CERTAIN OPEN SUBSETS

From the previous section, we see that, if we let $\overline{\mathfrak{M}}_*(n) = \mathfrak{B}_* \cup \mathfrak{M}(n)$, then the classes $\Xi_1, \Xi_2, \Xi_3, \Xi_4$ lie in $\overline{\mathfrak{M}}_*(n)$ while Ξ_5 and Ξ_6 lie in the complement $\overline{\mathfrak{M}}(n) - \overline{\mathfrak{M}}_*(n)$. Since a stable map $[\mu: (D, p) \rightarrow \overline{\mathfrak{M}}(n)]$ in $ev_1^{-1}(\Xi_i) \subset \overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f})$ has $\mu(p) \in \Xi_i$ for $1 \leq i \leq 4$, we have $\mu(D) \subset \mathfrak{B}_*$. In this section, we use the geometric construction of $\mathfrak{B}_*$ in Subsect. 3.2 to effectively compute the virtual cycle restricted to $ev_1^{-1}(\mathfrak{B}_*)$. The result will be used to compute the Gromov-Witten invariants $\langle \alpha \rangle_{0, d\mathfrak{f}}$ when α is dual to the classes $\Xi_1, \Xi_2, \Xi_3, \Xi_4$.

Fix $d \geq 1$. Consider the open subset $\mathfrak{D}_0$ of $\overline{\mathfrak{M}}_{0,0}(\overline{\mathfrak{M}}(n), d\mathfrak{f})$ consisting of stable maps $[\mu: D \rightarrow \overline{\mathfrak{M}}(n)]$ such that $\mu(D) \subset \overline{\mathfrak{M}}_*(n)$. Similarly, take the open subset $\mathfrak{D}_1$

of $\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f})$ consisting of stable maps $[\mu: (D; p) \rightarrow \overline{\mathfrak{M}}(n)]$ such that $\mu(D) \subset \overline{\mathfrak{M}}_*(n)$. Clearly $\mathfrak{D}_1 = f_{1,0}^{-1}(\mathfrak{D}_0)$. Let $[\mu: (D; p) \rightarrow \overline{\mathfrak{M}}(n)] \in \mathfrak{D}_1$. Since $\mu(D) = d\mathfrak{f}$ in $H_2(\overline{\mathfrak{M}}(n); \mathbb{Z})$, $\mu(D) \subset \mathfrak{B}_*$ and $\mu(D)$ is a fiber of the projection π . Moreover, the composition $\Psi \circ ev_1$ sends the stable map $[\mu: (D; p) \rightarrow \overline{\mathfrak{M}}(n)]$ to a point in $\mathfrak{M}(n-1) \times X \subset \overline{\mathfrak{U}}(n)$, which is independent of the marked point p on D . Hence ev_1 induces a morphism ϕ from $\mathfrak{D}_0$ to $\mathfrak{M}(n-1) \times X$. Putting $\tilde{ev}_1 = ev_1|_{\mathfrak{D}_1}$ and $\tilde{f}_{1,0} = f_{1,0}|_{\mathfrak{D}_1}$, we have the following commutative diagram:

$$(5.1) \quad \begin{array}{ccccc} \mathfrak{D}_1 & \xrightarrow{\tilde{ev}_1} & \mathfrak{B}_* & & \\ \downarrow \tilde{f}_{1,0} & & \downarrow \pi & & \\ \mathfrak{D}_0 & \xrightarrow{\phi} & \mathfrak{M}(n-1) \times X & \xrightarrow{\tilde{\pi}_3} & X. \end{array}$$

Note that the fiber $\phi^{-1}(V_1, x)$ over a point $(V_1, x) \in \mathfrak{M}(n-1) \times X$ is simply

$$\overline{\mathfrak{M}}_{0,0}(\pi^{-1}(V_1, x), d[\pi^{-1}(V_1, x)]),$$

which is isomorphic to $\overline{\mathfrak{M}}_{0,0}(\mathbb{P}^1, d[\mathbb{P}^1])$ via the isomorphism $\pi^{-1}(V_1, x) \cong \mathbb{P}^1$. Hence the complex dimension of the open subset $\mathfrak{D}_0 \subset \overline{\mathfrak{M}}_{0,0}(\overline{\mathfrak{M}}(n), d\mathfrak{f})$ is equal to

$$\begin{aligned} & \dim \overline{\mathfrak{M}}_{0,0}(\mathbb{P}^1, d[\mathbb{P}^1]) + \dim \mathfrak{M}(n-1) + \dim X \\ &= (2d-2) + [4(n-1)-4] + 2 \\ &= 2d + 4n - 8. \end{aligned}$$

Since $K_{\overline{\mathfrak{M}}(n)} \cdot d\mathfrak{f} = 0$, the expected dimension of $\mathfrak{D}_0 \subset \overline{\mathfrak{M}}_{0,0}(\overline{\mathfrak{M}}(n), d\mathfrak{f})$ is $4n-7$ by (2.2). Hence the excess dimension of $\mathfrak{D}_0$ is $e = 2d-1$.

Let $\mathcal{V}$ be the restriction of $R^1(f_{1,0})_*(ev_1)^*T_{\overline{\mathfrak{M}}(n)}$ to $\mathfrak{D}_0$.

Lemma 5.1. (i) *The sheaf $\mathcal{V}$ is locally free of rank $2d-1$;*

(ii) $\mathcal{V} \cong R^1(\tilde{f}_{1,0})_*(\tilde{ev}_1)^*\mathcal{O}_{\mathfrak{B}_*}(\mathfrak{B}_*)$.

Proof. (i) Take a stable map $u = [\mu: D \rightarrow \overline{\mathfrak{M}}(n)]$ in $\mathfrak{D}_0$, and consider

$$H^1(f_{1,0}^{-1}(u), (ev_1^*T_{\overline{\mathfrak{M}}(n)})|_{f_{1,0}^{-1}(u)}) \cong H^1(D, \mu^*T_{\overline{\mathfrak{M}}(n)}).$$

Since $\mu(D) \cong \mathbb{P}^1$ is a fiber of the projection π , we see from Lemma 3.4 (iii) that

$$T_{\overline{\mathfrak{M}}(n)}|_{\mu(D)} = \mathcal{O}_{\mu(D)}^{\oplus(4n-6)} \oplus \mathcal{O}_{\mu(D)}(-2) \oplus \mathcal{O}_{\mu(D)}(2).$$

Thus $H^1(D, \mu^*T_{\overline{\mathfrak{M}}(n)}) \cong H^1(D, \mu^*\mathcal{O}_{\mu(D)}(-2))$ whose dimension equals the excess dimension $e = 2d-1$. Therefore, the restriction $\mathcal{V}$ of $R^1(f_{1,0})_*(ev_1)^*T_{\overline{\mathfrak{M}}(n)}$ to $\mathfrak{D}_0$ is a locally free sheaf of rank $2d-1$.

(ii) Since $ev_1(\mathfrak{D}_1) \subset \mathfrak{B}_*$, we have $((ev_1)^*T_{\overline{\mathfrak{M}}(n)})|_{\mathfrak{D}_1} = (\tilde{ev}_1)^*(T_{\overline{\mathfrak{M}}_*(n)}|_{\mathfrak{B}_*})$ and

$$\begin{aligned} \mathcal{V} &= (R^1(f_{1,0})_*(ev_1)^*T_{\overline{\mathfrak{M}}(n)})|_{\mathfrak{D}_0} = R^1(\tilde{f}_{1,0})_*(((ev_1)^*T_{\overline{\mathfrak{M}}(n)})|_{\mathfrak{D}_1}) \\ &= R^1(\tilde{f}_{1,0})_*(\tilde{ev}_1)^*(T_{\overline{\mathfrak{M}}(n)}|_{\mathfrak{B}_*}). \end{aligned}$$

Since $\mathfrak{B}_*$ is smooth of codimension 1 in $\overline{\mathfrak{M}}(n)$, we obtain the exact sequence

$$(5.2) \quad 0 \rightarrow T_{\mathfrak{B}_*} \rightarrow T_{\overline{\mathfrak{M}}(n)}|_{\mathfrak{B}_*} \rightarrow \mathcal{O}_{\mathfrak{B}_*}(\mathfrak{B}_*) \rightarrow 0.$$

Applying $(\tilde{ev}_1)^*$ and $(\tilde{f}_{1,0})_*$ to the exact sequence (5.2), we get

$$(5.3) \quad R^1(\tilde{f}_{1,0})_*(\tilde{ev}_1)^*T_{\mathfrak{B}_*} \rightarrow \mathcal{V} \rightarrow R^1(\tilde{f}_{1,0})_*(\tilde{ev}_1)^*\mathcal{O}_{\mathfrak{B}_*}(\mathfrak{B}_*) \rightarrow 0,$$

where we have used $R^2(\tilde{f}_{1,0})_*(\tilde{ev}_1)^*T_{\mathfrak{B}_*} = 0$ since $\tilde{f}_{1,0}$ is of relative dimension 1.

If $[\mu: D \rightarrow \overline{\mathfrak{M}}(n)]$ is a stable map in $\mathfrak{D}_0$, then $\mu(D)$ is a fiber of the projection π in (3.9). Hence the normal bundle of $\mu(D)$ in $\mathfrak{B}_*$ is trivial. Therefore we have

$$T_{\mathfrak{B}_*}|_{\mu(D)} \cong \mathcal{O}_{\mu(D)}^{\oplus(4n-6)} \oplus \mathcal{O}_{\mu(D)}(2),$$

and

$$H^1(D, \mu^* T_{\mathfrak{B}_*}) \cong H^1(D, \mu^*(\mathcal{O}_{\mu(D)}^{\oplus(4n-6)} \oplus \mathcal{O}_{\mu(D)}(2))) = 0.$$

It follows that $R^1(\tilde{f}_{1,0})_*(\tilde{ev}_1)^* T_{\mathfrak{B}_*} = 0$. $\square$

Proposition 5.2. *Put $\mathcal{L} = \det(\mathcal{E}_{n-1}^0)$. Then $\mathcal{V}$ sits in the exact sequence*

$$\begin{aligned} 0 &\rightarrow \phi^*(\mathcal{L}^{-1} \otimes \tilde{\pi}_1^* \mathcal{O}_{\mathfrak{M}(n-1)}(\mathfrak{D}_{n-1}) \otimes \tilde{\pi}_2^* \mathcal{O}_X(2\ell)) \rightarrow \mathcal{V} \\ &\rightarrow R^1(\tilde{f}_{1,0})_* \tilde{ev}_1^*(\pi^*(\mathcal{E}_{n-1}^0)^* \otimes \mathcal{O}_{\mathfrak{B}_*}(-1)) \otimes \phi^*(\tilde{\pi}_1^* \mathcal{O}_{\mathfrak{M}(n-1)}(\mathfrak{D}_{n-1}) \otimes \tilde{\pi}_2^* \mathcal{O}_X(2\ell)) \rightarrow 0. \end{aligned}$$

Proof. Recall that $\mathfrak{B}_* = \mathbb{P}(\mathcal{E}_{n-1}^0)$. The kernel of the tautological surjection $\pi^* \mathcal{E}_{n-1}^0 \rightarrow \mathcal{O}_{\mathfrak{B}_*}(1) \rightarrow 0$ is a line bundle. By comparing the first Chern classes, we get

$$0 \rightarrow \pi^* \mathcal{L} \otimes \mathcal{O}_{\mathfrak{B}_*}(-1) \rightarrow \pi^* \mathcal{E}_{n-1}^0 \rightarrow \mathcal{O}_{\mathfrak{B}_*}(1) \rightarrow 0.$$

Tensoring with $\pi^* \mathcal{L}^{-1} \otimes \mathcal{O}_{\mathfrak{B}_*}(-1)$, we obtain the exact sequence

$$0 \rightarrow \mathcal{O}_{\mathfrak{B}_*}(-2) \rightarrow \pi^*(\mathcal{E}_{n-1}^0 \otimes \mathcal{L}^{-1}) \otimes \mathcal{O}_{\mathfrak{B}_*}(-1) \rightarrow \pi^* \mathcal{L}^{-1} \rightarrow 0.$$

Note that $\mathcal{E}_{n-1}^0 \otimes \mathcal{L}^{-1} \cong (\mathcal{E}_{n-1}^0)^*$. Applying $\tilde{ev}_1^*$ to the above exact sequence yields

$$0 \rightarrow \tilde{ev}_1^* \mathcal{O}_{\mathfrak{B}_*}(-2) \rightarrow \tilde{ev}_1^*(\pi^*(\mathcal{E}_{n-1}^0)^* \otimes \mathcal{O}_{\mathfrak{B}_*}(-1)) \rightarrow (\pi \circ \tilde{ev}_1)^* \mathcal{L}^{-1} \rightarrow 0.$$

By (5.1), $\pi \circ \tilde{ev}_1 = \phi \circ \tilde{f}_{1,0}$. Rewriting the 3rd term, we have

$$(5.4) \quad 0 \rightarrow \tilde{ev}_1^* \mathcal{O}_{\mathfrak{B}_*}(-2) \rightarrow \tilde{ev}_1^*(\pi^*(\mathcal{E}_{n-1}^0)^* \otimes \mathcal{O}_{\mathfrak{B}_*}(-1)) \rightarrow \tilde{f}_{1,0}^* \phi^* \mathcal{L}^{-1} \rightarrow 0.$$

Applying the functor $(\tilde{f}_{1,0})_*$ to (5.4), we get the exact sequence

$$\begin{aligned} 0 &\rightarrow \phi^* \mathcal{L}^{-1} \rightarrow R^1(\tilde{f}_{1,0})_*(\tilde{ev}_1)^* \mathcal{O}_{\mathfrak{B}_*}(-2) \\ &\rightarrow R^1(\tilde{f}_{1,0})_* \tilde{ev}_1^*(\pi^*(\mathcal{E}_{n-1}^0)^* \otimes \mathcal{O}_{\mathfrak{B}_*}(-1)) \rightarrow 0, \end{aligned}$$

where we have used the projection formula, $(\tilde{f}_{1,0})_* \mathcal{O}_{\mathfrak{D}_1} \cong \mathcal{O}_{\mathfrak{D}_0}$, and

$$R^1(\tilde{f}_{1,0})_* \mathcal{O}_{\mathfrak{D}_1} = 0, \quad (\tilde{f}_{1,0})_* \tilde{ev}_1^*(\pi^*(\mathcal{E}_{n-1}^0)^* \otimes \mathcal{O}_{\mathfrak{B}_*}(-1)) = 0.$$

By Lemma 5.1 and Lemma 3.3, we obtain the desired exact sequence for $\mathcal{V}$. $\square$

Remark 5.3. Fix a point $(V_1, x) \in \mathfrak{M}(n-1) \times X$. Via $\phi^{-1}(V_1, x) \cong \overline{\mathfrak{M}}_{0,0}(\mathbb{P}^1, d[\mathbb{P}^1])$, the restriction of $R^1(\tilde{f}_{1,0})_* \tilde{ev}_1^*(\pi^*(\mathcal{E}_1^0)^* \otimes \mathcal{O}_{\mathfrak{B}_*}(-1))$ to $\phi^{-1}(V_1, x)$ is isomorphic to

$$R^1(f_{1,0})_*(ev_1)^*(\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)),$$

where by abusing notation, we still use $f_{1,0}$ and ev_1 to denote the forgetful map and the evaluation map from $\overline{\mathfrak{M}}_{0,1}(\mathbb{P}^1, d[\mathbb{P}^1])$ to $\overline{\mathfrak{M}}_{0,0}(\mathbb{P}^1, d[\mathbb{P}^1])$ and $\mathbb{P}^1$, respectively.

6. THE VIRTUAL FUNDAMENTAL CLASS $[\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f})]^{\text{vir}}$

As we saw in the construction of the classes Ξ_i , Ξ_5 and Ξ_6 don't lie in $\mathfrak{B}_*$. The method to compute the virtual cycle restricted to $ev_1^{-1}(\Xi_i)$ in the previous section won't work for $i = 5, 6$. In this section, we shall employ the localization method of Kiem-Li to find a sufficiently small closed subset of $\overline{\mathfrak{M}}(n)$ containing the image of the virtual cycle under the evaluation map ev_1 . The result will be used to show the vanishing of the Gromov-Witten invariants $\langle \alpha \rangle_{0, d\mathfrak{f}}$ when α is dual to Ξ_5, Ξ_6 .

Let $X = \mathbb{P}^2$, and let $C_0 \subset X$ be a smooth cubic curve. Recall that the Zariski tangent space of $\overline{\mathfrak{M}}(n)$ at $V \in \overline{\mathfrak{M}}(n)$ is canonically $\text{Ext}^1(V, V)$. Therefore, the natural map $K_X = \mathcal{O}_X(-C_0) \rightarrow \mathcal{O}_X$ induces a meromorphic 2-form Θ on the moduli space $\overline{\mathfrak{M}}(n)$ given point-wisely by

$$(6.1) \quad \text{Ext}^1(V, V)^* \cong \text{Ext}^1(V, V \otimes K_X) \rightarrow \text{Ext}^1(V, V).$$

Note that Θ is holomorphic at $V \in \overline{\mathfrak{M}}(n)$ if (6.1) is an isomorphism.

The constructions in [K-L, L-L] show that the meromorphic 2-form Θ on the moduli space $\overline{\mathfrak{M}}(n)$ induces a meromorphic homomorphism:

$$(6.2) \quad \eta : \mathcal{E} \rightarrow \mathcal{O}$$

over $\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f})$. Here $\mathcal{E}$ is a suitable bundle on $\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f})$ such that

$$[\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f})]^{\text{vir}} \in H_{2(4n-6)}(\Lambda),$$

where $\Lambda \subset \overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f})$ is the degeneracy loci consisting of points at which either the map η is undefined or not surjective.

Next, we analyze the degeneracy loci Λ . Let $[\mu : (D; p) \rightarrow \overline{\mathfrak{M}}(n)] \in \Lambda$. Since $\mu_*[D] = d\mathfrak{f}$, $\mu(D)$ is contracted to a point by the Gieseker-Uhlenbeck morphism Ψ .

Lemma 6.1. *Let $[\mu : (D; p) \rightarrow \overline{\mathfrak{M}}(n)] \in \Lambda$, and put*

$$(\Psi \circ \mu)(D) = (V_k; \zeta) \in \mathfrak{M}(n-k) \times \text{Sym}^k(X) \subset \overline{\mathfrak{U}}(n)$$

for some $1 \leq k \leq (n-1)$. Then, either $\text{Supp}(\zeta) \cap C_0 \neq \emptyset$ or $V_k|_{C_0}$ is not stable.

Proof. Assume that $\text{Supp}(\zeta) \cap C_0 = \emptyset$ and $V_k|_{C_0}$ is stable. We will draw a contradiction by showing that η is both defined and surjective at $[\mu : (D; p) \rightarrow \overline{\mathfrak{M}}(n)]$.

Note that η is defined at $[\mu : (D; p) \rightarrow \overline{\mathfrak{M}}(n)]$ if Θ is holomorphic along $\mu(D)$. Let $V \in \mu(D)$. Then, $V^{**} = V_k$ and we have $0 \rightarrow V \rightarrow V_k \rightarrow Q \rightarrow 0$, where Q is a torsion sheaf with $\sum_{x \in X} h^0(X, Q_x)x = \zeta$. So $V|_{C_0} \cong V_k|_{C_0}$ is locally free and stable. From $0 \rightarrow K_X \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_{C_0} \rightarrow 0$, we get $0 \rightarrow V \otimes K_X \rightarrow V \rightarrow V|_{C_0} \rightarrow 0$. Applying the functor $\text{Hom}(V, \cdot)$, we obtain a long exact sequence:

$$\begin{aligned} 0 \rightarrow \text{Hom}(V, V \otimes K_X) &\rightarrow \text{Hom}(V, V) \rightarrow \text{Hom}(V, V|_{C_0}) \\ &\rightarrow \text{Ext}^1(V, V \otimes K_X) \rightarrow \text{Ext}^1(V, V). \end{aligned}$$

Since V and $V_k|_{C_0}$ are stable, we have $\text{Hom}(V, V) \cong \mathbb{C}$, $\text{Hom}(V, V \otimes K_X) = 0$ and $\text{Hom}(V, V|_{C_0}) \cong \text{Hom}(V_k|_{C_0}, V_k|_{C_0}) \cong \mathbb{C}$. The above exact sequence is simplified to

$$0 \rightarrow \text{Ext}^1(V, V \otimes K_X) \rightarrow \text{Ext}^1(V, V).$$

Since $\dim \text{Ext}^1(V, V \otimes K_X) = \dim \text{Ext}^1(V, V)$, we obtain an isomorphism

$$\text{Ext}^1(V, V \otimes K_X) \cong \text{Ext}^1(V, V).$$

Hence the meromorphic 2-form Θ is defined at $V \in \mu(D)$. This proves that Θ is holomorphic along $\mu(D)$. So η is defined at $[\mu : (D; p) \rightarrow \overline{\mathfrak{M}}(n)]$.

The above argument also shows that $\Theta|_{\mu(D)}$ is an isomorphism. Since μ is not a constant map, the image of $\mu_* : T_{D_{\text{reg}}} \rightarrow T_{\overline{\mathfrak{M}}(n)}$ does not lie in the null space of $\Theta : T_{\overline{\mathfrak{M}}(n)} \dashrightarrow (T_{\overline{\mathfrak{M}}(n)})^*$, where D_{reg} denotes the smooth part of D . By the vanishing criterion in [K-L], η is surjective at $[\mu : (D; p) \rightarrow \overline{\mathfrak{M}}(n)]$. $\square$

Lemma 6.2. *Let $C_0 \subset X = \mathbb{P}^2$ be a smooth cubic curve. Let $n \geq 1$, and let $V \in \overline{\mathfrak{M}}(n)$ be generic. Then, the restriction $V|_{C_0}$ is stable.*

Proof. It is well known that the cotangent bundle Ω_X is stable. So $\Omega_X \otimes \mathcal{O}_X(1) \in \overline{\mathfrak{M}}(1)$. Let ξ consist of $n - 1$ distinct points away from C_0 . Choose a surjection $\Omega_X \otimes \mathcal{O}_X(1) \rightarrow \mathcal{O}_\xi$, and let V_0 be the kernel. Then $V_0 \in \overline{\mathfrak{M}}(n)$ and $V_0|_{C_0} \cong (\Omega_X \otimes \mathcal{O}_X(1))|_{C_0}$. Since $\overline{\mathfrak{M}}(n)$ is irreducible and the open subset $\mathfrak{M}(n)$ is nonempty, our lemma will follow if we can prove that $(\Omega_X \otimes \mathcal{O}_X(1))|_{C_0}$ is stable.

Let $\mathcal{O}_{C_0}(D)$ be any sub-line-bundle of $(\Omega_X \otimes \mathcal{O}_X(1))|_{C_0}$. Note that the degree of $(\Omega_X \otimes \mathcal{O}_X(1))|_{C_0}$ is -3 , and there is an exact sequence

$$(6.3) \quad 0 \rightarrow (\Omega_X \otimes \mathcal{O}_X(1))|_{C_0} \rightarrow \mathcal{O}_{C_0}^{\oplus 3} \rightarrow \mathcal{O}_X(1)|_{C_0} \rightarrow 0$$

induced from the exact sequence $0 \rightarrow \Omega_X \rightarrow \mathcal{O}_X(-1)^{\oplus 3} \rightarrow \mathcal{O}_X \rightarrow 0$. Thus

$$(6.4) \quad \deg(D) \leq 0.$$

If $\deg(D) = 0$, then D must be the trivial divisor and $\mathcal{O}_{C_0}(D) = \mathcal{O}_{C_0}$. So $H^0(C_0, (\Omega_X \otimes \mathcal{O}_X(1))|_{C_0}) \neq 0$. On the other hand, (6.3) induces an exact sequence

$$0 \rightarrow H^0(C_0, (\Omega_X \otimes \mathcal{O}_X(1))|_{C_0}) \rightarrow H^0(C_0, \mathcal{O}_{C_0}^{\oplus 3}) \xrightarrow{f_0} H^0(C_0, \mathcal{O}_X(1)|_{C_0}).$$

The image of f_0 is $H^0(X, \mathcal{O}_X(1))|_{C_0} = H^0(C_0, \mathcal{O}_X(1)|_{C_0})$. So f_0 is surjective. Since $H^0(C_0, \mathcal{O}_{C_0}^{\oplus 3})$ and $H^0(C_0, \mathcal{O}_X(1)|_{C_0})$ have the same dimension, f_0 is an isomorphism and $H^0(C_0, (\Omega_X \otimes \mathcal{O}_X(1))|_{C_0}) = 0$. Hence we obtain a contradiction.

If $\deg(D) = -1$, then $\mathcal{O}_{C_0}(D) = \mathcal{O}_{C_0}(-x)$ for a unique $x \in C_0$. So the map

$$H^0(C_0, \mathcal{O}_{C_0}(x)^{\oplus 3}) \xrightarrow{f_1} H^0(C_0, \mathcal{O}_X(1)|_{C_0} \otimes \mathcal{O}_{C_0}(x))$$

induced from (6.3) is not injective. On the other hand, since $H^0(C_0, \mathcal{O}_{C_0}(x)) \cong \mathbb{C}$,

$$\begin{aligned} \text{im}(f_1) &= H^0(X, \mathcal{O}_X(1))|_{C_0} \otimes H^0(C_0, \mathcal{O}_{C_0}(x)) \\ &= H^0(C_0, \mathcal{O}_X(1)|_{C_0}) \otimes H^0(C_0, \mathcal{O}_{C_0}(x)) \\ &\subset H^0(C_0, \mathcal{O}_X(1)|_{C_0} \otimes \mathcal{O}_{C_0}(x)). \end{aligned}$$

So the linear system corresponding to $\text{im}(f_1)$ consists of all the elements $\ell|_{C_0} + x$, where ℓ denotes lines in X . In particular, the dimension of $\text{im}(f_1)$ is 3. Thus f_1 must be injective. Again, we obtain a contradiction.

By (6.4), $\deg(D) \leq -2$. Therefore, $(\Omega_X \otimes \mathcal{O}_X(1))|_{C_0}$ is stable. $\square$

Definition 6.3. For $n \geq 2$, we define $\mathfrak{T}_{C_0}(n)$ (respectively, $\mathfrak{U}_{C_0}(n)$) to be the subset of $\overline{\mathfrak{M}}(n)$ consisting of all the nonlocally free sheaves V such that $V|_{C_0}$ contains torsion (respectively, $V|_{C_0}$ is torsion-free and unstable).

Lemma 6.4. *Let $V \in \overline{\mathfrak{M}}(n) - \mathfrak{M}(n)$ and $\Psi(V) = (V_k; \zeta)$. Then,*

- (i) $V \in \mathfrak{T}_{C_0}(n)$ if and only if $\text{Supp}(\zeta) \cap C_0 \neq \emptyset$;
- (ii) $V \in \mathfrak{U}_{C_0}(n)$ if and only if $\text{Supp}(\zeta) \cap C_0 = \emptyset$ and $V_k|_{C_0}$ is unstable. $\square$

Lemma 6.5. (i) $(ev_1)(\Lambda) \subset \mathfrak{T}_{C_0}(n) \amalg \mathfrak{U}_{C_0}(n)$.

- (ii) Both $\mathfrak{T}_{C_0}(n)$ and $\mathfrak{U}_{C_0}(n)$ are closed subsets of $\overline{\mathfrak{M}}(n)$.

Proof. By Lemma 6.1, the image of a point in $(ev_1)(\Lambda)$ under Ψ is of the form

$$(V_k; \zeta) \in \mathfrak{M}(n-k) \times \text{Sym}^k(X) \subset \overline{\mathfrak{U}}(n),$$

where $1 \leq k \leq n-1$, and either $\text{Supp}(\zeta) \cap C_0 \neq \emptyset$ or $V_k|_{C_0}$ is not stable. So we see from Lemma 6.4 that $(ev_1)(\Lambda) \subset \mathfrak{T}_{C_0}(n) \coprod \mathfrak{U}_{C_0}(n)$. This proves (i).

Since being torsion-free and being stable are open conditions, both $\mathfrak{T}_{C_0}(n)$ and $\mathfrak{U}_{C_0}(n)$ are closed subsets of $\overline{\mathfrak{M}}(n)$. This proves (ii). $\square$

7. THE 1-POINT GROMOV-WITTEN INVARIANTS

Now we are ready to compute the 1-point Gromov-Witten invariants

$$(7.1) \quad \langle \alpha \rangle_{0,df} = \int_{[\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n),df)]^{\text{vir}}} ev_1^* \alpha,$$

where $\alpha \in H^{8n-12}(\overline{\mathfrak{M}}(n); \mathbb{C})$ denotes the Poincaré duals of the classes $\Xi_1, \dots, \Xi_6 \in H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$. By abusing notation, we use Ξ_i to stand for both the class in $H_4(\overline{\mathfrak{M}}(n); \mathbb{C})$ and its Poincaré dual in $H^{8n-12}(\overline{\mathfrak{M}}(n); \mathbb{C})$.

Lemma 7.1. $\langle \Xi_1 \rangle_{0,df} = -6/d^2$.

Proof. Let $\alpha = \Xi_1$. By Subsection 4.1, $\Xi_1 = \pi^{-1}(\{V_1\} \times \ell) = \mathbb{P}(V_1|_\ell)$, where the stable vector bundle $V_1 \in \mathfrak{M}(n-1)$ and the line $\ell \subset X$ are fixed. Let $\mathfrak{M} = (ev_1)^{-1}(\Xi_1)$. Then $\mathfrak{M} \subset \mathfrak{D}_1$. By Proposition 5.2 and Proposition 2.1 for $k=1$ and $\mathfrak{D} = \mathfrak{D}_1$, we obtain

$$(7.2) \quad \begin{aligned} \langle \alpha \rangle_{0,df} &= \int_{[\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n),df)]^{\text{vir}}} (ev_1)^* \alpha = \int_{(\tilde{f}_{1,0})^* c_{2d-1}(\mathcal{V})} (ev_1)^* \alpha \\ &= \int_{(\tilde{f}_{1,0})^* \phi^*(-\mathcal{L} + \tilde{\pi}_1^* \mathfrak{D}_{n-1} + 2\tilde{\pi}_2^* \ell) \cdot (\tilde{f}_{1,0})^* c_{2d-2}(\mathbb{E})} (ev_1)^* \alpha, \end{aligned}$$

where $\mathbb{E} = R^1(\tilde{f}_{1,0})_* \tilde{ev}_1^*(\pi^*(\mathcal{E}_1^0)^* \otimes \mathcal{O}_{\mathfrak{B}_*}(-1)) \otimes \phi^*(\tilde{\pi}_1^* \mathcal{O}_{\mathfrak{M}(n-1)}(\mathfrak{D}_{n-1}) \otimes \tilde{\pi}_2^* \mathcal{O}_X(2\ell))$. Note that $\Xi_1 \subset \mathfrak{B}_*$ and $\phi \circ \tilde{f}_{1,0} = \pi \circ \tilde{ev}_1$. So we obtain

$$\begin{aligned} \langle \alpha \rangle_{0,df} &= (\tilde{ev}_1)^* \pi^*(-\mathcal{L} + \tilde{\pi}_1^* \mathfrak{D}_{n-1} + 2\tilde{\pi}_2^* \ell) \cdot (\tilde{f}_{1,0})^* c_{2d-2}(\mathbb{E}) \cdot (\tilde{ev}_1)^*([\Xi_1] \cdot c_1(\mathcal{O}_{\mathfrak{B}_*}(\mathfrak{B}_*))) \\ &= (\tilde{f}_{1,0})^* c_{2d-2}(\mathbb{E}) \cdot (\tilde{ev}_1)^* \left(\pi^*(-\mathcal{L} + \tilde{\pi}_1^* \mathfrak{D}_{n-1} + 2\tilde{\pi}_2^* \ell) \cdot [\Xi_1] \cdot c_1(\mathcal{O}_{\mathfrak{B}_*}(\mathfrak{B}_*)) \right). \end{aligned}$$

Recall the definitions of $\mathcal{L}$ and $\mathfrak{D}_{n-1}$ in Proposition 5.2 and Lemma 3.3. We have

$$(7.3) \quad \mathcal{L} = \det(\tilde{\mathcal{E}}_{n-1}) = (2, -1) \in \text{Pic}(\Gamma \times X),$$

$$(7.4) \quad \mathfrak{D}_{n-1} = 2a_{n-1,1} - a_{n-1,2} - a_{n-1,0} = 2 \in \text{Pic}(\Gamma),$$

$$(7.5) \quad c_1(\mathcal{O}_{\mathfrak{B}_*}(\mathfrak{B}_*))|_{\mathbb{P}} = -2c_1(\mathcal{O}_{\mathbb{P}}(1)) + \pi^*(2, 2) \in \text{Pic}(\mathbb{P})$$

in view of the exact sequence (4.1), the degrees in (4.3) and Lemma 3.3. Thus,

$$(7.6) \quad \langle \alpha \rangle_{0,df} = (\tilde{f}_{1,0})^* c_{2d-2}(\mathbb{E}) \cdot (\tilde{ev}_1)^* \left(\pi^*(0, 3) \cdot [\Xi_1] \cdot (-2c_1(\mathcal{O}_{\mathbb{P}}(1)) + \pi^*(2, 2)) \right).$$

Since $[\Xi_1] = \pi^*(\{V_1\} \times \ell)$, it follows immediately that

$$(7.7) \quad \langle \alpha \rangle_{0,df} = -6 \cdot (\tilde{f}_{1,0})^* c_{2d-2}(\mathbb{E}) \cdot (\tilde{ev}_1)^*[\xi],$$

where $[\xi]$ denotes the cycle of a fixed point ξ in $\mathbb{P} \subset \mathfrak{B}_*$.

Let $\mathfrak{M}'_1 = (\tilde{ev}_1)^{-1}(\xi) = (ev_1)^{-1}(\xi)$. If $[\mu: (D; p) \rightarrow \overline{\mathfrak{M}}(n)] \in \mathfrak{M}'_1$, then $\mu(p) = \xi$ and $\pi(\mu(D)) = \pi(\mu(p)) = \pi(\xi) = (V_1, x) \in \mathfrak{M}(n-1) \times X$. So $\mu(D) = \mathfrak{f}$, which

denotes the unique fiber of π from (3.9) containing the point $\xi \in \mathfrak{B}_*$. Thus the restriction of the forgetful map $\tilde{f}_{1,0}$ to $\mathfrak{M}'_1$ gives a degree- d morphism from $\mathfrak{M}'_1$ to

$$\mathfrak{M}'_0 \stackrel{\text{def}}{=} \tilde{f}_{1,0}(\mathfrak{M}'_1) = \phi^{-1}(V_1, x).$$

Hence, as algebraic cycles, we have $(\tilde{f}_{1,0})_*[\mathfrak{M}'_1] = d[\mathfrak{M}'_0] = d \cdot \phi^*[(V_1, x)]$. By (7.7),

$$\begin{aligned} \langle \alpha \rangle_{0,d\mathfrak{f}} &= -6 \cdot c_{2d-2}(\mathbb{E}) \cdot (\tilde{f}_{1,0})_*[\mathfrak{M}'_1] = -6d \cdot c_{2d-2}(\mathbb{E}) \cdot \phi^*[(V_1, x)] \\ (7.8) \quad &= -6d \cdot c_{2d-2}(\mathbb{E}|_{\phi^{-1}(V_1, x)}). \end{aligned}$$

By Remark 5.3, $\mathbb{E}|_{\phi^{-1}(V_1, x)} \cong R^1(f_{1,0})_*(ev_1)^*(\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1))$, where $f_{1,0}$ and ev_1 denote the forgetful map and the evaluation map from the moduli space $\overline{\mathfrak{M}}_{0,1}(\mathbb{P}^1, d[\mathbb{P}^1])$ to $\overline{\mathfrak{M}}_{0,0}(\mathbb{P}^1, d[\mathbb{P}^1])$ and $\mathbb{P}^1$, respectively. We have

$$c_{2d-2}(R^1(f_{1,0})_*(ev_1)^*(\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1))) = \frac{1}{d^3}$$

by Theorem 9.2.3 in [C-K]. Therefore, $\langle \alpha \rangle_{0,d\mathfrak{f}} = -6/d^2$ by (7.8). $\square$

For $i = 2, 3$ or 4 , we may assume that the classes Ξ_2, Ξ_3 and Ξ_4 are represented by complex surfaces in $\mathbb{P} \subset \mathfrak{B}_*$. The same proofs of (7.6) and (7.7) show that

$$\langle \Xi_i \rangle_{0,d\mathfrak{f}} = a_i \cdot (\tilde{f}_{1,0})^* c_{2d-2}(\mathbb{E}) \cdot (\tilde{ev}_1)^*[\xi],$$

where $[\xi]$ denotes the cycle of a fixed point ξ in $\mathbb{P} \subset \mathfrak{B}_*$, and

$$(7.9) \quad a_i = \pi^*(0, 3) \cdot [\Xi_i] \cdot (-2c_1(\mathcal{O}_{\mathbb{P}}(1)) + \pi^*(2, 2))$$

is the intersection number in $\mathbb{P}$. Note from the last two paragraphs in the proof of Lemma 7.1 that $(\tilde{f}_{1,0})^* c_{2d-2}(\mathbb{E}) \cdot (\tilde{ev}_1)^*[\xi] = 1/d^2$. Therefore,

$$(7.10) \quad \langle \Xi_i \rangle_{0,d\mathfrak{f}} = \frac{a_i}{d^2}.$$

Theorem 7.2. *Let $d \geq 1$ and $n \geq 3$. The Gromov-Witten invariants $\langle \alpha \rangle_{0,d\mathfrak{f}}$ for the classes $\alpha = \text{PD}(\Xi_1), \dots, \text{PD}(\Xi_6) \in H^{8n-12}(\overline{\mathfrak{M}}(n); \mathbb{C})$ are respectively equal to*

$$-6/d^2, \quad 12/d^2, \quad 0, \quad -6/d^2, \quad 0, \quad 0.$$

Proof. First of all, $\langle \Xi_1 \rangle_{0,d\mathfrak{f}} = -6/d^2$ is Lemma 7.1.

Next, $\langle \Xi_2 \rangle_{0,d\mathfrak{f}} = 12/d^2$ follows from the computation of the number in (7.9):

$$a_2 = 3\pi^*\ell \cdot c_1(\mathcal{O}_{\mathbb{P}(V_1)}(1)) \cdot (-2c_1(\mathcal{O}_{\mathbb{P}(V_1)}(1)) + 2\pi^*\ell) = 12$$

by (4.8), where $\pi : \mathbb{P}(V_1) \rightarrow X$ denotes the tautological projection.

Since $\Xi_3 = \pi^{-1}(\Gamma \times \{y\})$ with $y \in X$, $\langle \Xi_3 \rangle_{0,d\mathfrak{f}} = 0$ comes from the computation

$$(7.11) \quad a_3 = \pi^*(0, 3) \cdot [\Xi_3] \cdot (-2c_1(\mathcal{O}_{\mathbb{P}}(1)) + \pi^*(2, 2)) = 0.$$

Similarly, $\langle \Xi_4 \rangle_{0,d\mathfrak{f}} = -6/d^2$ follows from the computation

$$a_4 = \pi^*(0, 3) \cdot c_1(\mathcal{O}_W(1)) \cdot (-2c_1(\mathcal{O}_W(1)) + \pi^*(2, 2)) = -6$$

by (4.11), where $W = \pi^{-1}(\Gamma \times \ell) \subset \mathbb{P} \subset \overline{\mathfrak{M}}(n)$ for a fixed line $\ell \subset X$.

To prove $\langle \Xi_5 \rangle_{0,d\mathfrak{f}} = 0$, choose the vector bundle $V_2 \in \mathfrak{M}(n-2)$ and the distinct points $x_1, x_2 \in X$ in Subsection 4.3 such that $V_2|_{C_0}$ is stable and $x_1, x_2 \notin C_0$. By Lemma 6.4, $\Xi_5 \cap (\mathfrak{T}_{C_0}(n) \cup \mathfrak{U}_{C_0}(n)) = \emptyset$. Hence $\Xi_5 \cap (ev_1)(\Lambda) = \emptyset$ by Lemma 6.5 (i).

Since $[\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f})]^{\text{vir}} \in H_{2(4n-6)}(\Lambda)$, we get

$$(7.12) \quad \langle \Xi_5 \rangle_{0,d\mathfrak{f}} = \int_{[\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f})]^{\text{vir}}} ev_1^* \Xi_5 = ev_{1*}[\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f})]^{\text{vir}} \cdot \Xi_5 = 0.$$

Finally, to prove $\langle \Xi_6 \rangle_{0,d\mathfrak{f}} = 0$, choose the vector bundle $V_2 \in \mathfrak{M}(n-2)$ and the point $x \in X$ in Subsection 4.4 such that $V_2|_{C_0}$ is stable and $x \notin C_0$. Now our result follows from the same proof in the previous paragraph. $\square$

Remark 7.3. Let $d \geq 1$ and $n \geq 3$. Using the theorem above, one can show that

$$ev_{1*}[\overline{\mathfrak{M}}_{0,1}(\overline{\mathfrak{M}}(n), d\mathfrak{f})]^{\text{vir}} = 1/d^2 \mathfrak{T}_{C_0}(n).$$

ACKNOWLEDGMENTS

The authors would like to thank Jun Li for helpful discussions. We also would like to thank the referee for suggestions which helped to remove the condition $n \geq 5$ in the original draft.

REFERENCES

- [Bea] A. Beauville, *Sur la cohomologie de certains espaces de modules de fibrés vectoriels*, Geometry and Analysis (Bombay, 1992), 37-40, Tata Inst. Fund. Res., Bombay, 1995. MR1351502 (96f:14011)
- [Beh] K. Behrend, *Gromov-Witten invariants in algebraic geometry*, Invent. Math. **127** (1997) 601-617. MR1431140 (98i:14015)
- [B-F] K. Behrend, B. Fantechi, *The intrinsic normal cone*, Invent. Math. **128** (1997) 45-88. MR1437495 (98e:14022)
- [C-K] D. Cox, S. Katz, *Mirror symmetry and algebraic geometry*, Mathematical Surveys and Monographs **68**, Amer. Math. Soc., Providence, RI, 1999. MR1677117 (2000d:14048)
- [E-S] G. Ellingsrud, S. Strømme, *Towards the Chow ring of the Hilbert scheme of $\mathbb{P}^2$* , J. reine angew. Math. **441** (1993), 33-44. MR1228610 (94i:14004)
- [Ful] W. Fulton, *Intersection Theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete 3. Folge **2**. Springer, Berlin Heidelberg New York Tokyo, 1994. MR732620 (85k:14004)
- [F-P] W. Fulton, R. Pandharipande, *Notes on stable maps and quantum cohomology*. Algebraic Geometry—Santa Cruz 1995, 45-96, Proc. Sympos. Pure Math. **62**, Amer. Math. Soc., Providence, RI, 1997. MR1492534 (98m:14025)
- [Get] E. Getzler, *Intersection theory on $\overline{M}_{1,4}$ and elliptic Gromov-Witten invariants*, J. Amer. Math. Soc. **10** (1997) 973-998. MR1451505 (98f:14018)
- [Gro] A. Grothendieck, *Étude cohomologique des faisceaux cohérents*, Éléments de géométrie algébrique. III, Publ. Inst. Hautes Études Sci. Math, No. **17** (1963). MR0163911 (29:1210)
- [H-H] A. Hirschowitz, K. Hulek, *Complete families of stable vector bundles over $\mathbb{P}_2$* . With an appendix by K. Hulek and S. A. Strømme. Lecture Notes in Math. **1194**, Complex Analysis and Algebraic Geometry (Gottingen, 1985), 19-40, Springer, Berlin, 1986. MR855874 (87j:14019)
- [K-L] Y. Kiem, J. Li, *Gromov-Witten invariants of varieties with holomorphic 2-forms*. Preprint.
- [LJ1] J. Li, *Algebraic geometric interpretation of Donaldson's polynomial invariants*, J. Differ. Geom. **37** (1993), 417-466. MR1205451 (93m:14007)
- [LJ2] J. Li, *Kodaira dimension of moduli space of vector bundles on surfaces*, Invent. Math. **115** (1994), 1-40. MR1248077 (94i:14016)
- [L-L] J. Li, W.-P. Li, *Two point extremal Gromov-Witten invariants of Hilbert schemes of points on surfaces*. Preprint.
- [LT1] J. Li, G. Tian, *Virtual moduli cycles and Gromov-Witten invariants of algebraic varieties*, J. Amer. Math. Soc. **11** (1998) 19-174. MR1467172 (99d:14011)
- [LT2] J. Li, G. Tian, *Virtual moduli cycles and Gromov-Witten invariants of general symplectic manifolds*, Topics in symplectic 4-manifolds (Irvine, CA, 1996), First Int. Press Lect. Ser., I, Internat. Press, Cambridge, MA, 1998, 47-83. MR1635695 (2000d:53137)
- [L-Q] W.-P. Li, Z. Qin, *On 1-point Gromov-Witten invariants of the Hilbert schemes of points on surfaces*. Proceedings of 8th Gökova Geometry-Topology Conference (2001). Turkish J. Math. **26** (2002), 53-68. MR1892799 (2003m:14006)

- [Mar] E. Markman, *Integral generators for the cohomology ring of moduli spaces of sheaves over Poisson surfaces*. Adv. Math. **208** (2007), 622-646. MR2304330 (2008b:14018)
- [Mor] J.W. Morgan, *Comparison of the Donaldson polynomial invariants with their algebro-geometric analogues*, Topology **32** (1993), 449-488. MR1231956 (94m:57066)
- [Q-Z] Z. Qin, Q. Zhang, *On the crepancy of the Gieseker-Uhlenbeck morphism*. Asian J. Math. **12** (2008), 213-224. MR2439261
- [Str] S.A. Strømme, *Ample divisors on fine moduli spaces on the projective plane*, Math. Z. **187** (1984), 405-423. MR757480 (85i:14004)
- [Uhl] K. Uhlenbeck, *Removable singularity in Yang-Mills fields*, Comm. Math. Phys. **83** (1982), 11-29. MR648355 (83e:53034)
- [Yos] K. Yoshioka, *The Betti numbers of the moduli space of stable sheaves of rank 2 on $\mathbb{P}^2$* , J. reine angew. Math. **453** (1994), 193-220. MR1285785 (95f:14022)

DEPARTMENT OF MATHEMATICS, HONG KONG UNIVERSITY OF SCIENCE AND TECHNOLOGY,
CLEAR WATER BAY, KOWLOON, HONG KONG
E-mail address: `mawpli@ust.hk`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MISSOURI 65211
E-mail address: `qinz@missouri.edu`