REGULARITY OF MORREY COMMUTATORS

DAVID R. ADAMS AND JIE XIAO

ABSTRACT. This paper is devoted to presenting a new proof of boundedness of the commutator $bI_\alpha - I_\alpha b$ (in which I_α and b are regarded as the Riesz and multiplication operators) acting on the Morrey space $L^{p,\lambda}$ under $b \in \text{BMO}$, and naturally, developing a regularity theory of commutators for Morrey-Sobolev spaces $I_\alpha(L^{p,\lambda})$ via a completely original iteration of I_α . Even in the special case of $I_\alpha(L^p)$, this is a new theory.

1. INTRODUCTION

This beginning section outlines our current work and collects the fundamental notation that will be used throughout the paper.

1.1. Overview. This article is two-fold: first to give an alternative argument (based on the new dual principle for Morrey spaces in [10]) for the following implication (cf. [19], [13], [14], [22]):

$$b \in \text{BMO} \implies [b, I_\alpha] : L^{p,\lambda} \rightarrow L^{\frac{p\lambda}{\lambda-\alpha p}, \lambda} \text{ being bounded under } \lambda > \alpha p,$$

thereby establishing the corresponding property for some constant $c > 0$:

$$b \in \text{BMO} \implies \sup_{\|f\|_{L^{p,\lambda}} \leq 1} \int_{B(0,1)} \exp(c|[b, I_\alpha]f(x)|) dx < \infty \text{ under } \lambda = \alpha p;$$

second, to develop, with much more effort on relating the traces of Morrey potentials $I_\alpha(L^{p,\lambda})$ in [11] to an entirely new way of constructing both the non-negative function h enjoying

$$|[b, I_\alpha]f| \lesssim I_\alpha h, \quad \|h\|_{L^{p,\lambda}} \lesssim \|f\|_{L^{p,\lambda}}, \quad |\nabla b| \in L^N$$

and its iterated form, a regularity (or trace, or integrability) theory of the commutators associated with Riesz potential integral operators I_α acting on Morrey spaces $L^{p,\lambda}$ covering Lebesgue spaces L^p , namely, to explore the precise values of $q > 0$ and the essential properties of both non-negative Borel measures μ on $\mathbb{R}^N$ and BMO symbols b such that

$$(1.1) \quad \sup_{\|f\|_{L^{p,\lambda}} \leq 1} \int_{B(0,1)} |[b, I_\alpha]f|^q d\mu < \infty \text{ holds when } \lambda > \alpha p$$

and for some constant $c > 0$,

$$(1.2) \quad \sup_{\|f\|_{L^{p,\lambda}} \leq 1} \int_{B(0,1)} \exp(c|[b, I_\alpha]f|^q) d\mu < \infty \text{ holds when } \lambda = \alpha p.$$

Received by the editors November 4, 2010.

2010 *Mathematics Subject Classification.* Primary 42B35, 46E35, 47G10.

Key words and phrases. Morrey-Sobolev spaces, commutators, traces, weights, Choquet integrals, fractional Laplacians, Riesz integrals, maximal operators.

The second author was supported in part by NSERC of Canada.

1.2. Conventions. Before going into the next two sections detailing the above overview, let us agree to some conventions. The N -dimensional Euclidean space $\mathbb{R}^N$, the set of positive integers $\mathbb{Z}^+$, and the Euclidean open ball $B(x, r)$ with center x and radius r are our working base. On $\mathbb{R}^N$ the Morrey functions and John-Nirenberg functions – $f \in L^{p,\lambda}$ (where $1 \leq p < \infty$ and $0 < \lambda \leq N$) and $b \in \text{BMO}$ (cf. [18]) mean respectively:

$$\|f\|_{L^{p,\lambda}} = \left[\sup_{x \in \mathbb{R}^N \text{ \& } r > 0} r^{\lambda-N} \int_{B(x,r)} |f(y)|^p dy \right]^{\frac{1}{p}} < \infty$$

and

$$\|b\|_{\text{BMO}} = \sup_{x \in \mathbb{R}^N \text{ \& } r > 0} \left| \int_{B(x,r)} b - \int_{B(x,r)} b \right| < \infty,$$

where $\int_{B(x,r)} b$ denotes the Lebesgue integral mean of b over $B(x, r)$.

Given $\alpha \in (0, N)$, denote the α -th order Riesz commutator with symbol b by

$$[b, I_\alpha]f(x) = b(x)I_\alpha f(x) - [I_\alpha(bf)](x) = \int_{\mathbb{R}^N} \frac{[b(x) - b(y)]f(y)}{|x - y|^{N-\alpha}} dy,$$

where

$$I_\alpha f(x) = \int_{\mathbb{R}^N} f(y)|x - y|^{\alpha-N} dy$$

represents the α -th order Riesz operator acting on a given function f in $\mathbb{R}^N$.

The notation $(\partial/\partial x)^s U$ represents the distributional derivative of U of order $\mathbf{s} = (s_1, \dots, s_N)$ with $s_j \in \mathbb{Z}^+$, i.e.,

$$\left(\frac{\partial}{\partial x}\right)^{\mathbf{s}} = \left(\frac{\partial}{\partial x_1}\right)^{s_1} \cdots \left(\frac{\partial}{\partial x_N}\right)^{s_N},$$

and given $m \in \mathbb{Z}^+$, $|D^m U|$ is used as the Euclidean norm of the m -th order gradient

$$D^m U = \left\{ \left(\frac{\partial}{\partial x}\right)^{\mathbf{s}} U \right\}_{|\mathbf{s}|=m}, \quad \text{where } |\mathbf{s}| = \sum_{j=1}^N s_j.$$

In the above and below, $X \lesssim Y$, $X \gtrsim Y$, and $X \approx Y$ are employed to denote that there exists a constant $c > 0$ such that $X \leq cY$, $X \geq cY$, and $c^{-1}Y \leq X \leq cY$, separately. Moreover, $\int \cdots$ is used whenever an integral domain is clear and the integral is with respect to the Lebesgue measure.

2. MORREY COMMUTATORS

By a Morrey commutator we mean $[b, I_\alpha]f$, where $f \in L^{p,\lambda}$. In this section, we revisit continuity of $[b, I_\alpha]$ with $b \in \text{BMO}$ under the standard setting $\lambda > p\alpha$ via giving a new proof, and find out some properties of the commutators under the endpoint setting $\lambda = p\alpha$. To do so, we need the following Adams' extension of the classical Sobolev imbedding to Morrey spaces (cf. [3, Theorem 3.1] or [5, Theorem 4.1 & Remark 4.1]):

Theorem 2.1 (Adams). *Let $\alpha \in (0, N)$ and $\lambda \in (0, N]$.*

(i) *If $1 < p < \frac{\lambda}{\alpha}$, then*

$$\|I_\alpha f\|_{L^{\frac{\lambda p}{\lambda - \alpha p}, \lambda}} \lesssim \|f\|_{L^{p,\lambda}}.$$

(ii) If $1 < p = \frac{\lambda}{\alpha}$, then

$$\|I_\alpha f\|_{\text{BMO}} \lesssim \|f\|_{L^{p,\lambda}}.$$

2.1. All exponents $p \in (1, \frac{\lambda}{\alpha})$. Following [10], which gives a new formulation of the predual to a Morrey space with $\lambda \in (0, N)$, we write

$$H^{p,\lambda} = \left\{ f \in L^p_{loc} : \|f\|_{H^{p,\lambda}} = \inf_w \left(\int_{\mathbb{R}^N} |f(y)|^p w(y)^{1-p} dy \right)^{\frac{1}{p}} < \infty \right\},$$

where L^p_{loc} is the class of all p -locally integrable functions on $\mathbb{R}^N$ and the infimum ranges over the class $A_1^{(N-\lambda)}$ of all non-negative weights w belonging to the weight class A_1 (whose definition and related A_p -weights may be read in e.g. [16], [25], and [26]) and satisfying

$$\int_{\mathbb{R}^N} w d\Lambda_{N-\lambda}^{(\infty)} = \int_0^\infty \Lambda_{N-\lambda}^{(\infty)}(\{x \in \mathbb{R}^N : w(x) > t\}) dt \leq 1.$$

In the above and below, for $0 < \lambda < N$ the symbol $\Lambda_{N-\lambda}^{(\infty)}$ is the Hausdorff capacity of order $N - \lambda$, as a set function on $\mathbb{R}^N$.

It is known that if $\alpha \in (0, N)$, $p \in (1, \infty)$, $0 < \alpha p < \lambda$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{\lambda}$, then

$$[b, I_\alpha] : L^{p,\lambda} \rightarrow L^{q,\lambda} \iff b \in \text{BMO}.$$

See also [19], [13], [14], [22] and the references therein.

Our contribution to this direction is to present an alternative proof to the above known result.

Theorem 2.2. *Let*

$$0 < \lambda, \alpha < N, \quad 1 < p < \frac{\lambda}{\alpha}, \quad p' = \frac{p}{p-1}, \quad \bar{p} = \frac{\lambda p}{\lambda - \alpha p}, \quad \& \quad \bar{p}' = \frac{\bar{p}}{\bar{p}-1}.$$

Then the following three statements are equivalent:

- (i) $b \in \text{BMO}$;
- (ii) $[b, I_\alpha] : H^{p,\lambda} \rightarrow H^{\bar{p},\lambda}$ *is bounded*;
- (iii) $[b, I_\alpha] : L^{\bar{p}',\lambda} \rightarrow L^{p',\lambda}$ *is bounded*.

Proof. Because (ii) $\Rightarrow$ (iii) $\Rightarrow$ (i) follows from the dual pairing

$$\int ([b, I_\alpha]f)g = - \int ([b, I_\alpha]g)f \quad \forall f \in H^{p,\lambda} \quad \& \quad g \in L^{\bar{p}',\lambda}$$

and the known implication in [14]-[13], it remains to prove (i) $\Rightarrow$ (ii). Suppose $b \in \text{BMO}$. We rely on the following basic estimate, which appeared in [14, Lemma 2], [13, Lemma 2] and [22, Lemma 1]:

$$(2.1) \quad ([b, I_\alpha]f)^\# \lesssim \|b\|_{\text{BMO}} \left([M(I_\alpha f)^r]^\frac{1}{r} + [M_{\alpha t} f^t]^\frac{1}{t} \right)$$

for

$$1 < r, t < p < \frac{N}{\alpha} \quad \& \quad f \geq 0.$$

Here and henceforth,

$$f^\#(x) = \sup_{x \in B} \int_B \left| f - \int_B f \right|$$

is the sharp function of f at x , where the supremum is taken over all Euclidean balls B containing x , and

$$M_\beta f(x) = \sup_{r>0} r^\beta \int_{B(x,r)} |f|, \quad 0 \leq \beta \leq N$$

is the fractional maximal function of f ; in particular, M_0 is the standard Hardy-Littlewood maximal operator M .

Instead of solving the original problem, we work on the predual space $H^{p,\lambda}$. To see this, without loss of generality, we may assume $0 \leq f \in H^{p,\lambda}$ and $0 \leq g \in L^{q',\lambda}$, where q' is the conjugate of $q = \bar{p}$. Due to the following duality inequality (see [25, p. 146] but also the proof of [10, Theorem 4.5]),

$$(2.2) \quad \left| \int ([b, I_\alpha]f)g \right| \lesssim \int ([b, I_\alpha]f)^\# Mg,$$

from (2.1) we need to first handle the following product:

$$P_1 := \left(\inf_{w \in A_1^{(N-\lambda)}} \int [M(I_\alpha f)^r]^{\frac{q}{r}} w^{1-q} \right)^{\frac{1}{q}} \|Mg\|_{L^{q',\lambda}}.$$

But $w \in A_1$, so $w^{1-q} \in A_q$. However, via the reverse Hölder inequality, we have $w^{1-q} \in A_{q/r}$ for some $r > 1$. So, via A_p weight theory, we use [10, Theorem 4.7 (ii)] to derive

$$\begin{aligned} P_1 &\lesssim \left(\inf_{w \in A_1^{(N-\lambda)}} \int (I_\alpha f)^q w^{1-q} \right)^{\frac{1}{q}} \|g\|_{L^{q',\lambda}} \\ &\lesssim \|f\|_{H^{p,\lambda}} \|g\|_{L^{q',\lambda}}. \end{aligned}$$

Next, we look at

$$P_2 := \left(\inf_{w \in A_1^{(N-\lambda)}} \int [M_{\alpha t}(f^t)]^{\frac{q}{t}} w^{1-q} \right)^{\frac{1}{q}} \|Mg\|_{L^{q',\lambda}}.$$

For this, we need the following.

Lemma 2.3. *If $w \in A_1$, then there exists a $\delta > 0$ such that $w^t \in A_1$ whenever $0 < t < 1 + \delta$. In particular, this implication holds for all t close to 1. Moreover, if c_1 denotes the “ A_1 constant for w ”, i.e.,*

$$\int_Q w \leq c_1 \inf_Q w \quad \forall \text{ coordinate cubes } Q \subseteq \mathbb{R}^N,$$

then, with c_t the “ A_1 constant for w^t ”, we have

$$c_t \leq \begin{cases} c_1^t, & 0 < t < 1, \\ (c_1 c_{RH})^t, & 1 < t < 1 + \delta, \end{cases}$$

where c_{RH} stands for the reverse Hölder constant.

In fact, when $0 < t < 1$, we have that for any coordinate cube Q ,

$$\int_Q w^t \leq \left(\int_Q w \right)^t \leq c_1^t \inf_Q w^t.$$

While there is a $\delta > 0$ such that the reverse Hölder inequality holds with constant $c_{RN} > 0$ (cf. [26, p. 230]), consequently, under $1 < t < 1 + \delta$,

$$\left(\int_Q w^t \right)^{\frac{1}{t}} \leq c_{RH} \int_Q w \leq c_{RH} c_1 \inf_Q w.$$

Now, we use this lemma as follows: Write

$$w^{1-q} = \Theta^{1-\frac{q}{t}}, \quad \text{where } \Theta = w^{\frac{q-1}{t-1}} \in A_1$$

since the map $w \rightarrow w^t$ is bijective on A_1 . Thus, we have

$$P_2 \lesssim \|M_{\alpha t}(f^t)\|_{H^{\frac{q}{t}, \lambda}}^{\frac{1}{t}} \|g\|_{L^{q', \lambda}} \lesssim \|f^t\|_{H^{\frac{p}{t}, \lambda}}^{\frac{1}{t}} \|g\|_{L^{q', \lambda}}$$

thanks to

$$M_{\alpha t}(f^t) \lesssim I_{\alpha t}(f^t)$$

and [10, Theorem 4.7]. But again using Lemma 2.3 we see that

$$\begin{aligned} P_3 &:= \left(\inf_{w \in A_1^{(N-\lambda)}} \int (f^t)^{\frac{p}{t}} w^{1-\frac{p}{t}} \right)^{\frac{1}{p}} \|g\|_{L^{q', \lambda}} \\ &\lesssim \left(\inf_{\Theta \in A_1^{(N-\lambda)}} \int f^p \Theta^{1-p} \right)^{\frac{1}{p}} \|g\|_{L^{q', \lambda}} \\ &\lesssim \|f\|_{H^{p, \lambda}} \|g\|_{L^{q', \lambda}}, \end{aligned}$$

thereby reaching

$$P_2 \lesssim P_3 \lesssim \|f\|_{H^{p, \lambda}} \|g\|_{L^{q', \lambda}}.$$

The previous estimates for P_1 and P_2 , along with duality, yield the desired result. $\square$

2.2. The borderline $p = \frac{\lambda}{\alpha} > 1$. The foregoing argument does not apply to the endpoint case $p = \frac{\lambda}{\alpha}$. But we can settle this via a direct application of (2.1).

Theorem 2.4. *Let $1 < p = \frac{\lambda}{\alpha}$ and $0 < \lambda, \alpha < N$. If $0 < \|b\|_{\text{BMO}} < \infty$, then there exists a constant $c > 0$ such that*

$$\sup_{\|f\|_{L^{p, \lambda}} \leq 1} \int_{B(0,1)} \exp \left[\frac{c|[b, I_\alpha]f(x)|}{\|b\|_{\text{BMO}}} \right] dx < \infty.$$

Proof. Without loss of generality, we may assume $f \geq 0$. We first prove that for each $\gamma > 0$ there is a constant $c > 0$ such that

$$(2.3) \quad \int_{B(0,1)} \exp[\gamma g(x)] dx \lesssim \int_{B(0,2)} \exp[c\gamma g^\#(x)] dx$$

holds for $g \geq 0$. In fact, following the arguments for [10, Remark 4.6], [16, Theorem 3.8] and [23, Lemmas 3.5-3.6 & Theorem 3.7], we can use the Calderón-Zygmund decomposition for a cube $2Q$ (cf. [17, p. 127]) with $Q \subseteq B(0, 1) \subseteq 2Q \subseteq B(0, 2)$ to

obtain two constants $c_1, c_2 > 0$ independent of both $0 \leq g \in L^{q_0}(B(0, 2))$ for some $q_0 \in [1, \infty)$ and $q \geq q_0$ such that

$$\int_{B(0,1)} [g(x)]^q dx \leq c_1 c_2^q \int_{B(0,2)} [g^\#(x)]^q dx.$$

This, along with

$$\exp t = \sum_{k=0}^{\infty} \frac{t^k}{k!} \quad \forall t > 0,$$

yields (2.3).

Next, (2.1) suggests to us to handle two terms. The first one is

$$\int_{B(0,2)} [M(I_\alpha f)^r]^{\frac{k}{r}} dx \leq c(k) \int_{B(0,2)} [I_\alpha f(x)]^k dx \quad \text{for integer } k > r,$$

where (cf. [20, p. 12])

$$c(k) = \frac{(k/r)2^{k/r}5^N}{(k/r) - 1}.$$

Now since $0 \leq f \in L^{p,\lambda}$ with $\lambda = \alpha p$, $I_\alpha f \in \text{BMO}$ follows from Theorem 2.1 (ii) with $\alpha = \frac{\lambda}{p}$. By the well-known John-Nirenberg theorem (cf. [18]), there is a constant $c_3 > 0$ such that

$$\sum_{k=0}^{\infty} \frac{c_3^k}{k!} \int_{B(0,2)} [I_\alpha f(x)]^k dx < \infty.$$

This, in turn, induces a constant $c_{3,*} > 0$ such that

$$(2.4) \quad \sum_{k=0}^{\infty} \frac{c_{3,*}^k}{k!} \int_{B(0,2)} [M(I_\alpha f)^r(x)]^{k/r} dx < \infty.$$

The second one is

$$\int_{B(0,2)} [M_{\alpha t} f^t(x)]^{\frac{k}{t}} dx \lesssim \int_{B(0,2)} [I_{\alpha t} f^t(x)]^{k/t} dx \quad \text{for } k = 0, 1, 2, \dots$$

This follows from $M_{\alpha t} f^t \lesssim I_{\alpha t} f^t$. Note also that

$$f^t \in L^{p/t,\lambda} \quad \& \quad (\alpha t)(p/t) = \lambda.$$

So $I_{\alpha t} f^t \in \text{BMO}$ follows by Theorem 2.1(ii). Consequently, there is a constant $c_4 > 0$ such that

$$(2.5) \quad \sum_{k=0}^{\infty} \frac{c_4^k}{k!} \int_{B(0,2)} [M_{\alpha t} f^t(x)]^{k/t} dx < \infty.$$

A combination of (2.1) and (2.3) yields that for $\gamma > 0$ and $\|f\|_{L^{p,\lambda}} \leq 1$,

$$\begin{aligned} & \int_{B(0,1)} \exp \left[\frac{\gamma |[b, I_\alpha]f(x)|}{\|b\|_{\text{BMO}}} \right] dx \\ & \lesssim \int_{B(0,2)} \exp \left[\frac{c\gamma ([b, I_\alpha]f)^\#(x)}{\|b\|_{\text{BMO}}} \right] dx \\ & \lesssim \int_{B(0,2)} \exp \left[cc'\gamma \left([M(I_\alpha f)^r(x)]^{\frac{1}{r}} + [M_{\alpha t} f^t(x)]^{\frac{1}{t}} \right) \right] dx \\ & \lesssim \left(\int_{B(0,2)} \exp \left[2cc'\gamma [M(I_\alpha f)^r(x)]^{\frac{1}{r}} \right] dx \right)^{\frac{1}{2}} \\ & \quad \times \left(\int_{B(0,2)} \exp \left[2cc'\gamma [M_{\alpha t} f^t(x)]^{\frac{1}{t}} \right] dx \right)^{\frac{1}{2}}, \end{aligned}$$

where $c' > 0$ is the constant in front of $\|b\|_{\text{BMO}}$ of (2.1).

The last estimate, together with (2.4) and (2.5), derives the desired inequality, via choosing a small $\gamma > 0$. $\square$

Remark 2.5. Let $p \geq \frac{\lambda}{\alpha} > 1$, $0 < \lambda \leq N$, $f \in L^{p,\lambda}$ be supported in $B(0,1)$ with $\|f\|_{L^{p,\lambda}} \leq 1$, and $b \in L^\infty$ with

$$\|b\|_{os} = \sup_{(x,r) \in \mathbb{R}^N \times (0,\infty)} \sup_{y \in B(x,r)} |b(x) - b(y)| > 0.$$

(i) If $p = \frac{\lambda}{\alpha}$, then

$$(2.6) \quad \int_{B(0,1)} \exp \left[c \frac{|[b, I_\alpha]f(x)|}{\|b\|_{os}} \right] dx < \infty$$

is valid for any non-negative constant c less than

$$c_{\alpha,\lambda} = \frac{\lambda}{N-\alpha} \left(\frac{\omega_{N-1}}{N} \right)^{\frac{1}{p}-1},$$

where ω_{N-1} is the surface area of $B(0,1)$.

(ii) If $p > \frac{\lambda}{\alpha}$, then

$$\sup_{x \in B(0,1)} \frac{|[b, I_\alpha]f(x)|}{\|b\|_{os}} \leq \left(\frac{Np - \lambda}{\alpha p - \lambda} \right) 2^{\alpha - \frac{\lambda}{p}}.$$

Moreover,

$$[b, I_\alpha] : L^{p,\lambda} \rightarrow C^{\alpha - \frac{\lambda}{p}}$$

under $\|b\|_{os} < \infty$ and $\lambda/p < \alpha \leq N/p$. This corresponds nicely to the Morrey type embedding below (cf. [10, Remark 3.4]),

$$I_\alpha : L^{p,\lambda} \rightarrow C^{\alpha - \frac{\lambda}{p}} \quad \text{under} \quad \frac{\lambda}{p} < \alpha \leq \frac{N}{p},$$

extending the famous Morrey's lemma (cf. [20, p. 30, 1.53 Theorem]):

$$|\nabla u| \in L^{p,\lambda} \ \& \ \lambda < p \implies u \in C^{1 - \frac{\lambda}{p}}.$$

3. REGULARITY PROPERTIES

Concerning the continuity of an operator, I_α and $[b, I_\alpha]$ (under $b \in \text{BMO}$) behave very similarly. Thus, it is reasonable to expect that the regularity estimates for the Morrey commutators are similar to those for the Morrey potentials under some further assumptions on b since BMO functions are only defined with respect to the Lebesgue N -measure. Below is a summary of the regularity estimates for the Morrey potentials established in [11] (see also [1] regarding an early result of the case $\lambda = N$).

Theorem 3.1 (Adams-Xiao). *Let μ be a non-negative Borel measure on $\mathbb{R}^N$ such that*

$$\sup_{(r,x) \in (0,\infty) \times \mathbb{R}^N} \frac{\mu(B(x,r))}{r^d} < \infty \quad \text{for } N \geq d > \lambda - \alpha p \geq 0.$$

(i) *If $1 < p < \frac{\lambda}{\alpha}$ and $0 < \lambda < N$, then*

$$\sup_{\|f\|_{L^{p,\lambda}} \leq 1} \int_{B(0,1)} |I_\alpha f|^q d\mu < \infty \quad \text{for } q < \frac{dp}{\lambda - \alpha p}$$

and

$$\sup_{\|f\|_{L^{p,\lambda}} \leq 1} \int_{B(0,1)} \frac{|I_\alpha f|^{\tilde{p}}}{[\ln(1 + |I_\alpha f|)]^\gamma} d\mu < \infty \quad \text{for } \tilde{p} = \frac{dp}{\lambda - \alpha p} \text{ \& } \gamma > 2.$$

(ii) *If $1 < p = \frac{\lambda}{\alpha}$ and $0 < \lambda \leq N$, then there exists a constant $c > 0$ such that*

$$\sup_{\|f\|_{L^{p,\lambda}} \leq 1} \int_{B(0,1)} \exp(c|I_\alpha f|^q) d\mu < \infty$$

holds for

$$\text{either } (\lambda, q) \in (0, N) \times (0, 1] \quad \text{or } (\lambda, q) = \left(N, \frac{N}{N-1}\right).$$

3.1. All exponents $p \in (1, \frac{\lambda}{\alpha})$. Based on Theorem 3.1 (i), we discover the following result.

Theorem 3.2. *Let $|\nabla b| \in L^N$, $\alpha \in (0, N)$, $p > 1$, and μ be a non-negative Borel measure on $\mathbb{R}^N$ such that*

$$\sup_{(r,x) \in (0,\infty) \times \mathbb{R}^N} \frac{\mu(B(x,r))}{r^d} < \infty \quad \text{holds for } N \geq d > \lambda - \alpha p > 0.$$

Then

$$\sup_{\|f\|_{L^{p,\lambda}} \leq 1} \int_{B(0,1)} |[b, I_\alpha]f|^q d\mu < \infty$$

holds for

$$\text{either } (\lambda, q) \in (0, N) \times \left(0, \frac{dp}{\lambda - \alpha p}\right) \quad \text{or } (\lambda, q) = \left(N, \frac{dp}{\lambda - \alpha p}\right).$$

Proof. First of all, let us control the case

$$(\lambda, q) = \left(N, \frac{dp}{\lambda - \alpha p}\right).$$

Under this, we consider two subcases as follows.

Subcase 1: $\alpha = m \in \mathbb{Z}^+$. Set

$$U = [b, I_\alpha]f \quad \& \quad K_\alpha(x, y) = |x - y|^{\alpha-N}.$$

If $|\mathbf{s}| = m$, then according to [3, p. 769], we have

$$(3.1) \quad |U(x)| \lesssim I_m \left(\sum_{|\mathbf{s}|=m} \left| \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} U(x) \right| \right)$$

holds almost everywhere with respect to the N -Lebesgue measure. Note that if $m = 1$, then

$$\left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} [b, I_\alpha](x) = \left[b, \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} K_\alpha \right] f(x) + \left(\left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} b(x) \right) (I_\alpha f)(x).$$

Thus, if $f \in L^{\bar{q}, \lambda}$ (in which case $\bar{q} > 1$ will be selected later on, but $\lambda = N$ is always assumed), then Hölder's inequality and Theorem 2.1 yield

$$\begin{aligned} \left\| \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} [b, I_\alpha]f \right\|_{L^{\bar{q}, \lambda}} &\leq \left\| \left[b, \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} K_\alpha \right] f \right\|_{L^{\bar{q}, \lambda}} + \left\| \left(\left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} b(x) \right) (I_\alpha f) \right\|_{L^{\bar{q}, \lambda}} \\ &\lesssim \|b\|_{\text{BMO}} \|f\|_{L^{\bar{q}, \lambda}} + \| |D^1 b| \|_{L^{\lambda, \lambda}} \|f\|_{L^{\bar{q}, \lambda}} \\ &\lesssim \| |\nabla b| \|_{L^N} \|f\|_{L^{\bar{q}, \lambda}}. \end{aligned}$$

This last estimate and (3.1) give

$$(3.2) \quad |[b, I_\alpha]f| \lesssim I_\alpha h,$$

where

$$h = \sum_{|\mathbf{s}|=1} \left| \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} [b, I_\alpha]f \right|$$

and

$$\|h\|_{L^{\bar{q}, \lambda}} \lesssim \| |\nabla b| \|_{L^N} \|f\|_{L^{\bar{q}, \lambda}}.$$

Next, we take $\theta > 0$ and write the I_θ potential of the commutator as

$$\begin{aligned} I_\theta(bI_1f - I_1(bf)) &= I_\theta(bI_1f) - bI_{\theta+1}f + bI_{\theta+1}f - I_{\theta+1}(bf) \\ &= [b, I_{\theta+1}]f - [b, I_\theta](I_1f), \end{aligned}$$

or

$$(3.3) \quad [b, I_{\theta+1}]f = [b, I_\theta](I_1f) + I_\theta([b, I_1]f).$$

Below, we see that identity (3.3) is the key. In fact, from the previous case $m = 1$, we have upon setting $\theta = 1$ in (3.3),

$$(3.4) \quad |[b, I_2]f| \lesssim I_1g + I_2h,$$

where

$$0 \leq g \in L^{\tilde{p}, \lambda}, \quad \tilde{p} = \frac{\lambda p}{\lambda - p}, \quad \& \quad 0 \leq h \in L^{p, \lambda};$$

i.e., we apply (3.2) first with $\bar{q} = \tilde{p}$ and then with $\bar{q} = p$. So, the trace of the commutator can be determined by (3.4). Notice that the remark (ii) of [4, Theorem 4] ensures that

$$\int_{B(0,1)} |I_2h|^q d\mu < \infty \quad \text{for} \quad q = \frac{dp}{\lambda - 2p}$$

and

$$\int_{B(0,1)} |I_1 g|^r d\mu < \infty \quad \text{for } r = \frac{d\tilde{p}}{\lambda - \tilde{p}} = \frac{dp}{\lambda - 2p}.$$

As a result, we obtain that under $|\nabla b| \in L^N$,

$$\int_{B(0,1)} |[b, I_m]f|^{\frac{dp}{N-mp}} d\mu < \infty \quad \text{for } m = 1, 2.$$

Now, assume that the assertion is valid for $\alpha = m$. When $\alpha = m + 1$, we apply (3.3) with $\theta = m$ to derive

$$(3.5) \quad |[b, I_{m+1}]f| \lesssim |[b, I_m](I_1 f)| + I_m([b, I_1]f) \lesssim \sum_{j=0}^m I_{m+1-j} f_j,$$

where

$$0 \leq f_j \in L^{\frac{\lambda p}{\lambda - j p}, \lambda} \quad \text{for } j = 0, 1, \dots, m$$

and

$$\int_{B(0,1)} |I_{m+1-j} f_j|^q d\mu < \infty \quad \text{for } q = \frac{dp}{\lambda - (m+1)p}.$$

Clearly,

$$\int_{B(0,1)} |[b, I_{m+1}]f|^q d\mu < \infty \quad \text{for } q = \frac{dp}{\lambda - (m+1)p}.$$

So, the general case follows by induction.

Subcase 2: $m - 1 < \alpha < m \in \mathbb{Z}^+$. If $m = 1$, i.e., $0 < \alpha < 1$, then according to [4, (3.2)] we have

$$(3.6) \quad |[b, I_\alpha]f| \lesssim I_\alpha \left(\sum_{|\mathbf{s}|=1} \left| \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} I_{1-\alpha}([b, I_\alpha]f) \right| \right).$$

Hence, it is enough to verify that under $|\nabla b| \in L^N$ and $|\mathbf{s}| = 1$ one has

$$(3.7) \quad f \in L^{p, \lambda} \implies h = \left| \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} I_{1-\alpha}([b, I_\alpha]f) \right| \in L^{p, \lambda}.$$

Let us form the operator

$$T_\alpha(b, f)(x) = \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} \left(I_{1-\alpha}[b, I_\alpha]f \right),$$

where f may be assumed to be smooth whenever needed. After extending $T_\alpha(b, f)$ to $T_\xi(b, f)$, where $\xi = \zeta + i\eta$ and $0 < \zeta < 1$, we see that $T_\xi(b, \cdot)$ is an analytic family of operators. It remains to check the boundedness of the extended operator on L^p in the cases $\zeta = 0, 1$.

For $\zeta = 0$, we have

$$\begin{aligned} T_{i\eta}(b, f) &= \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} \left(I_{1-i\eta}(b I_{i\eta} f - I_{i\eta}(b f)) \right) \\ &= \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} I_{1-i\eta}[b, I_{i\eta}]f. \end{aligned}$$

Note that $\left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} I_{1-i\eta}$ exists as a bounded linear operator on L^p . Thus

$$(3.8) \quad \|T_{i\eta}(b, f)\|_{L^{p, \lambda}} \lesssim \|[b, I_{i\eta}]f\|_{L^{p, \lambda}} \lesssim \|b\|_{\text{BMO}} \|f\|_{L^{p, \lambda}} \lesssim \| |\nabla b| \|_{L^N} \|f\|_{L^{p, \lambda}}.$$

For $\zeta = 1$, we get

$$\begin{aligned} T_{1+i\eta}(b, f) &= \left(\frac{\partial}{\partial x}\right)^s \left(I_{-i\eta}(b I_{1+i\eta}f - I_{1+i\eta}(bf)) \right) \\ &= I_{-i\eta} \left(\left(\frac{\partial}{\partial x}\right)^s b I_{1+i\eta}f \right) + I_{-i\eta}([b, \tilde{K}]f), \end{aligned}$$

where

$$\tilde{K}(x, \cdot) = \left(\frac{\partial}{\partial x}\right)^s |x - \cdot|^{1+i\eta-N}$$

is a Calderón-Zygmund integral kernel. As a result, we have

$$\|[b, \tilde{K}]\|_{L^{p,\lambda}} \lesssim \|b\|_{\text{BMO}} \|f\|_{L^{p,\lambda}} \lesssim \|\nabla b\|_{L^N} \|f\|_{L^{p,\lambda}}$$

and

$$\left\| \left(\frac{\partial}{\partial x}\right)^s b I_{1+i\eta}f \right\|_{L^{p,\lambda}} \lesssim \|\nabla b\|_{L^N} \|f\|_{L^{p,\lambda}}.$$

Also note that $I_{-i\eta}$ generates a Calderón-Zygmund integral kernel. So

$$(3.9) \quad \|T_{1+i\eta}(b, f)\|_{L^{p,\lambda}} \lesssim \|\nabla b\|_{L^N} \|f\|_{L^{p,\lambda}}.$$

Using (3.8) and (3.9) as well as Stein's interpolation theorem (cf. [12, p. 209, Theorem 3.3]), we obtain that under $|\nabla b| \in L^N$,

$$(3.10) \quad \|T_\xi(b, f)\|_{L^{p,\lambda}} \lesssim \|\nabla b\|_{L^N} \|f\|_{L^{p,\lambda}}, \quad 0 < \Re \xi = \zeta < 1.$$

Thus, T_ξ is a bilinear analytic family of operators. Consequently, (3.10) is true for $\xi = \alpha \in (0, 1)$. Therefore, (3.6), (3.7) and [8, p. 193, Theorem 7.2.2] yield the required result under $0 < \alpha < 1$.

If $m > 1$, then an application of (3.3) and the above treatment of $m = 1$ gives the assertion. As a matter of fact, since $1 \leq m - 1 < \alpha < m$, there is a $\theta \in (0, 1)$ such that $\alpha = m - 1 + \theta$. When $m = 2$, from (3.3) and the handling of $m = 1$ there are two non-negative functions

$$f_1 \in L^{\frac{\lambda p}{\lambda - p}, \lambda} \quad \& \quad f_0 \in L^{p, \lambda}$$

such that

$$|[b, I_\alpha]f| \lesssim I_\theta f_1 + I_{1+\theta} f_0 \approx I_\theta f_1 + I_\alpha f_0$$

with

$$\int_{B(0,1)} |I_\theta f_1|^{\frac{dp}{\lambda - \alpha p}} d\mu < \infty, \quad \int_{B(0,1)} |I_\alpha f_0|^{\frac{dp}{\lambda - \alpha p}} d\mu < \infty, \quad \& \quad \lambda = N.$$

Of course, this last argument verifies the result in the case $m = 2$. A similar argument works for other values of m through induction.

Next, let us turn to the case

$$(\lambda, q) \in (0, N) \times \left(0, \frac{dp}{\lambda - \alpha p}\right).$$

We settle this according to two subcases.

Subcase 1: $\alpha = m \in \mathbb{Z}^+$. This follows from Theorem 3.1(i) and the argument for the previously discussed *Subcase 1* with

$$q < \frac{dp}{\lambda - 2p} \quad \& \quad r < \frac{d\tilde{p}}{\lambda - \tilde{p}}$$

as well as

$$q < \frac{dp}{\lambda - (m+1)p}$$

in place of

$$q = \frac{dp}{\lambda - 2p} \quad \& \quad r = \frac{d\tilde{p}}{\lambda - \tilde{p}}$$

as well as

$$q = \frac{dp}{\lambda - (m+1)p}.$$

Subcase 2: $m-1 < \alpha < m \in \mathbb{Z}^+$. If $m = 1$, i.e., $0 < \alpha < 1$, then since it has just been proved that $|\nabla b| \in L^N$ ensures that

$$T_\alpha(b, \cdot) : L^p \rightarrow L^p \quad \text{for } 1 < p < \frac{N}{\alpha},$$

we now use the Garcia-Cuerva and Rubio de Francia extrapolation theorem (cf. [16, pp. 411-412]) to get boundedness of $T_\alpha(b, \cdot)$ on the weighted Lebesgue space $L^p(w)$ with w being an A_p weight:

$$T_\alpha(b, \cdot) : L^p(w) \rightarrow L^p(w) \quad \text{for } 1 < p < \frac{N}{\alpha}.$$

Consequently, $T_\alpha^*(b, \cdot)$ (the dual of $T_\alpha(b, \cdot)$) and $p' = \frac{p}{p-1}$ satisfy

$$T_\alpha^*(b, \cdot) : L^{p'}(w) \rightarrow L^{p'}(w).$$

Furthermore, choosing a weight $w \in A_p$ so that $w^{-p} \in A_1^{(N-\lambda)}$, we get

$$T_\alpha^*(b, \cdot) : H^{p', \lambda} \rightarrow H^{p', \lambda}.$$

Taking the dual of $T_\alpha^*(b, \cdot)$, we achieve

$$T_\alpha(b, \cdot) : L^{p, \lambda} \rightarrow L^{p, \lambda}, \quad \text{where } 1 < p < \frac{\lambda}{\alpha} < \frac{N}{\alpha}.$$

Thus, this last boundedness plus (3.6) yields a non-negative function h such that

$$|[b, I_\alpha]f| \lesssim I_\alpha h \quad \& \quad \|h\|_{L^{p, \lambda}} \lesssim \|f\|_{L^{p, \lambda}},$$

thereby implying the desired result via Theorem 3.1(i).

If $m > 1$, then there is a $\beta \in (0, 1)$ and m non-negative functions $\{f_j\}_{j=0}^{m-1}$ (by (3.3)) such that $\alpha = m - 1 + \beta$ and (see also (3.5))

$$|[b, I_\alpha]f| \lesssim \sum_{j=0}^{m-1} I_{m-1-j+\beta} f_j$$

with

$$f_j \in L^{\frac{\lambda p}{\lambda - j p}, \lambda} \quad \text{for } j = 0, 1, \dots, m-1$$

and

$$\int_{B(0,1)} |I_{m-1-j+\beta} f_j|^q d\mu < \infty \quad \forall q < \frac{dp}{\lambda - \alpha p}.$$

As a result, we get by induction (but also by the above-treated case $m = 1$, i.e., $0 < \alpha < 1$)

$$\int_{B(0,1)} |[b, I_\alpha]f|^q d\mu < \infty \quad \forall q < \frac{dp}{\lambda - \alpha p},$$

as desired. □

Remark 3.3. Two comments are in order:

(i) Under $|\nabla b| \in L^N$, $\lambda \in (0, N]$ and $\alpha \in (0, N)$, we can prove a bit more; in fact, since there is a capacitary strong-type inequality (CSI) for $C_\alpha(\cdot; L^p)$ (but not for $C_\alpha(\cdot; L^{p,\lambda})$, $0 < \lambda < N$; see also [9]), one can get the following new trace inequality, actually a CSI:

$$\int |[b, I_\alpha]f|^p dC_\alpha(\cdot; L^p) = \int_0^\infty C_\alpha(\{x : |[b, I_\alpha]f(x)| > t\}; L^p) dt^p \lesssim \|f\|_{L^p}^p.$$

(ii) It is conjectured that Theorem 3.2 is true for $b \in I_\alpha(L^{\frac{\lambda}{\alpha}, \lambda}) \subseteq \text{BMO}$ with $0 < \alpha < \lambda \leq N$ (cf. [4] & [6]).

3.2. The borderline $p = \frac{\lambda}{\alpha} > 1$. Regarding the regularity estimates for the Morrey commutators at the endpoint case $\alpha p = \lambda$, we first get the following result.

Theorem 3.4. *Let $|\nabla b| \in L^N$, $\alpha \in \mathbb{Z}^+ \cap (0, N)$, $p = \frac{\lambda}{\alpha} = \frac{N}{\alpha} > 1$, and μ be a non-negative Borel measure on $\mathbb{R}^N$ such that*

$$\sup_{(r,x) \in (0,\infty) \times \mathbb{R}^N} \frac{\mu(B(x,r))}{r^d} < \infty \quad \text{holds for } N \geq d > 0.$$

Then there is a constant $c > 0$ such that

$$\sup_{\|f\|_{L^{p,\lambda}} \leq 1} \int_{B(0,1)} \exp(c|[b, I_\alpha]f|^q) d\mu < \infty \quad \forall q \in \left(0, \frac{N}{N-1}\right).$$

Proof. First of all, we show that if

$$E_t = \{x \in \mathbb{R}^N : |[b, I_\alpha]f(x)| > t\} \cap B(0,1) \quad \forall t > 0,$$

then

$$C_1(E_t; L^N) \lesssim t^{-N} \quad \forall \text{ large } t > 0.$$

To get this, we use the special case $\lambda = N$ of (3.5):

$$|[b, I_m]f| \lesssim \sum_{j=0}^{m-1} I_{m-j} f_j, \quad \text{where } 0 \leq f_j \in L^{\frac{pN}{N-jp}}$$

and [8, Theorem 5.5.1 (d)] to obtain a constant $\kappa > 0$ such that if

$$E_{j,t} = \left\{x \in \mathbb{R}^N : I_{m-j} f_j(x) > \frac{t}{\kappa}\right\} \cap B(0,1), \quad j = 0, 1, \dots, m-1,$$

then

$$C_1(E_t; L^N) \lesssim \sum_{j=0}^{m-1} \left(C_{m-j}(E_{j,t}; L^{\frac{N}{m-j}})\right)^{\frac{(N-1)(m-j)}{N-m+j}}$$

and hence, it follows from [2] (cf. [20, 1.86 Lemma]) and CSI in [4] (cf. Remark 3.3 (i)) that

$$\begin{aligned} \int_{t_0}^\infty C_1(E_t; L^N) dt^q &\lesssim \sum_{j=0}^{m-1} \left(\int_{t_0}^\infty C_{m-j}(E_{j,t}; L^{\frac{N}{m-j}}) dt^{r_j} \right)^{\frac{(N-1)(m-j)}{N-m+j}} \\ (3.11) \quad &\lesssim \sum_{j=0}^{m-1} \|f\|_{L^{r_j}}^q \\ &< \infty \end{aligned}$$

holds for $t_0 > 1$ being big enough, where

$$r_j = \frac{\left(\frac{N}{m-j} - 1\right)q}{N-1} \quad \& \quad q < N.$$

Next, using [8, Theorem 5.1.12 & Corollary 5.1.14] it follows that for the given μ ,

$$(3.12) \quad C_1(E_t; L^N) \geq \left(\ln \frac{1}{\mu(E_t)} \right)^{1-N},$$

where $\mu(E_t)$ is of course assumed to be less than 1. As a result of (3.11) and (3.12), it follows that

$$(3.13) \quad \int_{t_0}^{2t_0} \left(\ln \frac{1}{\mu(E_t)} \right)^{1-N} dt^q \lesssim 1$$

for $t_0 > 1$ being big enough. Note that the constant on the right-hand side of (3.13) is independent of f , t_0 , and $\|f\|_{L^{p,\lambda}} \leq 1$. Thus (3.13) yields

$$\left(\ln \frac{1}{\mu(E_{t_0})} \right)^{1-N} t_0^q \lesssim 1,$$

whence

$$\mu(E_t) \lesssim \exp(-c_1 t^{\frac{q}{N-1}})$$

holds for some constant $c_1 > 0$ and $t \geq t_0$. Consequently, there is a constant $c \in (0, c_1)$ such that

$$\begin{aligned} \int_{B(0,1)} \exp(c|[b, I_\alpha]f|^r) d\mu &= \int_0^\infty \mu(E_t) d \exp(ct^r) \\ &= \left(\int_0^{t_0} + \int_{t_0}^\infty \right) \mu(E_t) d \exp(ct^r) \\ &\lesssim \mu(B(0,1)) (\exp(ct_0^r) - 1) \\ &\quad + c \int_{t_0}^\infty \exp(-c_1 t^{\frac{q}{N-1}} + ct^r) dt^r \\ &< \infty \end{aligned}$$

holds for all $r < \frac{q}{N-1} < \frac{N}{N-1}$. □

Remark 3.5. Theorem 3.4 is optimal in the sense that if f is a nice function, say $f \in C_0^\infty(\mathbb{R}^N)$, then $[b, I_\alpha]f$ can blow up on a zero set of $C_1(\cdot, L^N)$, i.e.,

$$C_1(\{x \in \mathbb{R}^N : |[b, I_\alpha]f(x)| = \infty\}; L^N) = 0.$$

The proof uses (3.2) and [8, Theorem 5.5.1 (d)], yielding that if

$$E_j = \{x \in \mathbb{R}^N : I_{m-j}f_j(x) = \infty\} \quad \& \quad E = \{x \in \mathbb{R}^N : |[b, I_\alpha]f(x)| = \infty\},$$

then $E \subseteq \bigcup_{j=0}^m E_j$ and hence

$$C_1(E; L^N) \lesssim \sum_{j=0}^{m-1} C_1(E_j; L^N) \lesssim \sum_{j=0}^{m-1} C_{m-j}(E_j; L^{\frac{N}{m-j}})^{\frac{(N-1)(m-j)}{N-m+j}} = 0.$$

When a hypothesis on b stronger than $|\nabla b| \in L^N$ is placed, the following becomes a natural thing.

Theorem 3.6. *Let $b \in I_\alpha(L^{\frac{N}{\alpha}})$, $\alpha \in \mathbb{Z}^+ \cap (0, N)$, $p > 1$, and μ be a non-negative Borel measure on $\mathbb{R}^N$ such that*

$$\sup_{(r,x) \in (0,\infty) \times \mathbb{R}^N} \frac{\mu(B(x,r))}{r^d} < \infty \quad \text{holds for } N \geq d > \lambda - \alpha p = 0.$$

Then there is a constant $c > 0$ such that

$$\sup_{\|f\|_{L^{p,\lambda}} \leq 1} \int_{B(0,1)} \exp(c|[b, I_\alpha]f|^q) d\mu < \infty$$

holds for

$$\text{either } (\lambda, q) \in (0, N) \times (0, 1] \quad \text{or} \quad (\lambda, q) = \left(N, \frac{p}{p-1}\right).$$

Proof. The assertion follows immediately from Theorem 3.1(ii) and the following.

Lemma 3.7. *Let $b \in I_\alpha(L^{\frac{N}{\alpha}})$. Then for any $f \in L^{p,\lambda}$ there is a non-negative function $h \in L^{p,\lambda}$ such that*

$$|[b, I_\alpha]f| \lesssim I_\alpha h \quad \text{with} \quad \|h\|_{L^{p,\lambda}} \lesssim \|f\|_{L^{p,\lambda}}$$

provided $\alpha = m \in \mathbb{Z}^+ \cap (0, N)$, $\lambda \in (0, N]$, and $1 < p \leq \frac{\lambda}{\alpha}$.

So, it remains to verify this lemma. Suppose

$$U = [b, I_\alpha]f, \quad K_\alpha(x, y) = |x - y|^{\alpha-N}, \quad \& \quad f \in L^{p,\lambda}.$$

By the Leibniz rule there are constants $c_{|\mathbf{t}|,m}$ depending only on $|\mathbf{t}|$ and m such that (cf. [7])

$$\begin{aligned} \left(\frac{\partial}{\partial x}\right)^{\mathbf{s}} U(x) &= \left[b, \left(\frac{\partial}{\partial x}\right)^{\mathbf{s}} K_\alpha\right] f(x) \\ &+ \sum_{\substack{|\mathbf{t}|=N-1 \\ |\mathbf{t}|=1}} c_{|\mathbf{t}|,m} \int \left(\frac{\partial}{\partial x}\right)^{\mathbf{t}} b(x) \left(\frac{\partial}{\partial x}\right)^{\mathbf{s}-\mathbf{t}} K_\alpha(x, y) f(y) dy \\ &+ \left(\left(\frac{\partial}{\partial x}\right)^{\mathbf{s}} b(x)\right) (I_\alpha f)(x) \\ &=: T_1 + T_2 + T_3. \end{aligned}$$

Note that

$$\left(\frac{\partial}{\partial x}\right)^{\mathbf{s}} K_\alpha(x, \cdot)$$

is a Calderón-Zygmund integral kernel, and $b \in I_\alpha(L^{\frac{N}{\alpha}}) \subseteq \text{BMO}$ thanks to Theorem 2.1(ii). So $f \in L^{p,\lambda}$ yields

$$\|T_1\|_{L^{p,\lambda}} = \left\| \left[b, \left(\frac{\partial}{\partial x}\right)^{\mathbf{s}} K_\alpha\right] f \right\|_{L^{p,\lambda}} \lesssim \|b\|_{\text{BMO}} \|f\|_{L^{p,\lambda}} \lesssim \| |D^m b| \|_{L^{N/m}} \|f\|_{L^{p,\lambda}} \quad (3.14)$$

by the well-known $L^{p,\lambda}$ -boundedness of the commutator $[b, K]$; see e.g. [15, Theorem 2.3] or [21, Theorem 3.3].

Next, the conclusion that $f \in L^{p,\lambda}$ and $b \in I_\alpha L^{\frac{N}{\alpha}}$ imply

$$(3.15) \quad T_3 = \left(\left(\frac{\partial}{\partial x}\right)^{\mathbf{s}} b\right) (I_\alpha f) \in L^{p,\lambda}$$

follows from the fact that $|D^m b| \in L^{N/m}$ and the following two situations:

Situation 1: If $N \geq \lambda > mp$, then an application of Hölder's inequality and Theorem 2.1(i) yields

$$\begin{aligned} \int_{B(z,r)} \left| \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} b(x) (I_\alpha f) \right|^p &\lesssim \left(\int_{B(z,r)} \left| \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} b(x) \right|^{\lambda/m} \right)^{\frac{pm}{\lambda}} \left(\int_{B(z,r)} |I_m f|^{\frac{p\lambda}{\lambda-mp}} \right)^{\frac{\lambda-pm}{\lambda}} \\ &\lesssim r^{N-\lambda} \| |D^m b| \|_{L^{N/m}}^p \|f\|_{L^{p,\lambda}}^p. \end{aligned}$$

Situation 2: If $N \geq \lambda = mp$, then the Hölder inequality and the Sobolev inequality for higher order derivatives on balls (cf. [24, Theorem 3.1]) are applied to derive that for $N > \lambda = mp$,

$$\begin{aligned} \int_{B(z,r)} \left| \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} b(x) (I_\alpha f) \right|^p &\lesssim \| |D^m b| \|_{L^{N/m}}^p \left(\int_{B(z,r)} |I_m f|^{\frac{pN}{N-pm}} \right)^{\frac{N-pm}{N}} \\ &\lesssim \| |D^m b| \|_{L^{N/m}}^p \int_{B(z,r)} |f|^p \\ &\lesssim r^{N-\lambda} \| |D^m b| \|_{L^{N/m}}^p \|f\|_{L^{p,\lambda}}^p. \end{aligned}$$

The case $N = \lambda = mp$ can be treated by using $\|I_m f\|_\infty$.

Finally, let us control the middle terms T_2 . By Hölder's inequality, Theorem 2.1(i), and the Sobolev embedding for higher order derivatives over balls once again, we get

$$\begin{aligned} \int_{B(z,r)} \left| \left(\frac{\partial}{\partial x} \right)^{\mathbf{t}} b(x) \left(\int \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}-\mathbf{t}} K_\alpha(x,y) f(y) dy \right) \right|^p &\lesssim \int_{B(z,r)} \left| \left(\frac{\partial}{\partial x} \right)^{\mathbf{t}} b(x) \right|^p |I_{|\mathbf{t}|} f|^p \\ &\lesssim \int_{B(z,r)} |D^{|\mathbf{t}|} b|^p |I_{|\mathbf{t}|} f|^p \\ &\lesssim \left(\int_{B(z,r)} |D^{|\mathbf{t}|} b|^{\frac{\lambda}{|\mathbf{t}|}} \right)^{\frac{p|\mathbf{t}|}{\lambda}} \left(\int_{B(z,r)} |I_{|\mathbf{t}|} f|^{\frac{p\lambda}{\lambda-p|\mathbf{t}|}} \right)^{\frac{\lambda-p|\mathbf{t}|}{\lambda}} \\ &\lesssim r^{N-\lambda} \| |D^m b| \|_{L^{N/m}}^p \|f\|_{L^{p,\lambda}}^p, \end{aligned}$$

whence

$$(3.16) \quad \left\| \left(\frac{\partial}{\partial x} \right)^{\mathbf{t}} b(x) \left(\int \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}-\mathbf{t}} K_\alpha(x,y) f(y) dy \right) \right\|_{L^{p,\lambda}} \lesssim \| |D^m b| \|_{L^{N/m}} \|f\|_{L^{p,\lambda}}.$$

Putting (3.14)-(3.15)-(3.16) together, we get

$$(3.17) \quad h = \sum_{|\mathbf{s}|=m} \left| \left(\frac{\partial}{\partial x} \right)^{\mathbf{s}} [b, I_m] f \right| \in L^{p,\lambda},$$

whence completing the argument for the lemma via (3.1). $\square$

Remark 3.8. It is conjectured that Theorem 3.6 is valid for non-integral $\alpha \in (0, N)$. On the other hand, it follows readily from

$$|[b, I_\alpha] f| \leq \|b\|_{os} |I_\alpha f| \quad \& \quad \|f\|_{L^{p,\lambda}} = \|f\|_{L^{p,\lambda}}$$

that Theorems 3.4 and 3.2 are trivially true for $b \in L^\infty$.

ACKNOWLEDGEMENTS

The authors are grateful for the referee's comments on the paper.

REFERENCES

- [1] D. R. Adams, Traces of potentials arising from translation invariant operators, *Ann. Scuola Norm. Sup. Pisa* (3) 25 (1971), 203–217. MR0287301 (44:4508)
- [2] D. R. Adams, A trace inequality for generalized potentials, *Studia Math.* 48 (1973), 99–105. MR0336316 (49:1091)
- [3] D. R. Adams, A note on Riesz potentials, *Duke Math. J.* 42 (1975), 765–778. MR0458158 (56:16361)
- [4] D. R. Adams, On the existence of capacitary strong type estimates, *Ark. Mat.* 14 (1976), 125–140. MR0417774 (54:5822)
- [5] D. R. Adams, *Lectures on L^p -Potential Theory*, Volume 2, Department of Mathematics, University of Umeå, 1981.
- [6] D. R. Adams, A sharp inequality of J. Moser for higher order derivatives, *Ann. of Math. (2)* 128 (1988), 385–398. MR960950 (89i:46034)
- [7] D. R. Adams and M. Frazier, Composition operators on potential spaces, *Proc. Amer. Math. Soc.* 114 (1992), 155–165. MR1076570 (92e:46061)
- [8] D. R. Adams and L. I. Hedberg, *Function Spaces and Potential Theory*, Springer-Verlag, Berlin, Heidelberg, 1996. MR1411441 (97j:46024)
- [9] D. R. Adams and J. Xiao, Nonlinear analysis on Morrey spaces and their capacities, *Indiana Univ. Math. J.* 53 (2004), 1629–1663. MR2106339 (2005h:31015)
- [10] D. R. Adams and J. Xiao, Morrey spaces in harmonic analysis, *Ark. Mat.* DOI:10.1007/s11512-010-0134-0, 30 pp.
- [11] D. R. Adams and J. Xiao, Morrey potentials and harmonic maps, *Comm. Math. Phys.* 308 (2011), 439–456. MR2851148
- [12] C. Bennett and R. Sharpley, *Interpolation of Operators*, Academic Press, Inc., 1988. MR928802 (89e:46001)
- [13] Y. Ding, A characterization of BMO via commutators for some operators, *Northeast. Math. J.* 13(4) (1997), 422–432. MR1612454 (2000d:42006)
- [14] G. Di Fazio and M. A. Ragusa, Commutators and Morrey spaces, *Bollettino U. M.I.* 7(5-A) (1991), 321–332. MR1138545 (93b:42028)
- [15] G. Di Fazio and M. A. Ragusa, Interior estimates in Morrey spaces for strong solutions to nondivergence form equations with discontinuous coefficients, *J. Funct. Anal.* 112 (1993), 241–256. MR1213138 (94e:35035)
- [16] J. Garcia-Cuerva and J. L. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, North-Holland Mathematics Studies, 116. Notas de Matemática [Mathematical Notes], 104. North-Holland Publishing Co., Amsterdam, 1985. MR807149 (87d:42023)
- [17] M. Giaquinta, *Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems*, *Annals of Math. Studies* 105, Princeton University Press, Princeton, N.J., 1983. MR717034 (86b:49003)
- [18] F. John and L. Nirenberg, On functions of bounded mean oscillation, *Comm. Pure Appl. Math.* 14 (1961), 415–426. MR0131498 (24:A1348)
- [19] Y. Komori and T. Mizuhara, Notes on commutators and Morrey spaces, *Hokkaido Math. J.* 32 (2003), 345–353. MR1996283 (2004f:42038)
- [20] J. Malý and W. P. Ziemer, *Fine Regularity of Solutions of Elliptic Partial Differential Equations*, Mathematical Surveys and Monographs 51, Amer. Math. Soc., 1997. MR1461542 (98h:35080)
- [21] M. A. Ragusa, Regularity of solutions of divergence form elliptic equations, *Proc. Amer. Math. Soc.* 128 (1999), 533–540. MR1641085 (2000c:35034)
- [22] M. A. Ragusa, Commutators of fractional integral operators on vanishing-Morrey spaces, *J. Glob. Optim.* 40 (2008), 361–368. MR2373563 (2008m:42026)
- [23] C. Sadosky, *Interpolation of Operators and Singular Integrals*. Pure and Appl. Math., Marcel Dekker, Inc. New York and Basel, 1979. MR551747 (81d:42001)
- [24] J. Shatah and M. Struwe, *Geometric Wave Equations*, Courant Institute of Math. Sci., New York Univ., 1998. MR1674843 (2000i:35135)

- [25] E. M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton Univ. Press, Princeton, New Jersey, 1993. MR1232192 (95c:42002)
- [26] A. Torchinsky, *Real-variable Methods in Harmonic Analysis*, Dover Publications, Inc., 2004. MR2059284

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KENTUCKY, LEXINGTON, KENTUCKY 40506-0027

E-mail address: dave@ms.uky.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, MEMORIAL UNIVERSITY OF NEWFOUNDLAND, ST. JOHN'S, NEWFOUNDLAND & LABRADOR, CANADA A1C 5S7

E-mail address: jxiao@mun.ca