CONE AVOIDING CLOSED SETS

LU LIU

ABSTRACT. We prove that for an arbitrary subtree T of $2^{<\omega}$ with each element extendable to a path, a given countable class $\mathcal{M}$ closed under disjoint union, and any set A , if none of the members of $\mathcal{M}$ strongly k -enumerate T for any k , then there exists an infinite set contained in either A or $\bar{A}$ such that for every $C \in \mathcal{M}$, $C \oplus G$ also does not strongly k -enumerate T . We give applications of this result, which include: (1) RT_2^2 doesn't imply WWKL_0 ; (2) [Ambos-Spies et al., 2004] DNR is strictly weaker than WWKL_0 ; (3) [Kjos-Hanssen, 2009] for any Martin-Löf random set A , either A or $\bar{A}$ contains an infinite subset that does not compute any Martin-Löf random set; etc. We also discuss further generalizations of this result.

1. INTRODUCTION

The interrelationship between topological properties and computability theoretic properties (usually computational power) of a class is widely studied in various branches of recursion theory. Here topological property is a fairly feeble term; for example such properties could involve

- all infinite homogeneous sets of a coloring of $[N]^n$;
- all infinite subsets of A or $\bar{A}$;
- all paths through a tree T . For this case, different combinatorial or topological conditions on the tree yield different topological conditions on the corresponding class. For example: T is finitely branching; homogeneous; $[T]$ is of positive measure, positive Hausdorff measure; etc.;
- all representations of a continuous function: $[0, 1] \rightarrow [0, 1]$.

This paper focuses on two kinds of classes, Part_k and $[Q]$, where

$$\begin{aligned} \text{Part}_k(A_1, A_2, \dots, A_{k-1}) \\ = \{X \in 2^\omega : (\exists i \leq k-1, X \subseteq A_i \vee X \subseteq \omega - \bigcup_{j=1}^{k-1} A_j) \wedge |X| = \infty\} \end{aligned}$$

and $[Q]$ is a closed set of 2^ω . On the positive side, Hirschfeldt et al. [10] proved that there exists $A \in \Delta_2^0$ such that $\text{DNR} \leq_u \text{Part}_2(A)$, where $\text{DNR} = \{f \in \omega^\omega : \varphi_n(n) \uparrow \vee \varphi_n(n) \neq f(n)\}$, and $\leq_u$ denotes the Muchnik reducibility (i.e. $\mathcal{C}_2 \leq_u \mathcal{C}_1$ means $\forall X \in \mathcal{C}_1 \exists Y \in \mathcal{C}_2$ such that $Y \leq_T X$), while on the negative side Dzhafarov and Jockusch [6] showed that for any $0 <_T C$ and any $A \in 2^\omega$, $\{C\} \not\leq_u \text{Part}_2(A)$. Kjos-Hanssen [13] showed that every real of positive effective Hausdorff dimension

Received by the editors November 20, 2010 and, in revised form, December 12, 2012.

2010 *Mathematics Subject Classification.* Primary 03B30; Secondary 03F35, 03C62, 68Q30, 03D32, 03D80, 28A78.

Key words and phrases. Ramsey's theorem, weak König's lemma, Martin-Löf randomness, randomness extraction.

computes an infinite subset of a Martin-Löf random set, and Greenberg and Miller [9] showed the same for every f -DNR (an f -DNR is a DNR g such that $(\forall n)g(n) \leq f(n)$), where f is a sufficiently “slow growing” computable function. Cholak et al. [4] showed that for any $A \in \Delta_2^0$ there exists an infinite low₂ set G contained in either A or $\bar{A}$.

We say we can cone avoid $\mathcal{C}_2$ within $\mathcal{C}_1$ iff $\mathcal{C}_2 \not\leq_u \mathcal{C}_1$. In this paper we generalize a result of the author [11], which proves the cone avoidance result for arbitrary $Part_k$ and $[Q] = 2\text{-DNR}$ (where $2\text{-DNR} = \{X \in 2^\omega : \forall n \varphi_n(n) \downarrow \rightarrow \varphi_n(n) \neq X(n)\}$).

The following is the main theorem in this paper. For a closed set $[Q]$ (also denoted as $\mathcal{Q}$) of 2^ω let $Q = \{\rho \in 2^{<\omega} : [\rho]^\preceq \cap [Q] \neq \emptyset\}$ (where $[\rho]^\preceq = \{X \in 2^\omega : X \supset \rho\}$).

Definition 1.1 (Beigel et al. [2]). Fix the canonical representation of finite sets, with each finite set denoted by D_n , where n is the (canonical) index of this finite set. Let k be a positive integer. Let $\{X_i\}_{i \in \omega}$ be an array of non-empty sets.

A *strong k -enumeration* of $\{X_i\}_{i \in \omega}$ is a function $h \in \omega^\omega$ such that $|D_{h(n)}| \leq k$ and $X_n \cap D_{h(n)} \neq \emptyset$. A *strong constant-bound-enumeration* of $\{X_i\}_{i \in \omega}$ is a strong k -enumeration for *some* k .

Let Q be a set of strings, say a tree. In the following text we also regard Q as an array of sets, i.e. $Q = \bigcup_{n \in \omega} Q_n$, where Q_n is the set of n -length strings of Q . A strong k -enumeration of Q means a strong k -enumeration of $\{Q_i\}_{i \in \omega}$.

Note that if Q is a tree and $W \subseteq \omega$ is infinite, then a strong k -enumeration of $\{Q_{r_n}\}_{r_n \in W}$ can be effectively translated to a strong k -enumeration of $\{Q_i\}_{i \in \omega}$.

Theorem 1.2. Suppose $\mathcal{M} = \{C^{(1)}, C^{(2)}, \dots\}$ is a countable class of sets and $\mathcal{Q} = [Q]$ is a closed set (of 2^ω) such that

- $\forall C^{(i)}, C^{(j)} \in \mathcal{M}, C^{(i)} \oplus C^{(j)} \in \mathcal{M}$ and
- Q doesn't admit a strong constant-bound-enumeration computable in $\mathcal{M}$ (i.e. computable in some member of $\mathcal{M}$).

Then for any set A , there exists an infinite set G such that

- $G \subseteq A \vee G \subseteq \bar{A}$ and
- $\forall C \in \mathcal{M}, G \oplus C$ also doesn't compute any strong constant-bound-enumeration of Q . In particular, $G \oplus C$ does not compute any member of $[Q]$, since otherwise clearly $G \oplus C$ computes a strong 1-enumeration of Q .

To gain some interest, we first introduce some applications in reverse mathematics and algorithm randomness theory, which will be derived from Theorem 1.2 in Section 5.

Reverse mathematics studies the proof theoretic strength of various second order arithmetic statements. Several statements are so important and fundamental that they serve as level lines. Many mathematical theorems are found to be equivalent to these statements and are unchanged under small perturbations of themselves. The relationships between these statements and “other” statements draw much attention. WKL_0 is one of these statements. WKL_0 states that every infinite binary tree admits an infinite path. It is well known that as a second order arithmetic statement, WKL_0 is equivalent to the statement that for any set C there exists $B \gg C$, where $B \gg C$ means B is of PA-degree relative to C . A good survey of reverse mathematics is [23] or [7], [8]. One of the second order arithmetic statements close to WKL_0 is RT_2^2 .

Definition 1.3. Let $[X]^k$ denote $\{F \subseteq X : |F| = k\}$. A k -coloring f is a function $[X]^n \rightarrow \{1, 2, \dots, k\}$. A set $H \subseteq [X]^k$ is homogeneous for f iff f is constant on $[H]^k$. A stable coloring f is a 2-coloring of $[\mathbb{N}]^2$ such that $(\forall n \in \mathbb{N})(\exists N)(\forall m > N) f(\{m, n\}) = f(\{N, n\})$. For a stable coloring f , $f_1 = \{n \in \mathbb{N} : (\exists N)(\forall m > N), f(m, n) = 1\}$, $f_2 = \mathbb{N} - f_1$.

Ramsey's theorem (Ramsey [20]). *For any n and k , every k -coloring of $[\mathbb{N}]^n$ admits an infinite homogeneous set.*

Let RT_k^n denote Ramsey's theorem for k -colorings of $[\mathbb{N}]^n$ and SRT_k^2 denote Ramsey's theorem restricted to a stable coloring of pairs. It is clear that RT_k^2 implies SRT_k^2 .

Jockusch [12] showed that for $n > 2$, RT_2^n is equivalent to ACA_0 , while Seetapun and Slaman [22] showed that RT_2^2 does not imply ACA_0 . As to WKL_0 , Jockusch [12] proved that WKL_0 does not imply RT_2^2 . Whether RT_2^2 implies WKL_0 remained open. A more detailed survey of Ramsey's theorem in view of reverse mathematics can be found in Cholak, Jockusch and Slaman [4]. Say a set S cone avoids a class $\mathcal{M}$ iff $(\forall C \in \mathcal{M})[C \not\leq_T S]$.

This problem has been a major focus in reverse mathematics in the past twenty years. The first important progress was made by Seetapun and Slaman [22], where they showed that

Theorem 1.4 (Seetapun and Slaman [22]). *For any countable class of sets $\{C_j\}$, $j \in \omega$, such that each C_i is non-computable, any computable 2-coloring of pairs admits an infinite cone avoiding (for $\{C_j\}$) homogeneous set.*

Parallel to this result, using Mathias forcing in a different manner, Dzhafarov and Jockusch [6, Lemma 3.2] proved that

Theorem 1.5 (Dzhafarov and Jockusch [6]). *For any set A and any countable class $\mathcal{M}$ such that each member of $\mathcal{M}$ is non-computable, there exists an infinite set G contained in either A or its complement such that G is cone avoiding for $\mathcal{M}$.*

The main idea is to restrict the computational complexity (computability power) of the homogeneous set as much as possible, with complexity measured by various measurements. Along this line, with simplicity measured by extent of lowness, Cholak, Jockusch and Slaman [4, Theorem 3.1] showed, by a fairly ingenious argument,

Theorem 1.6 (Cholak, Jockusch, and Slaman [4]). *For any computable coloring of the unordered pairs of natural numbers with finitely many colors, there is an infinite low₂ homogeneous set X .*

The author proved [11] that RT_2^2 does not imply WKL_0 . Meanwhile, it had also been wondered whether RT_2^2 implies WWKL_0 . Here WWKL_0 is a weaker version of WKL_0 as follows.

Definition 1.7. WWKL_0 : $\forall T \subseteq 2^{<\omega}$, if T is a tree s.t. $(\exists a > 0 \exists b \forall n) \frac{|\{\rho \in T_n\}|}{2^n} > \frac{a}{b}$, then there exists an infinite path $X \in [T]$.

Intuitively, WWKL_0 states: for every infinite tree $T \subseteq 2^{<\omega}$, if $\mu([T]) > 0$, then there is a path through T . It is also considered as a statement in the language of

second order arithmetic. For background on reverse mathematics see [23] and [4], which focus on the proof theoretic strength of Ramsey's theorem for pairs.

Our major application is that,

Corollary 1.8. *Over RCA_0 , RT_2^2 does not imply WWKL_0 .*

Here RCA_0 is the axiom which says that for any member X of a model of arithmetic, all sets computable in X should also be included in the model. This axiom corresponds to the wide intuition in mathematics that things that are derivable, constructible (in some sense) from a given object that is known to exist, should also be considered to exist.

Let $\text{DNR} = \{f \in \omega^\omega : \forall e, f(e) \neq \Phi_e(e) \vee \Phi_e(e) \uparrow\}$, $\text{DNR}^X = \{f \in \omega^\omega : \forall e, f(e) \neq \Phi_e^X(e) \vee \Phi_e^X(e) \uparrow\}$, and $g\text{-DNR} = \{f \in \omega^\omega : f \in \text{DNR} \wedge (\forall n) f(n) \leq g(n)\}$. DNR as a second order arithmetic statement says: $\forall X \exists f \in \text{DNR}^X$. Hirschfeldt et al. [10, Theorem 2.3] showed that SRT_2^2 implies DNR ; they constructed a Δ_2^0 set A such that any infinite subset of A or $\bar{A}$ computes a DNR function. It is also well known that WWKL_0 implies DNR . Combining this with Corollary 1.8 yields:

Corollary 1.9 (Ambos-Spies et al. [1]). *DNR is implied by WWKL_0 but not vice versa.*

Actually, Corollary 1.8 yields more. For definitions of basic notions from algorithmic randomness and dimension used below, see [5] or [19]. Lutz [16] first studied the effective version of Hausdorff dimension. Later Mayordomo [17] and Ryabko [21] gave a characterization using Kolmogorov complexity. Earlier than them, some results of Levin, Cai, Hartmanis and Staiger also indicated the deep relationship between Hausdorff dimension and Kolmogorov complexity.

Definition 1.10. $\dim(A) = \liminf_{n \rightarrow \infty} \frac{K(A \upharpoonright n)}{n}$.

Clearly, for any constant $0 < d \leq 1$, $T_d = \{\rho \in 2^{<\omega} : \frac{K_U(\rho)}{|\rho|} \geq d\}$ is a Π_1^0 set that does not admit a computable strong constant-bound-enumeration (the proof proceeds exactly the same as Lemma 5.3); therefore the induced Π_1^0 class satisfies the conditions given in Theorem 1.2.

The following corollary related to algorithmic randomness theory answers a question of Joe Miller.

Corollary 1.11. *There exists a DNR function that does not compute any binary sequence with positive effective Hausdorff dimension.*

Another interesting application in algorithmic randomness theory is the following.

Corollary 1.12 (Kjos-Hanssen [13]). *For every Martin-Löf random set $A \in 2^\omega$, there exists an infinite set $G \subseteq A \vee G \subseteq \bar{A}$ such that G does not compute any Martin-Löf random set.*

In Section 3 we illustrate the basic ideas and demonstrate the vital part of the construction in an informal style for Step 1 and Step s , each parallel with the other. The frame of the proof, “tree-forcing”, together with a review of Mathias forcing, is given in Subsection 3.2. Definitions that simplify statements and needed notions are given in Section 2, and the basic facts concerned are given in Section 3. In

Section 4 we first define some operations used in the construction which are basic processes in the construction, then in Subsection 4.2 give a concrete construction stated via operations introduced in Subsection 4.1, and in Subsection 4.3 prove that the construction does provide the desired set. Section 5 gives some applications of this result. Section 6 briefly discusses generalizations of this result and proposes related questions.

2. PRELIMINARIES

Ψ is sometimes not a single Turing functional but a pair of Turing functionals, $\langle \Psi_l, \Psi_r \rangle$. $\vec{\Psi}$ denotes an array of pairs of Turing functionals. $\vec{\Psi}_i$ denotes the i^{th} component. Similarly ρ is sometimes not a single string but a pair, $\vec{\rho}$ is a sequence of pairs, and $\vec{\rho}_i = \langle \rho_{il}, \rho_{ir} \rangle$. Ψ^ρ means $\langle \Psi_l^{\rho_l}, \Psi_r^{\rho_r} \rangle$ and $\vec{\Psi}^{\vec{\rho}}$ means $(\langle \Psi_{1l}^{\rho_{1l}}, \Psi_{1r}^{\rho_{1r}} \rangle, \langle \Psi_{2l}^{\rho_{2l}}, \Psi_{2r}^{\rho_{2r}} \rangle, \dots)$.

V denotes a clopen set of 2^ω . We equate V with a finite set of strings, i.e. $[V]^\preceq = \bigcup_{i=1}^n [\rho_i]^\preceq$, where $\rho_1, \rho_2, \dots, \rho_n$ are mutually incompatible. Write $\text{height}(V)$ for $\max\{|\rho| : \rho \in V\}$. For a closed set $\mathcal{Q}$ (of 2^ω) identify $\bar{\mathcal{Q}}$ with $\{\rho : [\rho] \subseteq \bar{\mathcal{Q}}\}$ so that $\rho \in \bar{\mathcal{Q}}$ makes sense.

Let $\text{set}(\rho)$ denote $\{i \in \omega : \rho(i) = 1\}$, and let ρ/σ (Z/σ) be ρ (Z) with the first $|\sigma|$ many values replaced by σ .

For a set X , view X as an infinite binary string, and let $\pi_i^n(X) = X_i$ where $X_i \in 2^\omega$ is such that $X_i(j) = X((j-1)n + i)$, i.e. $X = \bigoplus_{i=1}^n X_i$. π can be defined on finite strings in the same way.

We say X codes an ordered k -partition of W iff $\bigcup_{i=1}^k \pi_i^k(X) = W$, and say that $\pi_i^k(X)$ is the i^{th} part of this partition. A tree $T \subseteq 2^{<\omega}$ is an ordered k -partition tree of W iff $\forall X \in [T]$ X is an ordered k -partition of W . Note that in this paper, “partition” does not mean a piecewise disjoint partition.

Definition 2.1. For $i = 1, 2, \dots, u$, let $\mathcal{K}_i = \{K_{i,1}, K_{i,2}, \dots, K_{i,m_i}\}$, where each $K_{i,j}$ is a subset of $\{1, 2, \dots, n\}$, and let $\vec{\mathcal{K}} = (\mathcal{K}_1, \mathcal{K}_2, \dots, \mathcal{K}_u)$. We call $\vec{\mathcal{K}}$ a u -supporter of $\{1, 2, \dots, n\}$ iff for every ordered u -partition (not necessarily pairwise disjoint) of $\{1, 2, \dots, n\}$, namely $P^{(1)} \cup P^{(2)} \cup \dots \cup P^{(u)} = \{1, 2, \dots, n\}$, there exists some $K_{i,j} \in \mathcal{K}_i$ such that $K_{i,j} \subseteq P^{(i)}$.

A sequence of n clopen sets $V^{(1)}, V^{(2)}, \dots, V^{(n)}$ is u -disperse iff for any ordered u -partition (not necessarily pairwise disjoint) of $\{1, 2, \dots, n\}$, $P^{(1)} \cup P^{(2)} \cup \dots \cup P^{(u)} = \{1, 2, \dots, n\}$, there exists $i \leq u$ such that $\bigcap_{j \in P^{(i)}} [V^{(j)}]^\preceq = \emptyset$.

Definition 2.2. For n many ordered u -partitions $X^{(1)}, \dots, X^{(n)}$ and for $\vec{\mathcal{K}} = \{\mathcal{K}_1, \mathcal{K}_2, \dots, \mathcal{K}_u\}$, where each $\mathcal{K}_i$ is a finite class of finite subsets of $\{1, 2, \dots, n\}$ whose members are denoted by $K_{i,j}$, let

$$\begin{aligned} \text{Cross}(X^{(1)}, X^{(2)}, \dots, X^{(n)}; \vec{\mathcal{K}}) &= Y \\ &= \left(\bigoplus_{1 \leq j \leq |\mathcal{K}_1|} Y_{K_{1,j}} \right) \oplus \left(\bigoplus_{1 \leq j \leq |\mathcal{K}_2|} Y_{K_{2,j}} \right) \oplus \dots \oplus \left(\bigoplus_{1 \leq j \leq |\mathcal{K}_u|} Y_{K_{u,j}} \right), \end{aligned}$$

where $Y_{K_{i,j}} = \bigcap_{p \in K_{i,j} \in \mathcal{K}_i} \pi_i^u(X^{(p)})$; i.e. $Y_{K_{i,j}}$ is the intersection of the i^{th} parts of those u -partitions $X^{(p)}$ for which $p \in K_{i,j}$. The order of each $Y_{K_{i,j}}$ in Y “does not matter”; the point is that given Y, i, j and $\vec{\mathcal{K}}$ one could uniformly (in Y, i, j and $\vec{\mathcal{K}}$) compute each $Y_{K_{i,j}}$.

If each $\mathcal{K}_i$ consists of all of the m -element subsets of $\{1, 2, \dots, n\}$, we also abbreviate $Cross(X^{(1)}, X^{(2)}, \dots, X^{(n)}; \vec{\mathcal{K}})$ by $Cross(X^{(1)}, X^{(2)}, \dots, X^{(n)}; m)$. For n classes $\mathcal{C}^{(1)}, \mathcal{C}^{(2)}, \dots, \mathcal{C}^{(n)}$ of ordered u -partitions,

$$Cross(\mathcal{C}^{(1)}, \mathcal{C}^{(2)}, \dots, \mathcal{C}^{(n)}; \vec{\mathcal{K}}) = \{Y \in 2^\omega : \exists X^{(i)} \in \mathcal{C}^{(i)} \text{ for } 1 \leq i \leq n, \\ Y = Cross(X^{(1)}, \dots, X^{(n)}; \vec{\mathcal{K}})\}.$$

Note that the operation $Cross$ can be defined on binary strings in a natural way; therefore if $T^{(1)}, \dots, T^{(n)}$ are computable trees, then $Cross([T^{(1)}], \dots, [T^{(n)}]; \vec{\mathcal{K}})$ is a Π_1^0 class. Let $Cross(T^{(1)}, \dots, T^{(n)}; \vec{\mathcal{K}})$ denote the corresponding computable tree.

Fixing a bijection from ω to representations of finite subsets of $2^{<\omega}$, for any X , we view each function Ψ_e^X as enumeration, identifying $\Psi_e(n)$ with the finite subset of $2^{<\omega}$ it represents, so that $|\Psi_e(n)|, \Psi_e(n) \cap X, [\Psi_e(n)]^\preceq = \bigcup_{i=1}^{|\Psi_e(n)|} [\rho_i]^\preceq$, etc. make sense (if $\Psi_e(n) \uparrow$ we view $\Psi_e(n)$ as the empty set).

3. BASIC IDEAS AND SOME STEPS

We will construct a set G with $G \subseteq A \vee G \subseteq \bar{A}$, satisfying for all $e, j \in \mathbb{N}$:

- $R(e; j)$: $\Psi_e^{C^{(j)} \oplus G}$ is not a strong e -enumeration of Q . I.e. $\Psi_e^{C^{(j)} \oplus G}$ is not total, or $(\exists n)(|\Psi_e^{C^{(j)} \oplus G}(n)| > e \vee \Psi_e^{C^{(j)} \oplus G}(n) \cap Q = \emptyset)$, or $(\exists l)(\forall n)(\exists \rho \in \Psi_e^{C^{(j)} \oplus G}(n))|\rho| < l$.
- P_e : $|G| \geq e$.

3.1. Step 1: Satisfying $R(1; j)$.

Case i. If we can find some $\rho \in 2^{<\omega}$ such that $set(\rho) \subset A \vee set(\rho) \subset \bar{A}$ and some n such that $\Psi_1^{C^{(j)} \oplus \rho}(n) \downarrow \cap Q = \emptyset \vee |\Psi_1^{C^{(j)} \oplus \rho}(n)| > 1$, then we can satisfy $R(1; j)$ by finitely extending our initial segment requirement to ρ . In this case we say Case i occurs to $R(1; j)$.

Case ii. If there is no such ρ , i.e. Case i doesn't occur to $R(1; j)$. Then we make $\Psi_1^{C^{(j)} \oplus G}$ non-total; we can achieve this by *trying* to find three disjoint clopen sets $[V^{(1)}]^\preceq, [V^{(2)}]^\preceq, [V^{(3)}]^\preceq$ such that the following classes are non-empty:

$$[T_{V^{(i)}}] = \{X = X_l \oplus X_r : X_l \cup X_r = \omega \wedge [(\forall Z \text{ s.t. } Z \subseteq X_l \vee Z \subseteq X_r)(\forall n) \\ \Psi_1^{C^{(j)} \oplus Z}(n) \downarrow \rightarrow (|\Psi_1^{C^{(j)} \oplus Z}(n)| \leq 1 \wedge [\Psi_1^{C^{(j)} \oplus Z}(n)]^\preceq \cap [V^{(i)}]^\preceq \neq \emptyset)]\}.$$

Let

$$[T_1] = Cross([T_{V^{(1)}}], [T_{V^{(2)}}], [T_{V^{(3)}}]; 2),$$

i.e. $\forall X \in [T_1]$, $X = Y_1 \oplus Y_2 \oplus Y_3 \oplus Y_4 \oplus Y_5 \oplus Y_6$, where

$$\begin{aligned} Y_1 &= \pi_l^2(X') \cap \pi_l^2(X'') && \text{for some } X' \in [T_{V^{(1)}}], X'' \in [T_{V^{(2)}}], \\ Y_2 &= \pi_l^2(X') \cap \pi_l^2(X'') && \text{for some } X' \in [T_{V^{(2)}}], X'' \in [T_{V^{(3)}}], \\ Y_3 &= \pi_l^2(X') \cap \pi_l^2(X'') && \text{for some } X' \in [T_{V^{(3)}}], X'' \in [T_{V^{(1)}}], \\ Y_4 &= \pi_r^2(X') \cap \pi_r^2(X'') && \text{for some } X' \in [T_{V^{(1)}}], X'' \in [T_{V^{(2)}}], \\ Y_5 &= \pi_r^2(X') \cap \pi_r^2(X'') && \text{for some } X' \in [T_{V^{(2)}}], X'' \in [T_{V^{(3)}}], \\ Y_6 &= \pi_r^2(X') \cap \pi_r^2(X'') && \text{for some } X' \in [T_{V^{(3)}}], X'' \in [T_{V^{(1)}}]. \end{aligned}$$

Note:

- (1) $[T_{V^{(i)}}]$ is a $\Pi_1^{0,C^{(j)}}$ class. For any general Ψ (instead of Ψ_1) and V (instead of $V^{(i)}$) the index of $[T_V]$, which is induced by Ψ, V , can be uniformly computed from indices for Ψ and V .
- (2) If $[V]^\preceq \supseteq [Q]$, then $T_V \neq \emptyset$, as otherwise Case i occurs: For $X_1 = A$ and $X_2 = \bar{A}$, if we assume that $X_1 \oplus X_2 \notin T_V$, then there are a ρ such that $\text{set}(\rho) \subseteq A \vee \text{set}(\rho) \subseteq \bar{A}$ and an $n \in \mathbb{N}$ with $[\Psi_1^{C^{(j)} \oplus \rho}(n) \downarrow]^\preceq \cap [V]^\preceq = \emptyset \vee |\Psi_1^{C^{(j)} \oplus \rho}(n)| > 1$, which implies that $[\Psi_1^{C^{(j)} \oplus \rho}(n)]^\preceq \cap Q = \emptyset \vee |\Psi_1^{C^{(j)} \oplus \rho}(n)| > 1$.
- (3) For any $i \leq 6$, $G \subseteq Y_i$, $\Psi_1^{C^{(j)} \oplus G}$ is not a strong 1-enumeration of Q . To see this, suppose on the contrary that for some $G \subseteq Y_1$, $\Psi_1^{C^{(j)} \oplus G}$ is a strong 1-enumeration of Q . Then there exists n such that $\forall \rho \in \Psi_1^{C^{(j)} \oplus G}(n)$, $|\rho| = n > \max\{\text{height}(V^{(1)}), \text{height}(V^{(2)})\}$. Then $\forall \rho \in \Psi_1^{C^{(j)} \oplus G}(n)$ either $[\rho]^\preceq \cap [V^{(1)}]^\preceq = \emptyset$ or $[\rho]^\preceq \cap [V^{(2)}]^\preceq = \emptyset$ (since $[V^{(1)}], [V^{(2)}]$ are disjoint), say $[\rho]^\preceq \cap [V^{(1)}]^\preceq = \emptyset$, which implies $[\Psi_1^{C^{(j)} \oplus G}(n)] \cap [V^{(1)}]^\preceq = \emptyset$ (note that at this step there is at most one $\rho \in \Psi_1^{C^{(j)} \oplus G}(n)$). But $G \subseteq Y_1 \subseteq \pi_i^2(X')$ for some $X' \in [T_{V^{(1)}}]$, and by the definition of $[T_{V^{(1)}}]$, $\forall n, [\Psi_1^{C^{(j)} \oplus G}(n)]^\preceq \cap [V^{(1)}]^\preceq \neq \emptyset$ if $\Psi_1^{C^{(j)} \oplus G}(n) \downarrow$.
- (4) $[T_1]$ is a $\Pi_1^{0,C^{(j)}}$ class (note that *Cross* can be applied to binary strings) whose index can be $C^{(j)}$ -computed from indices for Ψ , $V^{(1)}$, $V^{(2)}$, and $V^{(3)}$.
- (5) $\bigcup_{i=1}^6 Y_i = \omega$ (see Fact 3.6; this is just the pigeonhole principle). This is why we try to choose *three* mutually disjoint clopen sets at *this* step; i.e. if we choose, say, two disjoint clopen sets, then the union of all parts of some path through T_1 might be finite, which leads to difficulties at the next step of the construction. The advantage of requiring T to be an ordered-partition-tree will be clear in Lemma 4.5. In general, such three mutually disjoint clopen sets can't be found, so instead we *try* to find a 2-disperse sequence of clopen sets for some 2-supporter at *this* step, while at later steps 2-disperse sequences will not be enough and will have to be replaced by k' -disperse sequences.

How the requirement $R(1; j)$ is satisfied: If Case i occurs, we require that if finally $G \subseteq A(\bar{A}) \wedge \text{set}(\rho) \subset A(\bar{A})$, then $G \supset \rho$. If Case ii occurs, we require that for some path $X \in [T_1]$, G will be contained in some Y_i such that $Y_i = \pi_i^6(X)$. By the above notes clearly $R(1; j)$ is satisfied. Furthermore, Lemma 4.6 will prove that if no such 2-disperse sequence exists such that all induced classes are non-empty (i.e. Case ii does not occur), then Case i occurs; otherwise one can compute a strong k -enumeration of Q .

3.2. Tree forcing. A Mathias forcing condition is a pair (ρ, L) where $\rho \in 2^{<\omega}$ and $L \in 2^\omega$. We write $(\rho, L) \geq (\rho', L')$ and say that (ρ', L') extends (ρ, L) iff $\rho' \supset \rho \wedge L' \cup \text{set}(\rho') \subseteq L \cup \text{set}(\rho)$. A set G satisfies (ρ, L) iff $G \supset \rho \wedge G \subseteq \text{set}(\rho) \cup L$.

In this paper, a tree forcing condition is a triple $(\vec{\rho}, T, k)$, where $T \subseteq 2^{<\omega}$ is a tree and $\vec{\rho} = (\langle \rho_{1l}, \rho_{1r} \rangle, \langle \rho_{2l}, \rho_{2r} \rangle, \dots, \langle \rho_{kl}, \rho_{kr} \rangle)$ is an array of k pairs of binary strings,

such that:

- (1) each $X \in [T]$ together with $\vec{\rho}$ codes in a uniform way $2k$ Mathias forcing conditions, $\langle \rho_{1l}, \pi_1^k(X) \rangle, \langle \rho_{1r}, \pi_1^k(X) \rangle, \dots, \langle \rho_{kl}, \pi_k^k(X) \rangle, \langle \rho_{kr}, \pi_k^k(X) \rangle$, i.e. there is *one* initial segment requirement pair for *each* part.
- (2) $\bigcup_{i=1}^k \pi_i^k(X) =^* \omega$ (i.e. $\bigcup_{i=1}^k \pi_i^k(X)$ is ω minus some finite set).
- (3) $(\forall i \leq k, X \in [T]) \pi_i^k(X) \cap \{1, 2, \dots, \max\{|\rho_{il}|, |\rho_{ir}|\}\} = \emptyset$.
- (4) $\text{set}(\rho_{il}) \subseteq A \wedge \text{set}(\rho_{ir}) \subseteq \bar{A}$, where $A \in 2^\omega$ is the given set in Theorem 1.2.

A set G satisfies $(\vec{\rho}, T, k)$ iff there exist $X \in [T]$ and $i \leq k$ such that G satisfies either $\langle \rho_{il}, \pi_i^k(X) \rangle$ or $\langle \rho_{ir}, \pi_i^k(X) \rangle$. $(\vec{\rho}', T', k')$ extends $(\vec{\rho}, T, k)$, written as $(\vec{\rho}', T', k') \leq (\vec{\rho}, T, k)$, iff there exists a function $p : \{1, 2, \dots, k'\} \rightarrow \{1, 2, \dots, k\}$ such that

- (1) $\forall i \leq k', \rho'_{il} \supset \rho_{p(i),l} \wedge \rho'_{ir} \supset \rho_{p(i),r}$,
- (2) $\forall i \leq k', \forall X' \in [T'] \exists X \in [T], \text{set}(\rho'_{il}) \cup \pi_i^{k'}(X') \subseteq \text{set}(\rho_{p(i),l}) \cup \pi_{p(i)}^k(X)$ and similarly for the right-hand side. We call part i of T' a child part of part $p(i)$ of T .

We will construct a sequence $(\vec{\rho}^1, T_1, k_1) \geq (\vec{\rho}^2, T_2, k_2) \geq \dots \geq (\vec{\rho}^s, T_s, k_s) \geq \dots$ of tree forcing conditions. Each part $i \leq k_s$ of $(\vec{\rho}^s, T_s, k_s)$ will correspond to a progress information sequence $(j; e_{i,r}^{s,j}, e_{i,l}^{s,j})$ for $j \in \mathbb{N}$, which means that for all $i \leq k_s$ and $X \in [T_s]$, if G satisfies $\langle \rho_{il}^s, \pi_i^{k_s}(X) \rangle$, then G satisfies $R(e; j)$ for all $j \in \mathbb{N}$ and $e < e_{i,l}^{s,j}$, and if G satisfies $\langle \rho_{ir}^s, \pi_i^{k_s}(X) \rangle$, then G satisfies $R(e; j)$ for all $j \in \mathbb{N}$ and $e < e_{i,r}^{s,j}$. Furthermore, we will have $[T_s] \neq \emptyset$, and each T_s will be computable in some $C \in \mathcal{M}$ where $\mathcal{M}$ is the given countable class in Theorem 1.2.

Let $C^{(j)}$ for $j \in \mathbb{N}$ be a sequence of sets of $\mathcal{M}$ that is cofinal in $\mathcal{M}$, i.e. $(\forall j \in \mathbb{N})[C^{(j)} \in \mathcal{M} \wedge (\forall C \in \mathcal{M} \exists C^{(j)}, C \leq_T C^{(j)})]$, and such that $\forall j \in \mathbb{N}, C^{(j)} \leq_T C^{(j+1)}$. Fix a sequence C^{t_s} for $s \in \mathbb{N}$ of elements of $\{C^{(j)}\}_{j \in \mathbb{N}}$ such that each $C^{(j)}$ appears infinitely often in this sequence, i.e. $\exists^\infty s, t_s = j$. Step s will be devoted to $R(\dots; t_s)$, which means we make sure that if part i' of T_s has a child in T_{s+1} , then on each child of part i' , for example part i of T_{s+1} , for $j = t_s$ either $e_{ir}^{s,j} > e_{i'r}^{s-1,j}$ or $e_{il}^{s,j} > e_{i'l}^{s-1,j}$.

To avoid too many indices we use formal parameters $T, \vec{\rho}, \vec{\Psi}$. Their values are updated at each step, so $\vec{\Psi} = \vec{e}^{s-1,j}$ means we assign $e_{il}^{s-1,j}$ to be the index of Ψ_{il} , and writing $T = \{\rho \in 2^{<\omega} : \dots \sigma \in T \dots\}$ makes sense. Within each step, and from one step to the next, progress information may also need to be updated, so writing $e_{il}^{s-1,j} = e_{il}^{s,j} + 1$ or $e_{il}^{s,j} = e_{il}^{s-1,j} + 1$ also makes sense.

3.3. Some definitions and facts. Before we get a glimpse at Step s , we make the following definition in order to simplify our statements:

Definition 3.1.

- (1) $\Psi^{C \oplus \rho}$ *abandons* V on a set Y iff there exist $Z \subseteq Y$ and $n \in \mathbb{N}$ such that either $\Psi^{C \oplus Z/\rho}(n) \downarrow = D$ for a finite set D with $[D]^\perp \cap [V]^\perp = \emptyset$, or $|\Psi^{C \oplus Z/\rho}(n)|$ is greater than the index of Ψ .
- (2) $\langle \Psi_l^{C \oplus \rho_l}, \Psi_r^{C \oplus \rho_r} \rangle$ (also denoted by $\Psi^{C \oplus \rho}$) *abandons* V on $X_1 \oplus X_2$ iff *either* $\Psi_l^{C \oplus \rho_l}$ *abandons* V on X_1 *or* $\Psi_r^{C \oplus \rho_r}$ *abandons* V on X_2 .

- (3) $\Psi^{C \oplus \rho}$ (which means $\langle \Psi_l^{C \oplus \rho_l}, \Psi_r^{C \oplus \rho_r} \rangle$) *abandons* V on X iff for *all* ordered 2-partitions $X_1 \cup X_2 = X$, $\langle \Psi_l^{C \oplus \rho_l}, \Psi_r^{C \oplus \rho_r} \rangle$ *abandons* V on $X_1 \oplus X_2$. I.e. $\langle \Psi_l^{C \oplus \rho_l}, \Psi_r^{C \oplus \rho_r} \rangle$ doesn't abandon V on X iff there exists an ordered partition $X_1 \cup X_2 = X$ such that $\forall Y \supset \rho_l, Y \subseteq X_1 \cup \text{set}(\rho_l) \Rightarrow (\forall n) [(\Psi_l^{C \oplus Y}(n) \uparrow) \vee ([\Psi_l^{C \oplus Y}(n)]^\perp \cap [V]^\perp \neq \emptyset \wedge |\Psi_l^{C \oplus Y}(n)| \leq l)]$ and $\forall Y \supset \rho_r, Y \subseteq X_2 \cup \text{set}(\rho_r) \Rightarrow (\forall n) [(\Psi_r^{C \oplus Y}(n) \uparrow) \vee ([\Psi_r^{C \oplus Y}(n)]^\perp \cap [V]^\perp \neq \emptyset \wedge |\Psi_r^{C \oplus Y}(n)| \leq r)]$.

(In the above definitions, “abandons on a set” can be replaced by “abandons on a finite string”; the definitions generalize naturally.)

- (4) $\tilde{\Psi}^{C \oplus \vec{\rho}}$ *abandons* V on $X = \bigoplus_{i=1}^k X_i$ iff $\exists i \leq k$ such that the pair $\Psi_i^{C \oplus \rho_i}$ *abandons* V on X_i . I.e. $\tilde{\Psi}^{C \oplus \vec{\rho}}$ doesn't abandon V on $X = \bigoplus_{i=1}^k X_i$ iff for all i , there exist X_{il}, X_{ir} with $X_{il} \cup X_{ir} = X_i$ such that $\Psi_{il}^{C \oplus \rho_{il}}$ doesn't abandon V on X_{il} and $\Psi_{ir}^{C \oplus \rho_{ir}}$ doesn't abandon V on X_{ir} .

We also say that $\tilde{\Psi}^{C \oplus \vec{\rho}}$ doesn't abandon V on $\bigoplus_{i=1}^k (X_{il} \oplus X_{ir})$ to indicate that for all i , $\Psi_{il}^{C \oplus \rho_{il}}$ doesn't abandon V on X_{il} and $\Psi_{ir}^{C \oplus \rho_{ir}}$ doesn't abandon V on X_{ir} . (Note that this is a little abuse of notation, but the lower index il, ir shall avoid confusion.) Let $\pi_{il}^k(X) = X_{il}$ and $\pi_{ir}^k(X) = X_{ir}$.

The following two simple facts illustrate the central idea of the construction.

Fact 3.2. Let $V^{(1)}, V^{(2)}, \dots, V^{(m)}$ be an e -disperse sequence of clopen sets and let Ψ be a Turing functional such that for each i , Ψ doesn't abandon $V^{(i)}$ on X . Then, for any $Y \subseteq X$, Ψ^Y is not total or is not a strong e -enumeration. (The proof also holds if Ψ is relativized, i.e. it also holds if for some oracle C and some ρ we replace Ψ by $\Psi^{C \oplus \rho}$ and Ψ^Y by $\Psi^{C \oplus Y/\rho}$.)

Proof. Here and below Ψ abandoning V means $\Psi^{\emptyset \oplus \varepsilon}$ abandoning V , where ε is the empty string. Suppose not. Then there is some n and some $Y \subseteq X$ such that $|\Psi^Y(n)| \leq e$ and

$$\forall \rho \in \Psi^Y(n), |\rho| > \max\{\text{height}(V^{(1)}), \text{height}(V^{(2)}), \dots, \text{height}(V^{(m)})\},$$

and furthermore for each $i \leq m$, $[V^{(i)}]^\perp \cap [\Psi^Y(n)]^\perp \neq \emptyset$. For $1 \leq i \leq e$, let $P^{(i)} = \{V^{(j)} : [V^{(j)}]^\perp \cap [\rho_i]^\perp \neq \emptyset, \text{ where } \rho_i \text{ is the } i^{\text{th}} \text{ string in } \Psi^Y(n)\}$ (let $P^{(i)} = P^{(1)}$ if $i > |\Psi^Y(n)|$). Note that since $|\rho_i| > \text{height}(V^{(r)})$, we have that $[\rho_i]^\perp \cap [V^{(r)}]^\perp \neq \emptyset$ implies $[\rho_i]^\perp \subseteq [V^{(r)}]^\perp$. The $P^{(i)}$ form an e -partition of $V^{(1)}, V^{(2)}, \dots, V^{(m)}$, but $\forall i, \bigcap_{r \in P^{(i)}} [V^{(r)}]^\perp \supseteq [\rho_i]^\perp \neq \emptyset$, which contradicts the assumption that $V^{(1)}, V^{(2)}, \dots, V^{(m)}$ is an e -disperse sequence. $\square$

Fact 3.3. If a single Turing functional $\Psi_e^{C \oplus \rho}$ doesn't abandon V on X , then for any $Y \subseteq X$, $\Psi_e^{C \oplus \rho}$ doesn't abandon V on Y .

Proof. By the definition of “abandon”, item (1). $\square$

Combining Fact 3.2 and Fact 3.3:

Fact 3.4. Let $V^{(1)}, V^{(2)}, \dots, V^{(m)}$ be an e -disperse sequence of clopen sets and let Ψ be a Turing functional such that for each i , Ψ as a strong e -enumeration doesn't abandon $V^{(i)}$ on a set $X^{(i)}$. Then, for any $Y \subseteq \bigcap_{i=1}^m X^{(m)}$, Ψ^Y is not total or is not a strong e -enumeration. (As Fact 3.2 this also holds for the relativized version.)

Together with Fact 3.4, the following two facts tell us how to apply the *Cross* operation in order to ensure that T_s is an ordered-partition-tree.

Fact 3.5. Let $e_1, e_2, \dots, e_u$ be u many positive numbers and let $k' = \sum_{i=1}^u e_i$. If $V^{(1)}, V^{(2)}, \dots, V^{(n)}$ is a k' -disperse sequence of clopen sets, for each $1 \leq i \leq u$, let $\mathcal{K}_i = \{K \subseteq \{1, 2, \dots, n\} : \{V^{(j)}\}_{j \in K} \text{ is an } e_i\text{-disperse class}\}$. Then $\vec{\mathcal{K}} = (\mathcal{K}_1, \dots, \mathcal{K}_u)$ is a u -supporter of $\{1, 2, \dots, n\}$.

Proof. Suppose, on the contrary, there is an ordered partition $P^{(1)}, P^{(2)}, \dots, P^{(u)}$ of $\{1, 2, \dots, n\}$ such that for all $1 \leq i \leq u$, $P^{(i)} \notin \mathcal{K}_i$, i.e. $\{V^{(j)}\}_{j \in P^{(i)}}$ is not an e_i -disperse sequence of clopen sets. Then for each i , there exists a partition $P^{(i,1)} \cup P^{(i,2)} \cup \dots \cup P^{(i,e_i)} = P^{(i)}$ such that $(\forall e \leq e_i)(\bigcap_{j \in P^{(i,e)}} [V^{(j)}]^\perp \neq \emptyset)$. However, then $P^{(1,1)}, P^{(1,2)}, \dots, P^{(1,e_1)}, \dots, P^{(i,j)}, \dots, P^{(K,e_u)}$ is a k' -partition of $\{1, 2, \dots, n\}$ that contradicts the assumption that $V^{(1)}, V^{(2)}, \dots, V^{(n)}$ is a k' -disperse class of clopen sets. $\square$

By the definition of u -supporter we have:

Fact 3.6. Let $\vec{\mathcal{K}} = (\mathcal{K}_1, \dots, \mathcal{K}_u)$ be a u -supporter of $\{1, 2, \dots, n\}$, let the sets $X^{(1)}, X^{(2)}, \dots, X^{(n)}$ be ordered u -partitions of W , and let $K' = \sum_{i=1}^K |\mathcal{K}_i|$. Then $Cross(X^{(1)}, X^{(2)}, \dots, X^{(n)}; \vec{\mathcal{K}})$ is an ordered K' -partition of W .

Proof. This is straightforward by the definition of u -supporter. We show that for each $x \in W$ there exists an i and $K_{i,j} \in \mathcal{K}_i$, such that $x \in \bigcap_{p \in K_{i,j}} \pi_i^u(X^{(p)})$. Note that for each $m \leq n$, x belongs to some part of $X^{(m)}$, and therefore $P^{(i)} = \{m \leq n : x \in \pi_i^u(X^{(m)})\}$, $1 \leq i \leq u$, is a u -partition of $\{1, 2, \dots, n\}$. By the definition of u -supporter, there exists an i and $K_{i,j} \in \mathcal{K}_i$ such that $K_{i,j} \subseteq P^{(i)}$, which implies that $x \in \bigcap_{p \in K_{i,j}} \pi_i^K(X^{(p)})$. $\square$

Furthermore, the following plain fact tells us what kind of finite extension of an initial segment requirement is allowed.

Fact 3.7. If $\Psi^{C \oplus \rho}$ doesn't abandon V on Y and $\sigma \supset \rho$ is such that $set(\sigma) \subseteq Y \cup set(\rho)$, then $\Psi^{C \oplus \sigma}$ also doesn't abandon V on Y . Furthermore, if $\vec{\Psi}^{C \oplus \vec{\rho}}$ doesn't abandon V on $X = \bigoplus_{i=1}^k X_k$ and $\vec{\sigma} \supset \vec{\rho}$ is such that $(\forall i \leq k) set(\sigma_i) \subseteq X_i \cup set(\rho_i)$, then $\vec{\Psi}^{C \oplus \vec{\sigma}}$ also doesn't abandon V on X .

3.4. Step s: Construct $(\vec{\rho}^s, T_s)$. Suppose we have a C -computable ordered k_{s-1} -partition tree T_{s-1} , for some $C \in \mathcal{M}$ (here we mean a k -partition of some $W =^* \omega$; note that in above argument T_1 is a 6-partition tree of ω) and we need to construct T_s ensuring that on each child part, either the left-hand side or the right-hand side steps forward; i.e. if G satisfies $(\rho_{il}, \pi_i^k(X))$ for some $X \in [T_{s-1}]$, then $\Psi_{il}^{C^{(j)} \oplus G}$ is not a strong $e_{i,l}^{s-1,j}$ -enumeration of Q (where $e_{i,l}^{s-1,j}$ is the index of Ψ_{il} as updated at step $s-1$), and if G satisfies $(\rho_{ir}, \pi_i^k(X))$ for some $X \in [T_{s-1}]$, then $\Psi_{ir}^{C^{(j)} \oplus G}$ is not a strong $e_{i,r}^{s-1,j}$ -enumeration of Q .

Case i. If there exists ρ and i satisfying either

$$\begin{aligned} \rho \supset \rho_{il} \wedge (\exists n, |\Psi_{il}^{C^{(j)} \oplus \rho}(n)| > e_{i,l}^{s-1,j} \vee [\Psi_{il}^{C^{(j)} \oplus \rho}(n) \downarrow] \cap [Q] = \emptyset) \\ \wedge (set(\rho) \subseteq A) \wedge (\exists X \in [T], set(\rho) \subseteq \pi_i^k(X)/\rho_{il}) \end{aligned}$$

or

$$\rho \supset \rho_{ir} \wedge (\exists n, |\Psi_{ir}^{C(j) \oplus \rho}(n)| > e_{ir}^{s-1,j} \vee [\Psi_{ir}^{C(j) \oplus \rho}(n) \downarrow] \cap [Q] = \emptyset) \\ \wedge (set(\rho) \subseteq \bar{A}) \wedge (\exists X \in [T], set(\rho) \subseteq \pi_i^k(X)/\rho_{ir}),$$

then we can extend ρ_{il} or ρ_{ir} to ρ , which ensures that either $\Psi_{il}^{C(j) \oplus G}$ or $\Psi_{ir}^{C(j) \oplus G}$ is not a strong constant-bound-enumeration of Q , assuming G satisfies $\langle \rho_{il}, \pi_i^k(X) \rangle$ or $\langle \rho_{ir}, \pi_i^k(X) \rangle$, respectively, for some $X \in [T]$. Note that this operation doesn't increase the total number of parts and that the new Π_1^0 class $[T]$ is still non-empty since $\exists X \in [T]$ such that $set(\rho) \subseteq \pi_i^k(X)/\rho_{il}$ (or similarly for the right-hand side). During the construction the above process will be repeatedly carried out within Step s until Case i fails to occur, which must happen as proved in Lemma 4.4.

Case ii. If Case i fails to occur. Let $k' = \sum_{i=1}^k (e_{i,l}^{s-1,j} + e_{i,r}^{s-1,j})$, where $k = k_{s-1}$, and each $e_{i,r}^{s-1,j}$ is as updated in previous applications of Case i this stage, if any, i.e. the present progress for $R(\cdots; j)$, part i . Try to find a k' -disperse sequence of clopen sets $V^{(1)}, V^{(2)}, \dots, V^{(n)}$ (in Step 1, $k' = 1 + 1 = 2$) such that the following classes are non-empty for $m \leq n$:

$$[T_{V^{(m)}}] = \{X = \bigoplus_{i=1}^k (X_{il} \oplus X_{ir}) : \bigoplus_{i=1}^k (X_{il} \cup X_{ir}) \in [T_s] \\ \wedge \vec{\Psi}^{C(j) \oplus \vec{\rho}} \text{ doesn't abandon } V^{(m)} \text{ on } \bigoplus_{i=1}^k (X_{il} \oplus X_{ir})\}.$$

The above condition means that

$$(\forall i \leq k \forall G \subseteq X_{il} \forall n) (|\Psi_{il}^{C(j) \oplus G/\rho_{il}}(n)| \leq e_{i,l}^{s-1,j} \\ \wedge (\Psi_{il}^{C(j) \oplus G/\rho_{il}}(n) \uparrow \vee [\Psi_{il}^{C(j) \oplus G/\rho_{il}}(n)]^\preceq \cap [V^{(m)}]^\preceq \neq \emptyset)),$$

and similarly for the right-hand side. Note that each $[T_{V^{(m)}}]$ is a $\Pi_1^{0,C(j)}$ class of ordered $2k$ -partitions.

For all $1 \leq i \leq 2k$, if i is even, then let $i' = \frac{i}{2}$ and define $\mathcal{K}_i = \{K \subseteq \{1, 2, \dots, n\} : \{V^{(m)}\}_{m \in K} \text{ is an } e_{i',l}^{s-1,j}\text{-disperse class}\}$; if i is odd, then let $i' = \frac{i+1}{2}$ and define $\mathcal{K}_i = \{K \subseteq \{1, 2, \dots, n\} : \{V^{(m)}\}_{m \in K} \text{ is an } e_{i',r}^{s-1,j}\text{-disperse class}\}$. Note that by Fact 3.5, $\vec{\mathcal{K}} = (\mathcal{K}_1, \dots, \mathcal{K}_{2k})$ is a $2k$ -supporter of $\{1, 2, \dots, n\}$.

Let

$$T_s = Cross(T_{V^{(1)}}, T_{V^{(2)}}, \dots, T_{V^{(n)}}; \vec{\mathcal{K}}),$$

i.e.

$$\forall X \in [T_s], X = \bigoplus_{i \leq 2k} (\bigoplus_{j \leq |\mathcal{K}_i|} Y_{K_{i,j}}),$$

where

$$Y_{K_{i,j}} = \bigcap_{p \in K_{i,j}} \pi_i^{2k}(X^{(p)}) \text{ for } X^{(p)} \in [T_{V^{(p)}}].$$

Then we try to satisfy positive requirements by appropriately extending the present initial segment requirements which were updated in previous applications of Case i of the P-Operation in the last stage. Then we update progress information (i.e. update $\vec{e}^j$).

Note:

- (1) $[T_{V^{(m)}}]$ is a $\Pi_1^{0, C \oplus C^{(j)}}$ class.
- (2) If $[V]^\preceq \supseteq [Q]$, then $[T_V] \neq \emptyset$, as otherwise Case i occurs.
- (3) For all i , if i is even, then $\Psi_{il}^{C^{(j)} \oplus G / \rho_{il}}$ is not a strong $e_{il}^{s-1, j}$ -enumeration of Q for any $G \subseteq Y_{K_{i,j}}$, and if i is odd, then $\Psi_{ir}^{C^{(j)} \oplus G / \rho_{ir}}$ is not a strong $e_{ir}^{s-1, j}$ -enumeration of Q for any $G \subseteq Y_{K_{i,j}}$ (by Fact 3.4 and the definition of $\vec{\mathcal{K}}$); i.e. on each child part $R(\cdots; j)$ steps forward on either the left-hand side or the right-hand side.
- (4) $[T_s]$ is a $\Pi_1^{0, C \oplus C^{(j)}}$ class.
- (5) $\bigcup_{i \leq 2k} (\bigcup_{j \leq |\mathcal{K}_i|} Y_{K_{i,j}}) = W$; i.e. T_s is an ordered- K' -partition tree, where $K' = \sum_{i=1}^K |\mathcal{K}_i|$. (See Fact 3.6; this is why we choose a k' -disperse class of clopen sets at *this* step.)

How the requirement $R(e; j)$ is satisfied: For some path $X \in [T_s]$, G will be contained in some $\pi_{i'}^{K'}(X)$, and we will have $G \supset \rho_{i'l}$ or $G \supset \rho_{i'r}$. Therefore either $R(e_{il}^{s-1, j}; j)$ or $R(e_{ir}^{s-1, j}; j)$ is satisfied on part i' of T_s , where part i of T_{s-1} is the parent of part i' of T_s .

4. CONSTRUCTION AND VERIFICATION

Given a countable class $\mathcal{M}$ closed under disjoint union, a set A , and a closed set $[Q]$ of 2^ω satisfying the conditions stated in Theorem 1.2, we give a concrete construction of an infinite set G with $G \subseteq A \vee G \subseteq \bar{A}$ such that $\forall C \in \mathcal{M}, G \oplus C$ does not compute a constant-bound-enumeration of Q , and we prove that the construction provides the desired G .

If $A \leq_T C^{(j)}$ for some $C^{(j)} \in \mathcal{M}$, then just let $G = A$ if A is infinite, and $G = \bar{A}$ otherwise. So without loss of generality assume $\forall C \in \mathcal{M}, A \not\leq_T C$.

4.1. Some operations. Suppose we are given $A \in 2^\omega$, and let T be an ordered k -partition tree of $2^{<\omega}$, with $\vec{\rho}$ being the corresponding initial segment requirement.

To satisfy positive requirements we will apply the P-Operation.

Definition 4.1 (P-Operation). The P-Operation applied to the left-hand side of T 's part i : Choose $\rho \supset \rho_{il}$ (if one exists) such that $[T_{\rho, i}] = \{X \in [T] : \text{set}(\pi_i^k(X)) \supset \text{set}(\rho)\} \neq \emptyset$ and $\emptyset \neq \text{set}(\rho) - \text{set}(\rho_{il}) \subset A$, and then update initial segments: let $\rho_{il} = \rho / \rho_{il}$, let $\rho_{jl} = \rho_{jl}$ for $j \neq i$, and let $\rho_{ir} = \rho_{ir}$ for all i . Finally let $T = T_{\rho, i}$.

If such a ρ exists, we say the P-Operation succeeds. Otherwise do nothing; in this case we say the P-Operation fails.

The P-Operation applied to the right-hand side of T 's part i is analogous, with A replaced by $\bar{A}$.

To satisfy $R(\cdots; j)$ we will apply the R-i-Operation or the R-ii-Operation depending on which of Case i or Case ii occurs.

Definition 4.2 (R-i-Operation to $(T, \vec{\rho}, j, \vec{\Psi} = \vec{e}^{s-1,j})$). Choose the least $i \leq k$ for which either

- there exists a $\rho \supset \rho_{il}$ such that

$$(\exists n, |\Psi_{il}^{C(j) \oplus \rho}(n)| > e_{il}^{s-1,j} \vee [\Psi_{il}^{C(j) \oplus \rho}(n) \downarrow] \cap [Q] = \emptyset) \\ \wedge (set(\rho) \subseteq A) \wedge (\exists X \in [T], set(\rho) \subseteq \pi_i^k(X)/\rho_{il})$$

or

- there exists a $\rho \supset \rho_{ir}$ satisfying a similar condition as above but with $set(\rho) \subseteq A$ replaced by $set(\rho) \subseteq \bar{A}$ and “ il ” replaced by “ ir ”.

If such a ρ exists we say Case i occurs to $R(\cdots; j)$. For each subcase we say that the left-hand side of part i steps forward and that $R(\cdots; j)$ steps forward on the left-hand side; or that the right-hand side of part i steps forward and that $R(\cdots; j)$ steps forward on the right-hand side. We say that $R(\cdots; j)$, part i steps forward no matter which side steps forward.

Then update initial segment requirements (according to ρ, i), i.e. set $\rho_{il}(\rho_{ir}) = \rho$ in the first (second) subcase, and update progress information:

$$e_{il}^{s-1,m} = e_{il}^{s-1,m} \text{ and } e_{ir}^{s-1,m} = e_{ir}^{s-1,m} \text{ if } m \neq j, \\ e_{ir}^{s-1,j} = e_{ir}^{s-1,j} + 1 \text{ and } e_{il}^{s-1,j} = e_{il}^{s-1,j} \text{ if the right-hand side steps forward,} \\ e_{il}^{s-1,j} = e_{il}^{s-1,j} + 1 \text{ and } e_{ir}^{s-1,j} = e_{ir}^{s-1,j} \text{ if the left-hand side steps forward}$$

and $\vec{\Psi} = \vec{e}^{s-1,j}$ in any case.

Definition 4.3 (R-ii-Operation to $(T, \vec{\rho}, j, \vec{\Psi} = \vec{e}^{s-1,j})$). For a clopen set W , let

$$[T_W] = \{X = \bigoplus_{i=1}^k (X_{il} \oplus X_{ir}) : \bigoplus_{i=1}^k (X_{il} \cup X_{ir}) \in [T] \\ \wedge \vec{\Psi}^{C(j) \oplus \vec{\rho}} \text{ doesn't abandon } W \text{ on } \bigoplus_{i=1}^k (X_{il} \oplus X_{ir})\} \neq \emptyset.$$

In this operation we find a k' -disperse sequence of clopen sets $V^{(1)}, V^{(2)}, \dots, V^{(n)}$, where $k' = \sum_{i=1}^k e_{il}^{s-1,j} + e_{ir}^{s-1,j}$ such that for all $m \leq n$, $[T_{V^{(m)}}] \neq \emptyset$.

If such a k' -disperse sequence can be found, we say Case ii occurs. We will show in Lemma 4.6 that if Case i fails to occur, then Case ii must occur, as otherwise we obtain a $C^{(j)}$ -computable constant-bound-enumeration of Q .

Define $\vec{\mathcal{K}}$: for $i = 1, 2, \dots, 2k$,

- (1) if i is even, then let $i' = \frac{i}{2}$ and define

$$\mathcal{K}_i = \{K \subseteq \{1, 2, \dots, n\} : \{V^{(m)}\}_{m \in K} \text{ is an } e_{i',l}^{s-1,j} \text{-disperse class}\};$$

- (2) if i is odd, then let $i' = \frac{i+1}{2}$ and define

$$\mathcal{K}_i = \{K \subseteq \{1, 2, \dots, n\} : \{V^{(m)}\}_{m \in K} \text{ is an } e_{i',r}^{s-1,j} \text{-disperse class}\};$$

and let $u = \sum_{i=1}^{2k} |\mathcal{K}_i|$.

Update T : $T = T_s = \text{Cross}(T_{V^{(1)}}, T_{V^{(2)}}, \dots, T_{V^{(n)}}; \vec{\mathcal{K}})$; update $k = k_s = u$. (Note that each new part is contained in an old part of some path through T .)

Call a new part i' a left child of an old part i iff for any $X \in [T_s]$, $\pi_{i'}^{k_s}(X)$ is the intersection of $\pi_{il}^{k_{s-1}}(X^{(p)})$, $p \in K_{2i,m}$ for some $K_{2i,m}$, where $X^{(p)} \in [T_{V(p)}]$; and similarly for the right-hand side with $2i$ replaced by $2i - 1$.)

Update progress information: for $i = 1, 2, \dots, u$, let

$$\begin{aligned} e_{il}^{s,m} &= e_{il}^{s-1,m}, \quad e_{ir}^{s-1,m} = e_{ir}^{s-1,m} \text{ if } m \neq j, \\ e_{ir}^{s,j} &= e_{ir}^{s-1,j} + 1, \quad e_{il}^{s-1,j} = e_{il}^{s-1,j} \text{ if part } i \text{ of } T_s \text{ is a right child of part } i' \text{ of } T_{s-1}, \\ e_{il}^{s,j} &= e_{il}^{s-1,j} + 1, \quad e_{ir}^{s-1,j} = e_{ir}^{s-1,j} \text{ if part } i \text{ of } T_s \text{ is a left child of part } i' \text{ of } T_{s-1}. \end{aligned}$$

Finally, assign the value $\vec{e}^{s,t_{s+1}}$ to $\vec{\Psi}$.

4.2. Construction. Let $C^{(j)}$ be a sequence of sets of $\mathcal{M}$ that is cofinal in $\mathcal{M}$, i.e. $(\forall j \in \mathbb{N})[C^{(j)} \in \mathcal{M} \wedge (\forall C \in \mathcal{M} \exists C^{(j)}, C \leq_T C^{(j)})]$; furthermore, we require that $\forall j \in \mathbb{N}, C^{(j)} \leq_T C^{(j+1)}$. Fix a sequence of elements of $\{C^{(j)}\}_{j \in \mathbb{N}}$, namely $C^{(t_s)}$, such that each $C^{(j)}$ appears infinitely often in this sequence, i.e. $\exists^\infty s, t_s = j$.

Now suppose we are at the beginning of Step s , we have a C -computable k_{s-1} -partition tree $T = T_{s-1}$ with $[T] \neq \emptyset$, and $C^{(j)}$ is the corresponding set in the s^{th} position of the sequence (i.e. $t_s = j$, so Step s is devoted to $R(\dots; j)$), with current progress $\vec{\Psi} = \vec{e}^{s-1,j}$ for $R(\dots; j)$ and initial segment requirements $\vec{\rho}$.

Begin Step s of the construction:

- (1) Repeatedly apply the R-i-Operation to $(T, \vec{\rho}, j, \vec{\Psi} = \vec{e}^{s-1,j})$ until Case i fails to occur to $R(\dots; j)$.
(We will show in Lemma 4.4 that after finitely many repetitions Case i will stop occurring).
- (2) Then apply the R-ii-Operation to $(T, \vec{\rho}, j, \vec{\Psi} = \vec{e}^{s-1,j})$.
(We will show in Lemma 4.6 that if Case i doesn't occur, then Case ii must occur, as otherwise we obtain a strong constant-bound-enumeration of Q .)
- (3) For each part i , $i \leq k_s$, if part i is a left child, then apply the P-Operation to the left-hand side of T_s 's part i ; otherwise apply the P-Operation to the right-hand side of T_s 's part i .
(Here the P-Operation is applied to each part in order of their indices. We will show in Lemma 4.5 that the P-Operation succeeds on at least one of the new parts. Note that on each "new" part at least one side of $R(\dots; j)$ has just stepped forward.)
- (4) Finally, go to the next step.

Note that in either case $R(\dots; j)$ only steps forward but not "backward", since each child part is a subset of its parent part, and the initial segment requirements are inherited.

4.3. Verification. We need the following three facts as mentioned above.

Lemma 4.4. *Within each step, Case i will not occur to $R(\dots; j)$ forever.*

Lemma 4.5. *For any A and any C -computable ordered k -partition tree T such that $[T] \neq \emptyset$, if $A \not\leq_T C$, then there exist $i \leq k$ and $X \in [T]$ such that both $\pi_i^k(X) \cap A$ and $\pi_i^k(X) \cap \bar{A}$ are non-empty, which implies that in Case ii, for at least one of the new parts, the P-Operation succeeds on the side of that part to which it is applied.*

Lemma 4.6. *Let E be a set of finite subsets of $2^{<\omega}$ c.e. in $\mathbf{e}$ such that, for all n :*

- (1) $\mathcal{W}_n = \{W \subseteq 2^{<\omega} : \forall \rho \in W, |\rho| = n \wedge W \not\subseteq E\}$ *is not a k' -disperse class.*
- (2) $\mathcal{Q}_n = \{\rho \in 2^{<\omega} : |\rho| = n \wedge [\rho] \cap [Q] \neq \emptyset\} \notin E$.

Then there exists a strong k' -enumeration $h : \omega \rightarrow \omega$ of Q computable in $\mathbf{e}$.

Actually, we could further require that $(\forall \rho \in D_{h(n)})|\rho| = n$. This implies that for any specific tuple $(T, \vec{\rho}, j, \vec{\Psi} = \vec{e}^{s-1,j})$ in the construction, either Case i or Case ii occurs since otherwise $E = \{W \subseteq 2^{<\omega} : \forall \rho, \sigma \in W, |\rho| = |\sigma|, [T_W] = \emptyset\}$ is a $C \oplus C^{(j)}$ -c.e. set which satisfies (1) and (2) (recall item (2) in the Note to Step s and the definition of R-ii-Operation).

Assuming these facts we now show that the construction does provide the desired G . Note that our construction yields a tree $\mathcal{T}$: Each node at the s^{th} level represents a part at the s^{th} step and corresponds to a tuple $(j; e_l, e_r)$ standing for the progress of Step s for $C^{(j)}$ (on this part), where $C^{(j)}$ is the set that this step is devoted to, i.e. the j such that $R(\cdots; j)$ steps forward at this step. The successor nodes of a node c are those representing parts that are descendant parts of the part that c represents. Clearly $\mathcal{T}$ is a finitely branching tree.

Definition 4.7. Say part i fades away on the left-hand side at Step s iff $\forall X \in [T_s], \pi_i^k(X) \cap A = \emptyset$ (and similarly for the right-hand side with $\bar{A}$ replacing A).

Note that if part i 's left-hand side fades away at Step s , then it fades away forever; i.e. all its descendants also fade away at the left-hand side. So by Lemma 4.5, there exists an infinite subtree $\mathcal{T}'$ of $\mathcal{T}$ such that each s^{th} level node of $\mathcal{T}'$ represents a part that has not yet faded away on either side at Step s . Therefore there exists a path f through this subtree. Note that for each $R(\cdots; j)$ there exists some side, say the right-hand side, such that along f , $R(\cdots; j)$ steps forward on this side infinitely often. Therefore there exists some side, say the right-hand side, such that for any $C^{(j)}$ there exists $C^{(j')}$ with $C^{(j)} \leq_T C^{(j')}$, and there exist infinitely many steps $s_m, m \in \mathbb{N}$, such that step s_m is devoted to $R(\cdots; j')$, $f(s_m)$'s represented part f_{s_m} steps forward at the right-hand side at step s_m , and the following P-Operation (applied to the new part corresponding to this part's right-hand side) succeeds. In other words, for any $C \in \mathcal{M}$ there exists $C^{(j')}$ such that $C \leq_T C^{(j')}$ and along f , $R(\cdots; j')$ steps forward on the right-hand side infinitely often (which implies that all $R(j; e_r)$, $e_r \in \mathbb{N}$ are satisfied) and the P-Operation succeeds (on the right-hand side) infinitely often. Therefore, $G = \bigcup_{s=1}^{\infty} \rho_{f_s, r}^s$ is infinite and $C^{(j')} \oplus G$ doesn't compute a constant-bound-enumeration of Q , so that $G \oplus C \leq_T G \oplus C^{(j')}$ also fails to compute a strong constant-bound-enumeration of Q , i.e. $\forall C \in \mathcal{M}, G \oplus C$ doesn't compute a strong constant-bound-enumeration of Q .

It remains to prove the three facts mentioned above.

Proof of Lemma 4.4. Recall that there are infinitely many Turing functionals Ψ_e such that, for any oracle $C \oplus G$, $\Psi_e^{C \oplus G}$ doesn't halt on any input. Call such Turing functionals trivial. Clearly, at Step s , if a part proceeds to (e_l, e_r) (during the Case i loop), where Ψ_{e_l} is a trivial Turing functional, then the R-i-Operation will never succeed on the left-hand side of this part, and similarly for the right-hand side if Ψ_{e_r} is trivial. Since there are only finitely many different parts and sides, and the R-i-Operation doesn't increase the total number of parts, either Case i fails to occur during the loop or all Turing functionals in the progress tuples will finally

be updated to trivial ones, which also causes Case i to fail to occur, since Case i occurring implies that at least some Turing functional halts on some input. $\square$

Proof of Lemma 4.5. Suppose on the contrary that $(\forall i \leq k) ([\forall X \in [T], \pi_i^k(X) \cap A = \emptyset] \vee [\forall X \in [T], \pi_i^k(X) \cap \bar{A} = \emptyset])$. Let $R = \{i \leq k : \forall X \in [T], \pi_i^k(X) \cap A = \emptyset\}$ and L be the set of all $i \leq k$ that are not in R . We show that A is computable in C . Note that T is a partition of $W =^* \omega = \omega - F$ for some finite set F , i.e. $\forall X \in [T], \bigcup_{i=1}^k \pi_i^k(X) \supseteq \omega - F$. To decide whether $n \in A$, suppose $n \notin F$, wait (using C) for a moment such that for some $m, \forall \rho, (|\rho| = m \wedge \rho \in T) \Rightarrow [(\forall i \in L, n \notin \pi_i^k(\rho)) \vee (\forall i \in R, n \notin \pi_i^k(\rho))] \vee [\exists N \in \mathbb{N}, \forall \sigma \supset \rho, |\sigma| = N \rightarrow \sigma \notin T]$. Note that such a moment must exist by the assumption that each part is contained in either A or $\bar{A}$, so $n \in A \rightarrow [(\forall i \in R, n \notin \pi_i^k(\rho)) \vee (\exists N \in \mathbb{N}, \forall \sigma \supset \rho, |\sigma| = N \rightarrow \sigma \notin T)]$ and similarly for $n \in \bar{A}$. Furthermore, since $\forall X \in [T], \bigcup_{i=1}^k \pi_i^k(X) \supseteq \omega - F$, we have $(\forall i \in L, n \notin \pi_i^k(\rho)) \rightarrow n \in \pi_j^k(\rho)$ for some $j \in R$ and $\rho \subset X \in [T]$. Therefore $n \in A$ iff $\exists m \forall \rho, (|\rho| = m \wedge \rho \in T) \Rightarrow (\forall i \in R, n \notin \pi_i^k(\rho))$ and $n \in \bar{A}$ iff $\exists m \forall \rho, (|\rho| = m \wedge \rho \in T) \Rightarrow (\forall i \in L, n \notin \pi_i^k(\rho))$. So we could compute $A - F$ using C , which is clearly equivalent to $A \leq_T C$. $\square$

Proof of Lemma 4.6. We first show that if neither Case i nor Case ii occurs, then $E = \{W \subseteq 2^{<\omega} : \forall \rho, \sigma \in W, |\rho| = |\sigma|, [T_W] = \emptyset\}$ is $C \oplus C^{(j)}$ -c.e. and satisfies (1) and (2). Note that E is clearly $C \oplus C^{(j)}$ -c.e.

Clearly, if Case ii does not occur, then E satisfies (1) by the definition of “Case ii occurs”. Suppose for some $Q_n, [T_{Q_n}] = \emptyset$. Consider an arbitrary $X \in [T]$. For $i \leq k$ let $X_{il} = \pi_i^k(X) \cap A$ and $X_{ir} = \pi_i^k(X) \cap \bar{A}$. Since $[T_{Q_n}] = \emptyset, X' = \bigoplus_{i \leq k} (X_{il} \oplus X_{ir}) \notin [T_{Q_n}]$. Therefore on some part $i, \vec{\Psi}_i^{C^{(j)} \oplus \rho_i}$ abandons Q_n on $X_{il} \oplus X_{ir}$. Thus, by the definition of abandoning, item (2), either $(\exists \rho \supset \rho_{il}, \text{set}(\rho) \subseteq X_{il} \subseteq A, \exists n \in \mathbb{N})[\Psi_i^{C^{(j)} \oplus \rho}(n) \downarrow] \cap [Q_n] = \emptyset \vee |\Psi_{il}^{C^{(j)} \oplus \rho}(n)| > \text{index of } \Psi_{il})$ or $(\exists \rho \supset \rho_{ir}, \text{set}(\rho) \subseteq X_{ir} \subseteq \bar{A}, \exists n \in \mathbb{N})[\Psi_{ir}^{C^{(j)} \oplus \rho}(n) \downarrow] \cap [Q_n] = \emptyset \vee |\Psi_{il}^{C^{(j)} \oplus \rho}(n)| > (\text{index of } \Psi_{il})$. However, this implies that Case i occurs.

Now we show that the algorithm of E (or equally speaking, the degree $\mathbf{e}$) can be used to compute a strong k -enumeration of Q . Let E_t denote E at stage t and $\mathcal{W}_{n,t} = \{W \subseteq 2^{<\omega} : \forall \rho \in W, |\rho| = n \wedge W \notin E_t\}$. Note that since E is an enumeration, $\mathcal{W}_{n,t+1} \subseteq \mathcal{W}_{n,t}$. Therefore, if there exists a k' -partition of $\mathcal{W}_n, P^{(1)} \cup P^{(2)} \cup \dots \cup P^{(k')} = \mathcal{W}_n$, such that $(\forall i)(\bigcap_{W \in P^{(i)}} W \neq \emptyset)$, then this partition can be found in a finite amount of time. Furthermore, $Q_n \in P^{(i)}$ for some i , so $\forall \rho \in \bigcap_{W \in P^{(i)}} W, \rho \in Q$. It follows that the function h defined below, which is computable using an algorithm for E (i.e. computable in any degree that computes E), is a strong k' -enumeration of Q :

$$D_{h(n)} = \{\rho_1, \rho_2, \dots, \rho_{k'} : \text{there exists some } t$$

$$\text{and a } k'\text{-partition of } \mathcal{W}_{n,t} = \bigcup_{i=1}^{k'} P^{(i)} \text{ s.t. } (\forall j) \left(\bigcap_{W \in P^{(j)}} W \neq \emptyset \right);$$

$$\text{for } i = 1, 2, \dots, k', \rho_i \text{ is the leftmost string in } \bigcap_{W \in P^{(i)}} W \}.$$

$\square$

5. APPLICATIONS

Corollary 5.1. *Over RCA_0 , RT_2^2 does not imply WWKL_0 .*

Before proving this corollary, we make some remarks. It is known that in ω -models WWKL_0 is equivalent to the assertion that for each X there is a set that is Martin-Löf random relative to X . It is known that $X \in 2^\omega$ is Martin-Löf random iff $(\forall n) K(X \upharpoonright n) \geq^+ n$, where $K(\rho)$ denotes the prefix-free Kolmogorov complexity of a string, and furthermore there exists a universal prefix-free machine U such that for every prefix-free machine M , $K_U \leq K_M + c$ for some constant depending on M . For more background on algorithmic randomness theory, see [5, 19]. Fix a universal prefix-free machine U in the following text.

Definition 5.2. A string ρ is c -incompressible $_U$ iff $K_U(\rho) \geq |\rho| - c$.

Note that $[T_c] = \{X \in 2^\omega : \forall n, X \upharpoonright n \text{ is } c\text{-incompressible}_U\}$ is a closed set of 2^ω .

Lemma 5.3. *For any positive integers k, c , there is no computable strong k -enumeration of the c -incompressible $_U$ strings.*

Proof. Suppose on the contrary that $h : \omega \rightarrow \omega$ is a computable strong k -enumeration of the c -incompressible $_U$ strings. Let $\sigma_n = \underbrace{00 \dots 0}_n 1$. We define a prefix-free machine M . The algorithm of M is as follows for each n : look for an $m \geq n^2$, and then output ρ_i on input σ_{kn+i} (note that $j \leq k$). Note that all strings $\rho_1, \dots, \rho_j \in D_{h(m)}$ have length $m = n^2$. Since U is universal, there exists a constant d such that $K_U(\rho_i) < kn + i + d$; therefore, for sufficiently large n , $\forall i \leq k, K_U(\rho_i) < |\rho_i| - c$, contradicting the assumption that h is a strong k -enumeration of the c -incompressible $_U$ strings. $\square$

Note that although Theorem 1.2 proves the cone avoidance result for *one* closed set $[Q]$, it can be easily adapted to construct G cone avoiding countably many closed sets as long as for every finite number of them, $Q_1, Q_2, \dots, Q_n$, the joint union of them, $[\bigvee_{i=1}^n Q_i] = \{X \in 2^\omega : (\exists j \leq n \exists Y \in [Q_j]) X = \rho_j * Y\}$, where ρ_j is determined by Q_j , satisfies the condition in Theorem 1.2.

Now we can prove the main theorem.

Proof of Corollary 5.1. It suffices to construct a countable class $\mathcal{M}$ satisfying four conditions: (a) $C, B \in \mathcal{M} \rightarrow C \oplus B \in \mathcal{M}$; (b) $(C \in \mathcal{M} \wedge B \leq_T C) \rightarrow B \in \mathcal{M}$; (c) $(\forall C \in \mathcal{M}) C$ does not compute a strong constant-bound-enumeration of any T_c ; (d) $\mathcal{M} \models \text{RT}_2^2$.

Note that (a) and (b) ensure that RCA_0 is satisfied, (c) ensures that WWKL_0 is not satisfied, and (d) ensures that RT_2^2 is satisfied.

It is shown in [4, Lemma 7.11] and later corrected in [3] that $\text{RCA}_0 + \text{RT}_2^2$ is equivalent to $\text{RCA}_0 + \text{SRT}_2^\infty + \text{COH}$. Here COH is the axiom saying that for any $C = \bigoplus_{i=0}^\infty C_i \in \mathcal{M}$ there exists an infinite set C^* cohesive for C ; i.e. for any component C_i either $C^* - C_i$ or $C^* - \bar{C}_i$ is finite.

Moreover, we now show that it can be proved by the finite extension method that if $\{R^{(i)}\}_{i \in \mathbb{N}}$ is a sequence such that no finite disjoint union of the $R^{(i)}$ computes a strong constant-bound-enumeration of any T_c , then there exists an infinite set G cohesive for $\{R^{(i)}\}_{i \in \mathbb{N}}$ that also does not compute any strong constant-bound-enumeration of any T_c . To construct such a cohesive set, we try to satisfy the

following requirements.

$$\begin{aligned} R_e : \Psi_e^G \text{ is not a strong } e\text{-enumeration of any } T_c, \\ N_i : G \subseteq^* R^{(i)} \vee G \subseteq^* \omega - R^{(i)}. \end{aligned}$$

Clearly we can inductively define a 0-1 sequence ν such that, letting $R^i = R^{(i)}$ if $\nu(i) = 1$ and $R^i = \omega - R^{(i)}$ if $\nu(i) = 0$, we have $(\forall n) |\bigcap_{i=0}^n R^i| = \infty$.

For each Step s , let $G_s = \bigcap_{i=0}^n R^i$; we construct an initial segment ρ_s such that $\rho_s \supset \rho_{s-1}$, $\text{set}(\rho_s) \subseteq \text{set}(G_{s-1}/\rho_{s-1})$, and furthermore, (ρ_s, G_s) , as a Mathias forcing condition, forces R_s and N_s . It is clear that for any ρ , (ρ, G_s) forces N_s . We now choose $\rho \supset \rho_{s-1} \wedge \text{set}(\rho) \subseteq G_{s-1}$ to satisfy R_s . There are two ways to do so: find a ρ with $\rho \supset \rho_{s-1} \wedge \text{set}(\rho) \subseteq G_{s-1}$ such that $\Psi_e^\rho(n)$ does not halt, or one such that $|\Psi_e^\rho(n) \downarrow| > e \vee \Psi_e^\rho(n) \cap T_c = \emptyset \vee ((\exists \sigma \in \Psi_e^\rho(n)) |\sigma| > n)$. Suppose we can't find such ρ ; this implies that for every n , there exists ρ with $\rho \supset \rho_{s-1} \wedge \text{set}(\rho) \subseteq G_{s-1}$ such that $|\Psi_e^\rho(n) \downarrow| \leq e$ and $\Psi_e^\rho(n)$ is a set of length n strings that has non-empty intersection with T_c . But then one could use this Ψ_e and oracle $\bigoplus_{i=0}^s R^{(i)}$ to compute a strong e -enumeration of T_c , a contradiction.

In the following text, for a class of sets $\mathcal{M}$, we say X is $\mathcal{M}$ -computable iff $\exists C \in \mathcal{M}$ s.t. $X \leq_T C$. The notions of $\mathcal{M}$ -effectively closed, $\Pi_1^{0,\mathcal{M}}$, etc. are defined similarly. Note that the conclusion just proved above is relativizable to any countable class of sets closed under join. Let $\mathcal{M}_0$ = all recursive sets. Let f be a computable stable coloring. By Theorem 1.2, there exists an infinite G_0 with $G_0 \subseteq f_1 \vee G_0 \subseteq f_2$ such that $\forall C \in \mathcal{M}_0$, for each c , $C \oplus G_0$ does not compute any strong constant-bound-enumeration of any infinite subset of any T_c . Let $\mathcal{M}_1 = \{X \in 2^\omega : X \leq_T C \oplus G_0 \text{ for some } C \in \mathcal{M}_0\}$. Clearly $\mathcal{M}_1$ satisfies (a)(b)(c). Let G_1 be cohesive for a sequence of uniformly $\mathcal{M}_1$ -computable sets (where $\mathcal{M}_1$ -computable means computable in some $C \in \mathcal{M}_1$), so that $(\forall C \in \mathcal{M}_1)$ $G_1 \oplus C$ does not compute any strong constant-bound-enumeration of any infinite subset of any T_c . Let $\mathcal{M}_2 = \{X \in 2^\omega : \exists C \in \mathcal{M}_1, X \leq_T C \oplus G_1\}$. Clearly $\mathcal{M}_2$ also satisfies (a)(b)(c). Iterate the above process to ensure that (1) for any uniformly $\mathcal{M}_j$ -computable sequence $C^{(1)}, C^{(2)}, \dots$, there exists an i and a $G_{i-1} \in \mathcal{M}_i$ such that G_{i-1} is cohesive for $C^{(1)}, C^{(2)}, \dots$ and (2) for any $C \in \Delta_2^{0,\mathcal{M}_j}$ ($\Delta_2^{0,\mathcal{M}_j}$ means $\Delta_2^{0,C}$ for some $C \in \mathcal{M}_j$), there exists an i and an infinite $G_{i-1} \in \mathcal{M}_i$ such that $G_{i-1} \subseteq C \vee G_{i-1} \subseteq \bar{C}$. It follows that $\mathcal{M} = \bigcup_{i=0}^\infty \mathcal{M}_i \models \text{RCA}_0 + \text{SRT}_2^2 + \text{COH} \Leftrightarrow \text{RCA}_0 + \text{RT}_2^2$, but clearly $\mathcal{M}$ satisfies (a)(b)(c). The conclusion follows. $\square$

We now prove Corollary 1.11.

Corollary 5.4. *There exists a DNR function that does not compute any binary sequence with positive effective Hausdorff dimension.*

Proof. Again note that the construction can be adapted to construct a set G cone avoiding countably many closed sets if every finite joint union of those sets satisfies the conditions of Theorem 1.2. Therefore, choose a sequence $d_n > 0$ with $\lim_{n \rightarrow \infty} d_n = 0$ and let A be a Δ_2^0 set such that every infinite subset of A or $\bar{A}$ computes a DNR function. Now apply Theorem 1.2 to construct $G \subseteq A \vee G \subseteq \bar{A}$ that cone avoids all corresponding closed sets $[T_{d_n}]$, and let $f \leq_T G$ be a DNR function. Then f does not compute any binary sequence of positive effective Hausdorff dimension. $\square$

For most “natural” trees that have been studied, if the induced closed set has Muchnik degree strictly above 0 as a mass problem, then the tree does not admit a strong constant-bound-enumeration. For example, this fact is true of homogeneous trees.

Definition 5.5. A tree T is homogeneous iff $(\forall \rho_1, \rho_2 \in T \ \forall \rho)(\rho/\rho_1 \in T \Leftrightarrow \rho/\rho_2 \in T)$.

Lemma 5.6. Let $T \subseteq \omega^{<\omega}$ be a finitely branching homogeneous tree, and let $\mathbf{a}$ be a Turing degree such that $[T]$ is $\mathbf{a}$ -effectively closed. Then $\mathbf{a}$ computes a strong k -enumeration of T for some $k \in \mathbb{N}$ iff $\mathbf{a}$ computes a path through T .

Proof. $(\Leftarrow)$: Straightforward, since an enumeration of the initial segments of a path through T is a strong 1-enumeration of T .

$(\Rightarrow)$: We prove this by induction. Let $h \leq_T \mathbf{a}$ be a strong k -enumeration of T . Clearly the conclusion follows when $k = 1$.

Suppose $\exists N \in \mathbb{N} \forall n > N \ D_{h(n)} \subset T$. Then clearly one can compute a path through T as follows: Let $g(n)$ be the first integer g_n we find such that $\exists m > N \exists \sigma \in D_{h(m)}, \sigma(n) = g_n$. Since T is homogeneous, g is a path through T .

Suppose $\forall N \in \mathbb{N} \exists n > N \ D_{h(n)} \not\subset T$. Since $[T]$ is $\mathbf{a}$ -effectively closed, which means $\bar{T}$ is $\mathbf{a}$ -c.e., $\{n \in \mathbb{N} : D_{h(n)} \not\subset T\}$ is an infinite $\mathbf{a}$ -c.e. set. Therefore one can compute a strong $(k-1)$ -enumeration of T as follows: Search for an m and a σ such that $\sigma \in D_{h(m)} \cap \bar{T}_m$ and $\forall \rho \in D_{h(m)} \ |\rho| \geq n$, and let $g(n)$ be the index of $\{\rho \upharpoonright_n : \rho \in D_{h(m)} - \{\sigma\}\}$.

Note that $|D_{g(n)}| \leq k-1$ and $D_{g(n)} \cap T \neq \emptyset$ (as otherwise $D_{h(m)} \cap T = \emptyset$), i.e. $g \in \omega^\omega$ is a strong $(k-1)$ -enumeration of T . The conclusion follows by induction. $\square$

Combining Theorem 1.2 and Lemma 5.6 yields:

Corollary 5.7. Let $\mathcal{M}$ be a countable class closed under disjoint union, and let $Q \subseteq \omega^{<\omega}$ be an infinite $\mathcal{M}$ -computably bounded homogeneous tree such that $[Q]$ is $\mathcal{M}$ -effectively closed. Then

$(\forall A \in 2^\omega)(\exists G \text{ s.t. } (G \subseteq A \vee G \subseteq \bar{A}) \wedge |G| = \infty)(\forall C \in \mathcal{M}, f \in [Q]) \ f \not\leq_T G \oplus C$
if and only if $[Q] \not\leq_u \mathcal{M}$. Here $\leq_u$ denotes the Muchnik reducibility.

Note that the computable boundedness is needed to ensure that in the proof of Lemma 4.6 the class $\mathcal{W}_n$ is $\Pi_1^{0,\mathcal{M}}$ uniformly in n , which makes sure that the enumeration algorithm is computable in $\mathcal{M}$.

For $f \in \mathbb{N}^\mathbb{N}$, $\mathbf{f}\text{-WKL}_0$ as a second order arithmetic statement says: for any set X there exists a $g \in f\text{-DNR}^X$. $\mathbf{WKL}_0$ is $\mathbf{f}\text{-WKL}_0$ restricted to $f \equiv 2$. Using Corollary 5.7, we have:

Corollary 5.8. For any computable $f \in \mathbb{N}^\mathbb{N}$ such that $f(n) > 1$ i.o., $\mathbf{RT}_2^2$ does not imply $\mathbf{f}\text{-WKL}_0$.

Proof. Exactly the same as Corollary 5.1. $\square$

We now mention an interesting result derived by Kjos-Hanssen in algorithmic randomness theory.

Corollary 5.9 (Kjos-Hanssen [13]). For every Martin-Löf random set $A \in 2^\omega$, there exists an infinite set $G \subseteq A \vee G \subseteq \bar{A}$ such that G does not compute any Martin-Löf random set.

Proof. We've already shown in Lemma 5.3 that every finite joint union of the countably many effectively closed sets $[T_c] = \{X \in 2^\omega : \forall n, X \upharpoonright n \text{ is c-incompressible}_U\}$ satisfies the condition of Theorem 1.2. So the conclusion follows from Theorem 1.2 directly. $\square$

6. REMARKS

Remark 6.1. Although at each step of the construction we hold an $\mathcal{M}$ -computable tree T , updated by the R-ii-operation and the P-operation, this is not essential. One could pre-choose a path through T . Nonetheless, the construction tree $\mathcal{T}$ is not avoidable. Furthermore, holding the tree requires less computational power of the constructor.

Remark 6.2. Can we cone avoid an *arbitrary* tree Q if Q is not Muchnik reducible to $\mathcal{M}$? We guess not, and a construction showing that the answer is no might be provided by modifying the proof of Theorem 2.3 of [10], and noting that $Part_k$ reduces a k -enumeration to a 1-enumeration.

Remark 6.3. We have modularized the construction into as many parts as we could, which makes it convenient for further applications and generalizations. For example, by modifying the R-ii-Operation and analyzing the proof of Lemma 4.5, one might strengthen Corollary 5.9 to prove Kjos-Hanssen's result in [14] that any Martin-Löf random set contains an infinite subset that does not compute any Martin-Löf random set.

Another kind of application is to let $Part_k$ serve as a mediate as in Corollaries 1.9 and 1.11; i.e. first one proves that some class $\mathbb{X}$ is Muchnik reducible to $Part_k$, and then it follows that some member of $\mathbb{X}$ does not compute any member of another class.

Remark 6.4. A not so trivial generalization is to reinterpret $\subseteq, \cap$ and therefore overload *Cross* and the definition of "abandon". This is explained as follows.

Say that we can strongly compute a class $\mathcal{Q}$ under the condition $\langle \mathcal{B}, \mathcal{C}(A) \rangle$, where $\mathcal{B}, \mathcal{C}(A)$ are classes, possibly with other "parameters", iff $\exists A \in \mathcal{B} \quad \mathcal{Q} \leq_u \mathcal{C}(A)$. (In this paper $\mathcal{B} = \mathcal{P}(\omega)$, $\mathcal{C}(A) = Part_2(A)$, $\mathcal{Q}$ is a closed set with Q non- k -enumerable in $\mathcal{M}$, and we could not strongly compute $\mathcal{Q}$ under the condition $\langle \mathcal{B}, \mathcal{C}(A) \rangle$ for any A .) Therefore $G \subseteq A$ is generalized as $G \in \mathcal{C}(A)$ and $X \cap Y$ is generalized as $\mathcal{C}(X) \cap \mathcal{C}(Y)$. Note also that in many applications, $\mathcal{B}, \mathcal{C}(A)$ are Π_1^0 classes, for example $\mathcal{B} = \{T \subseteq 2^{<\omega} : T \text{ is a tree satisfying certain property}\}$, say the property of being n -bushy, and $\mathcal{C}(T) = [T]$ (or almost Π_1^0 as $Part_2(A)$, which is a Π_1^0 class minus a countable class of sets). Therefore by redefining "abandon" and *Cross*, the main frame of the construction may still serve for this purpose. (Of course, different combinatorial facts will be needed in different cases and there would be no universal way to produce these combinatorial facts; that belongs to the creative part of mathematics. However, there might be some results with broad application, since what we have studied is only a stone beside the sea.) We hope that this frame could help in the study of homogeneous trees, especially f -DNR, and give universal solutions to various results that have arisen recently, for example in [9, 18]; see also [15].

Question 6.5. The most important question at hand is what kind of Π_3^0 class can be avoided as above. Note that this is related to the question of whether SRT_2^2

implies RT_2^2 , which is equivalent to whether SRT_2^2 implies COH. Here COH is the second order arithmetic statement that every sequence of sets $R^{(i)}$ has a cohesive set for it, i.e. an infinite set Y such that for all $i \in \mathbb{N}$ either $Y \cap R^{(i)}$ or $Y \cap \bar{R}^{(i)}$ is finite. Note that the class of cohesive sets of a given $R^{(i)}$ is a $\Pi_3^{0, R^{(i)}}$ class.

Question 6.6. What is the difference between the Muchnik lattice of Part_2 and that of $\text{Part}_3, \text{Part}_k$? (Here the Muchnik lattice of a collection of subsets of ω^ω refers to the lattice where the $\leq$ relation is induced by the Muchnik reducibility.)

For example, is there a $\text{Part}_3(A_1, A_2)$ such that there exists no $\text{Part}_2(A)$ to which $\text{Part}_3(A_1, A_2)$ is Muchnik reducible? For the latter question we guess the answer is yes.

ACKNOWLEDGEMENT

The author is grateful to Professor Peter Cholak for his help during the preparation of this paper.

REFERENCES

- [1] Klaus Ambos-Spies, Bjørn Kjos-Hanssen, Steffen Lempp, and Theodore A. Slaman, *Comparing DNR and WWKL*, J. Symbolic Logic **69** (2004), no. 4, 1089–1104, DOI 10.2178/jsl/1102022212. MR2135656 (2006c:03061)
- [2] Richard Beigel, Harry Buhrman, Peter Fejer, Lance Fortnow, Piotr Grabowski, Luc Longpre, Andrej Muchnik, Frank Stephan, and Leen Torenvliet, *Enumerations of the Kolmogorov function*, J. Symbolic Logic **71** (2006), no. 2, 501–528, DOI 10.2178/jsl/1146620156. MR2225891 (2007b:68086)
- [3] P.A. Cholak, C.G. Jockusch, and T.A. Slaman, *Corrigendum to: On the strength of Ramsey's theorem for pairs*, J. Symbolic Logic **74** (2009), no. 4, 1438–1439. MR2583829
- [4] ———, *On the strength of Ramsey's theorem for pairs*, Journal of Symbolic Logic **66** (2001), no. 1, 1–55. MR1825173
- [5] Rodney G. Downey and Denis R. Hirschfeldt, *Algorithmic randomness and complexity*, Theory and Applications of Computability, Springer, New York, 2010. MR2732288 (2012g:03001)
- [6] Damir D. Dzhalalov and Carl G. Jockusch Jr., *Ramsey's theorem and cone avoidance*, J. Symbolic Logic **74** (2009), no. 2, 557–578, DOI 10.2178/jsl/1243948327. MR2518811 (2010e:03052)
- [7] Harvey Friedman, *Some systems of second order arithmetic and their use*, Proceedings of the International Congress of Mathematicians (Vancouver, B. C., 1974), Vol. 1, Canad. Math. Congress, Montreal, Que., 1975, pp. 235–242. MR0429508 (55 #2521)
- [8] Harvey Friedman, *Systems of second order arithmetic with restricted induction, i, ii*, Journal of Symbolic Logic **41** (1976), no. 2, 557–559.
- [9] Noam Greenberg and Joseph S. Miller, *Diagonally non-recursive functions and effective Hausdorff dimension*, Bull. Lond. Math. Soc. **43** (2011), no. 4, 636–654, DOI 10.1112/blms/bdr003. MR2820150 (2012g:03112)
- [10] D. R. Hirschfeldt, C. G. Jockusch Jr., B. Kjos-Hanssen, S. Lempp, and T. A. Slaman, *The strength of some combinatorial principles related to Ramsey's theorem for pairs*, Computational Prospects of Infinity: Presented talks **14** (2008), 143–161.
- [11] Jiayi Liu, RT_2^2 does not imply WKL_0 , J. Symbolic Logic **77** (2012), no. 2, 609–620, DOI 10.2178/jsl/1333566640. MR2963024
- [12] Carl G. Jockusch Jr., *Ramsey's theorem and recursion theory*, J. Symbolic Logic **37** (1972), 268–280. MR0376319 (51 #12495)
- [13] Bjørn Kjos-Hanssen, *Infinite subsets of random sets of integers*, Math. Res. Lett. **16** (2009), no. 1, 103–110. MR2480564 (2010b:03051)
- [14] Bjørn Kjos-Hanssen, *A strong law of computationally weak subsets*, J. Math. Log. **11** (2011), no. 1, 1–10, DOI 10.1142/S0219061311000980. MR2833148
- [15] Masahiro Kumabe and Andrew E. M. Lewis, *A fixed-point-free minimal degree*, J. Lond. Math. Soc. (2) **80** (2009), no. 3, 785–797, DOI 10.1112/jlms/jdp049. MR2559129 (2011d:03064)

- [16] Jack H. Lutz, *Gales and the constructive dimension of individual sequences*, Automata, languages and programming (Geneva, 2000), Lecture Notes in Comput. Sci., vol. 1853, Springer, Berlin, 2000, pp. 902–913, DOI 10.1007/3-540-45022-X_76. MR1795945 (2001g:68046)
- [17] Elvira Mayordomo, *A Kolmogorov complexity characterization of constructive Hausdorff dimension*, Inform. Process. Lett. **84** (2002), no. 1, 1–3, DOI 10.1016/S0020-0190(02)00343-5. MR1926330 (2003h:68053)
- [18] Joseph S. Miller, *Extracting information is hard: a Turing degree of non-integral effective Hausdorff dimension*, Adv. Math. **226** (2011), no. 1, 373–384, DOI 10.1016/j.aim.2010.06.024. MR2735764 (2012a:03114)
- [19] André Nies, *Computability and randomness*, Oxford Logic Guides, vol. 51, Oxford University Press, Oxford, 2009. MR2548883 (2011i:03003)
- [20] F. P. Ramsey, *On a Problem of Formal Logic*, Proc. London Math. Soc. **S2-30**, no. 1, 264, DOI 10.1112/plms/s2-30.1.264. MR1576401
- [21] B. Ya. Ryabko, *Coding of combinatorial sources and Hausdorff dimension* (Russian), Dokl. Akad. Nauk SSSR **277** (1984), no. 5, 1066–1070. MR759967 (86g:94033)
- [22] David Seetapun and Theodore A. Slaman, *On the strength of Ramsey's theorem*, Notre Dame J. Formal Logic **36** (1995), no. 4, 570–582, DOI 10.1305/ndjfl/1040136917. Special Issue: Models of arithmetic. MR1368468 (96k:03136)
- [23] Stephen G. Simpson, *Subsystems of second order arithmetic*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1999. MR1723993 (2001i:03126)

DEPARTMENT OF MATHEMATICS, CENTRAL SOUTH UNIVERSITY, CHANGSHA 410083, PEOPLE'S REPUBLIC OF CHINA

E-mail address: g.jiayi.liu@gmail.com