

ESCAPE RATES AND SINGULAR LIMITING DISTRIBUTIONS FOR INTERMITTENT MAPS WITH HOLES

MARK F. DEMERS AND BASTIEN FERNANDEZ

ABSTRACT. We study the escape dynamics in the presence of a hole of a standard family of intermittent maps of the unit interval with neutral fixed point at the origin (and finite absolutely continuous invariant measure). Provided that the hole (is a cylinder that) does not contain any neighborhood of the origin, the surviving volume is shown to decay at polynomial speed with time. The associated polynomial escape rate depends on the density of the initial distribution, more precisely, on its behavior in the vicinity of the origin. Moreover, the associated normalized push forward measures are proved to converge to the point mass supported at the origin, in sharp contrast to systems with exponential escape rate. Finally, a similar result is obtained for more general systems with subexponential escape rates, namely that the Cesàro limit of normalized push forward measures is typically singular, invariant and supported on the asymptotic survivor set.

1. INTRODUCTION AND SETTING

The study of systems with holes finds its origin in the study of Markov chains with absorbing states [26, 36–38] and was introduced in deterministic dynamical systems by Pianigiani and Yorke [33]. It has focused on the establishment of escape rates and on the existence of conditionally invariant measures which describe the asymptotic distribution of mass conditioned on non-escape.

Since conditionally invariant measures are badly non-unique [18], physically relevant measures are usually characterized as the limit of normalized push forward iterates of a reference measure (usually Lebesgue). Such limiting distributions are typically eigenmeasures with maximal eigenvalue of the corresponding transfer operator defined on an appropriate function space. The maximal eigenvalue itself gives the exponential rate of escape of mass from the system. These limiting conditionally invariant measures have properties analogous to Sinai-Ruelle-Bowen (SRB) measures for the corresponding closed system. Under reasonable assumptions, they converge to the SRB measure as the size of the hole tends to zero, and this establishes stability under perturbations in the form of holes.

Examples begin with open systems admitting a finite Markov partition: expanding maps in \mathbb{R}^n [15, 33], Smale horseshoes [7], Anosov diffeomorphisms [8, 9], and some unimodal maps [28]. Subsequent attempts to substitute the Markov assumption by requiring that the holes be small have extended this analysis to Anosov diffeomorphisms with non-Markov holes [10, 11], to expanding maps of the interval [12, 16, 30], to multimodal maps satisfying a Collet-Eckmann condition [3, 17], to

Received by the editors November 6, 2013 and, in revised form, May 8, 2014 and May 30, 2014.
 2010 *Mathematics Subject Classification.* Primary 37C30, 37C40, 37D25, 37E05.

The first author was partially supported by NSF grant DMS 1101572.

The second author was partially supported by EU FET Project No. TOPDRIM 318121.

piecewise hyperbolic maps [19] and recently, to various classes of dispersing billiards [20, 22, 23].

The characteristic common to all these systems is that the rate of escape is exponential (the systems enjoy exponential decay of correlations before the introduction of the hole) so that the concept of conditionally invariant measure is well-defined.

Polynomial rates of escape have been studied numerically in some systems [24, 25] and via formal expansions to obtain leading order terms for the decay rate [4, 5]. However, to our knowledge, there are no analytical results regarding limiting distributions for open systems with polynomial rates of escape.

The purpose of the present paper is to initiate the rigorous mathematical analysis of open systems with subexponential rates of escape. For simplicity, we consider a family of intermittent maps T of the unit interval, with neutral fixed point at the origin [31]. For the hole, we take any element of a refined Markov partition for the map not adjacent to the origin. (Of note, [27] has also considered interval maps with neutral fixed point and very specific holes which are either a neighborhood of the neutral fixed point or its complement.)

In this context, we first prove that the rates of escape must be polynomial for a large class of initial distributions, and this rate depends on the behavior of the initial distribution in a neighborhood of the origin. In particular, the polynomial rate of escape with respect to the SRB measure (for the map before the introduction of the hole) differs from that with respect to Lebesgue measure.

In this setting, conditionally invariant measures are not physically meaningful (although plenty still exist with any desired eigenvalue between 0 and 1 [18]). Letting \hat{T} denote the map with the hole, we show that the limit of $\hat{T}_*^n \mu / |\hat{T}_*^n \mu|$ (NB: for the precise definition of this notation, see section 1.1 below) converges to the point mass at the neutral fixed point for a large class of initial distributions μ (including both Lebesgue and the SRB measures).

These results hold independently of the size of the hole. Thus from the point of view of the physical limit $\hat{T}_*^n \mu / |\hat{T}_*^n \mu|$, a hole of any size is always a large perturbation in the context of subexponentially mixing systems. In other words, the attracting property of the SRB measure under the action of T_*^n is unstable with respect to small leaks in the system.

Finally, we consider more general systems with subexponential rates of escape. The analysis of intermittent maps of the interval might suggest that the results are specific to this setting. Our final result, Theorem 2.3, shows that this is not the case: in contrast to situations with exponential escape, in systems with slow escape, the (Cesàro) limit of $\hat{T}_*^n \mu / |\hat{T}_*^n \mu|$ for reasonable reference measures μ will always be singular and will typically be supported on the survivor set of points that never escape.

1.1. Setting. We study the dynamics of the family of maps of the unit interval $T : I \rightarrow I$ where $I = [0, 1)$ and T is defined by (see [31] and Figure 1)

$$T(x) = \begin{cases} x + 2^\gamma x^{\gamma+1} & \text{if } x \in [0, \frac{1}{2}), \\ 2x - 1 & \text{if } x \in [\frac{1}{2}, 1), \end{cases}$$

with $0 < \gamma < 1$, after the introduction of a hole H into I . In this parameter range, T preserves a finite invariant measure, μ_{SRB} , absolutely continuous with respect to Lebesgue.




FIGURE 1. Graph of the map T for $\gamma = \frac{3}{4}$ (solid red branches), together with some of the intervals $J_n = [a_n, a_{n-1}]$.

In order to define the hole, we need to introduce the (standard) finite and countable Markov partitions of I . The finite partition is defined by $\mathcal{P} := \{J_L, J_R\}$ where $J_L = [0, \frac{1}{2})$ and $J_R = [\frac{1}{2}, 1)$. The countable partition is defined by $\mathcal{J} := \{J_n\}_{n \geq 0}$ where

$$J_n = \begin{cases} J_R & \text{if } n = 0, \\ [a_n, a_{n-1}) & \text{if } n \geq 1, \end{cases}$$

where $a_n = T_L^{-n}(\frac{1}{2})$ and T_L denotes the left branch of T . (Note that $T(J_n) = J_{n-1}$ for all $n \geq 1$; see Figure 1.)

Now, given $t \geq 0$, let $\mathcal{J}^{(t)}$ be the refined partition defined as follows:

$$\mathcal{J}^{(t)} := \mathcal{J} \vee \bigvee_{i=0}^t T^{-i}(\mathcal{P}).$$

The hole H is defined to be any element of $\mathcal{J}^{(\ell_H)}$ where $\ell_H \geq 0$ is arbitrary. We shall denote by $J_h \supseteq H$ with $h \geq 0$ the element of \mathcal{J} that contains H .

This assumption on the hole gives immediate access to a countable Markov partition for the open system. Notice that dynamically refining \mathcal{J} according to $T^{-i}(\mathcal{P})$, and not $T^{-i}(\mathcal{J})$, preserves 0 as the only accumulation point of the endpoints of elements of $\mathcal{J}^{(t)}$ for all $t \geq 0$. In particular, this property is convenient for the conditioning arguments in the proof of Lemma 3.3 and for the invariance of a certain function space used in the proof of Theorem 2.2 that implies control of the structure of the singular limit. Nevertheless, we believe the assumption that H be an element of $\mathcal{J}^{(\ell_H)}$ is purely technical and we expect our results to hold even when relaxed, although significant technical modifications will have to be made.

Define $\dot{I} = I \setminus H$ and given $t \geq 0$, let $\dot{I}^t = \bigcap_{i=0}^t T^{-i}(I \setminus H)$ represent those points which have not escaped by time t (NB: we have $\dot{I}^0 = \dot{I}$). We refer to $\dot{T} := T|_{\dot{I}^1}$ as the map with a hole, and its iterates $\dot{T}^t = T^t|_{\dot{I}^t}$ ($t \geq 1$) describe the dynamics of the open system before escape. Notice that $\mathcal{J}^{(\ell_H)}$ is also a countable Markov partition for \dot{T} .

One of the quantities we will be interested in studying is the rate of escape of mass from the open system. Given a measure μ on \dot{I} , we define the *polynomial rate*

of escape with respect to μ by

$$\mathfrak{e}_{\text{poly}}(\mu) = - \lim_{t \rightarrow +\infty} \frac{\log \mu(\mathring{I}^t)}{\log t}$$

whenever the limit exists.

We will also study the asymptotic evolution of absolutely continuous measures that are transported under the action of \mathring{T} . Given a measure μ on \mathring{I} and $t \geq 1$, let $\mathring{T}_*^t \mu$ be the push forward measure under the action of \mathring{T}^t .

Let m denote Lebesgue measure on I and given $f \in L^1(m)$, let μ_f be the absolutely continuous measure with density f . Let \mathcal{L} be the transfer operator associated with T defined by the expression

$$\mathcal{L}f(x) = \sum_{y \in T^{-1}(x)} \frac{f(y)}{DT(y)},$$

where $DT > 0$ is the (first) derivative of T . Consider the operators $\mathring{\mathcal{L}}^t f := \mathcal{L}^t(f1_{\mathring{I}^t})$ where \mathcal{L}^t are the iterates of \mathcal{L} and 1_A denotes the characteristic function of the set A . We have $\mathring{T}_*^t \mu_f = \mu_{\mathring{\mathcal{L}}^t f}$ for all $t \geq 0$, and the change of variable formula implies in this case the relation

$$|\mathring{\mathcal{L}}^t f|_1 = \mu_f(\mathring{I}^t), \quad \forall t \geq 0,$$

where $|\cdot|_1$ denotes the L^1 -norm with respect to Lebesgue measure m . It will be useful for us later that, with these definitions, the usual composition property of the transfer operators $\mathring{\mathcal{L}}^t$ holds; i.e. for any $j, k \geq 1$,

$$\begin{aligned} \mathring{\mathcal{L}}^j(\mathring{\mathcal{L}}^k f)(x) &= \sum_{y \in T^{-j}(x)} \frac{(\mathring{\mathcal{L}}^k f)(y)}{|DT^j(y)|} 1_{\mathring{I}^j}(y) \\ &= \sum_{z \in T^{-(j+k)}(x)} \frac{f(z)}{|DT^{j+k}(z)|} 1_{\mathring{I}^j}(T^k(z)) 1_{\mathring{I}^k}(z) = \mathring{\mathcal{L}}^{j+k} f(x), \end{aligned}$$

where the last equality follows from the fact that $1_{\mathring{I}^j}(T^k(z)) 1_{\mathring{I}^k}(z) = 1_{\mathring{I}^{j+k}}(z)$.

2. STATEMENT OF RESULTS

2.1. Results for the open system \mathring{T} . Throughout this section, the hole H is fixed as in the previous section (and so are ℓ_H and h). Our first result describes a common set of escape rates for initial distributions depending on their behavior near 0. Following [39], the notation $u_t \approx v_t$ (resp. $u_t \lesssim v_t$, $u_t \gtrsim v_t$) means there exists $C > 0$ such that $C^{-1}v_t \leq u_t \leq Cv_t$ (resp. $u_t \leq Cv_t$, $u_t \geq Cv_t$) for all t . These notations will also be employed as abbreviations for uniform estimates on sequences with multiple indices.

Theorem 2.1. *For any non-negative $f \in L^1(m)$ for which there exist $x_0 \in (0, 1)$ and $\alpha \in [0, 1)$ such that*

$$(1) \quad 0 < \inf_{x \in (0, x_0)} x^\alpha f(x) \leq \sup_{x \in (0, 1)} x^\alpha f(x) < +\infty,$$

we have

$$\mu_f(\mathring{I}^t) \approx t^{-\frac{1-\alpha}{\gamma}}.$$

Consequently, the associated measure μ_f has polynomial escape rate $\mathfrak{e}_{\text{poly}}(\mu_f) = \frac{1-\alpha}{\gamma}$.

The proof is given in Section 3.2. Of note, to obtain the lower bound on $\mu_f(\dot{I}^t)$ is rather immediate (see relation (10)). Moreover, ergodicity of the map T with respect to the absolutely continuous invariant measure μ_{SRB} implies $\mu_f(\dot{I}^t) \xrightarrow{t \rightarrow +\infty} 0$. Thus, most of the proof consists of proving the upper bound. This part is inspired by the proof in [39] of the speed of convergence to the equilibrium measure.

Theorem 2.1 implies in particular that the polynomial escape rate associated with Lebesgue measure is given by $\mathfrak{e}_{\text{poly}}(m) = \frac{1}{\gamma}$. Interestingly, since $\frac{d\mu_{\text{SRB}}}{dm}(x) \approx x^{-\gamma}$ for x near 0 [39], this rate differs from the one associated with the SRB measure, $\mathfrak{e}_{\text{poly}}(\mu_{\text{SRB}}) = \frac{1-\gamma}{\gamma}$.

That the escape rate is polynomial depends on the assumptions both on H and on the initial density f . Indeed, if the hole included a neighborhood of the neutral fixed point 0, then the corresponding open system \dot{T} would be uniformly expanding and the escape rate would be exponential for any initial density $f \in L^1(m)$; see [27, 33] for the Markov case and any of [12, 16, 30] for the non-Markov case. (Obviously, such holes do not belong to $\mathcal{J}^{(t)}$ for any $t \geq 0$.)

Alternatively, $H = [\frac{1}{2}, \frac{1}{2} + d_{\ell_H}) \in \mathcal{J}^{(\ell_H)}$ for some $d_{\ell_H} > 0$ can also create exponential behavior for some initial densities f . Indeed, the map $T|_{[2d_{\ell_H}, 1)}$ is uniformly expanding and no point in $[2d_{\ell_H}, 1)$ can enter the interval $[0, 2d_{\ell_H})$ without first falling into H . Hence, the measure associated with any smooth density f satisfying $f|_{[0, 2d_{\ell_H})} \equiv 0$ must experience an exponential rate of escape in this case. (Such densities do not satisfy the assumption of Theorem 2.1.)

However, for any hole not blocking repeated passes through a neighborhood of 0, $\dot{\mathcal{L}}^t f$ will eventually be positive in a neighborhood of 0 (and bounded) for any ‘typical’ smooth density f ; hence Theorem 2.1 implies that the associated measure will experience a polynomial escape rate $\frac{1}{\gamma}$.

Our next result describes the limiting behavior of the sequence $\left\{ \frac{\dot{T}_*^t \mu_f}{\mu_f(\dot{I}^t)} \right\}_{t \in \mathbb{N}}$ of push forward probability measures for initial densities f that are log-Hölder continuous on elements of the partition $\mathcal{J}^{(\ell_H)}$. To be precise, let $C^0(\mathcal{J}^{(\ell_H)})$ denote the set of functions defined in the interior of I and continuous on each element of $\mathcal{J}^{(\ell_H)}$. Given $f \in C^0(\mathcal{J}^{(\ell_H)})$, $f \geq 0$, $p \in \mathbb{R}^+$ and $J \in \mathcal{J}^{(\ell_H)}$, define the quantity $H_J^p(f)$ as follows:

$$H_J^p(f) = \begin{cases} 0 & \text{if } f \equiv 0 \text{ on } J, \\ +\infty & \text{if } f(x) = 0 < f(y) \text{ for some } x, y \in J, \\ \sup_{x \neq y \in J} \frac{\log f(x) - \log f(y)}{|x - y|^p} & \text{if } f > 0 \text{ on } J, \end{cases}$$

and let $\|f\|_p := \sup_{J \in \mathcal{J}^{(\ell_H)}} H_J^p(f)$. Consider also the set of functions

$$\mathcal{F}_p = \{f \in C^0(\mathcal{J}^{(\ell_H)}) : f \geq 0, \|f\|_1 = 1 \text{ and } \|f\|_p < +\infty\}$$

and its subset

$$\mathcal{F}_p^0 = \{f \in \mathcal{F}_p : \exists x_0 \in (0, 1) \text{ and } \alpha \in [0, 1) \text{ such that (1) holds}\}$$

of functions which are bounded away from zero in a neighborhood of 0.

Theorem 2.2. *Let $f \in \mathcal{F}_p^0$ for some $p > 0$. Then the sequence $\left\{ \frac{\hat{T}_*^t \mu_f}{\mu_f(\hat{I}^t)} \right\}_{t \in \mathbb{N}}$ of absolutely continuous measures behaves asymptotically as follows:*

$$\lim_{t \rightarrow +\infty} \frac{\hat{T}_*^t \mu_f}{\mu_f(\hat{I}^t)} = \delta_0,$$

where δ_0 denotes the point mass at 0 and the convergence is in the weak sense. Moreover, we have

$$\lim_{t \rightarrow +\infty} \frac{\mu_f(\hat{I}^{t+1})}{\mu_f(\hat{I}^t)} = 1.$$

Of note, this last expression of the theorem can be alternatively formulated as

$$\lim_{t \rightarrow +\infty} \frac{\mu_f(\hat{I}^t \setminus \hat{I}^{t+1})}{\mu_f(\hat{I}^t)} = 0.$$

Theorem 2.2 applies in particular to Lebesgue measure, since 1 belongs to \mathcal{F}_p^0 for every $p \in \mathbb{R}^+$. (More generally, one easily checks that any Hölder continuous density with exponent p which is bounded away from 0 on I belongs to \mathcal{F}_p^0 .) Theorem 2.2 also applies to μ_{SRB} since the density $f_{\text{SRB}} = \frac{d\mu_{\text{SRB}}}{dm}$ belongs to \mathcal{F}_p^0 for every $p \in (0, \frac{\gamma}{\gamma+1}]$ as does any normalized density $f/|f|_1$, where $f(x) = x^{-\alpha}$ for some $\alpha \in (0, 1)$ (see Lemma 3.8 in Section 3.3).

As mentioned in the introduction, this theorem implies that arbitrarily small holes in systems with polynomial rates of escape can act as large perturbations from the point of view of the physical limit $\frac{\hat{T}_*^t \mu_f}{\mu_f(\hat{I}^t)}$.

Furthermore, one may also consider the stability of open systems with respect to the location of a hole of a given size [1, 6, 21, 29]. In this framework, consider a family of holes of the form $\{H_{\varepsilon_i}\}$ where $\varepsilon_i > 0$, $\lim_{i \rightarrow +\infty} \varepsilon_i = 0$ and $H_{\varepsilon_i} = [\varepsilon_i, \varepsilon_i + \eta_i)$ all satisfy the assumptions above ($\eta_i > 0$ is small). Then our results state that, for each i , the sequence $\frac{\hat{T}_*^t m}{m(\hat{I}^t)}$ tends to δ_0 for large t . However, for any $n \geq 1$ and $H_0 = [0, a_n)$, i.e. $\varepsilon_i = 0$, the results of [33] imply that the escape rate is exponential and the sequence $\frac{\hat{T}_*^t m}{m(\hat{I}^t)}$ tends to a conditionally invariant measure that is absolutely continuous with respect to Lebesgue. From this point of view, a discontinuity occurs when the hole goes through the neutral fixed point.

2.2. General open systems: Consequence of a subexponential escape rate.

The convergence of $\frac{\hat{T}_*^t \mu_f}{\mu_f(\hat{I}^t)}$ to a singular limit as in Theorem 2.2 is not limited to the map \hat{T} above. Indeed, as we show now, this phenomenon occurs very generally when the rate of escape is subexponential.

To see this, let X be a compact, separable metric space and let $T : X \rightarrow X$ be now an arbitrary Borel measurable map. Assume there exists a Borel probability measure μ with respect to which T is non-singular (but not necessarily invariant). This will be our reference measure.

Let an open set $H \subset X$ be the hole and let $\hat{X}^t = \bigcap_{i=0}^t T^{-i}(X \setminus H)$ denote the survivor set up until time $t \in \mathbb{N} \cup \{+\infty\}$. As before, let $\hat{T} := T|_{\hat{X}^1}$. We have $\hat{T}^t = T^t|_{\hat{X}^t}$ for all $t \geq 1$. Our main assumption on this open system is that μ -almost every point escapes and that the escape rate is subexponential; i.e. we

assume

$$(2) \quad \mu \left(\bigcup_{i=1}^{\infty} \text{int} \left(\bigcup_{j=0}^i T^{-j}(H) \right) \right) = 1, \quad \mu(\dot{X}^t) > 0 \text{ for all } t \geq 0 \quad \text{and} \\ \lim_{t \rightarrow +\infty} \frac{\log \mu(\dot{X}^t)}{t} = 0,$$

where $\text{int}(A)$ denotes the interior of a set A . In particular, this includes both polynomial and stretched exponential rates of escape. We remark that if T is continuous, the first assumption in (2) is equivalent to $\mu(\dot{X}^\infty) = 0$.

By assumption, all push forward (probability) measures $\frac{\dot{T}_*^t \mu}{\mu(\dot{X}^t)}$ are non-singular with respect to μ . Hence, the same is true for

$$(3) \quad \mu_t = \frac{1}{t} \sum_{k=0}^{t-1} \frac{\dot{T}_*^k \mu}{\mu(\dot{X}^k)},$$

for all $t \geq 1$. As the next result shows, any limit point however must be singular. Convergence here is also understood in the weak sense.

Theorem 2.3. *Any limit point μ_∞ of the sequence $\{\mu_t\}_{t \in \mathbb{N}}$ is singular with respect to μ and is supported on $X \setminus \left(\bigcup_{i=1}^{\infty} \text{int} \left(\bigcup_{j=0}^i T^{-j}(H) \right) \right)$. If, in addition, μ_∞ gives zero measure to the discontinuity set of \dot{T} , then μ_∞ is T -invariant and supported on \dot{X}^∞ .*

Interestingly, the averaging method presented here does not work so easily in the case of exponential escape (unless a priori one knows that the limit of $\frac{\dot{T}_*^t \mu}{\mu(\dot{X}^t)}$ itself exists). Indeed, in this case, the ratio of consecutive normalizations $\frac{\mu(\dot{X}^{t+1})}{\mu(\dot{X}^t)}$ does not converge to 1, and the terms appearing in the sum must be weighed to compensate for this. For an example of an averaging method in the exponential case under stronger assumptions, see [13, 14]. Theorem 2.3 is proved in Section 3.6.

3. PROOFS

3.1. Preliminary estimates. In proving the theorems, we shall repeatedly use the following bounds [39]:

$$(4) \quad a_n \approx n^{-\frac{1}{\gamma}} \quad \text{and} \quad |J_n| \approx n^{-\frac{\gamma+1}{\gamma}}.$$

We shall also rely on the following lemma.

Lemma 3.1. *Given $n \geq 0$ and $t \geq 1$, let $x, y \in \dot{I}^t$ lie in the same element of $\mathcal{J}^{(t+1)}$ such that $T^t(x), T^t(y) \in J_n$. Then we have*

$$(a) \quad \frac{1}{DT^t(x)} \lesssim \left(\frac{n+1}{n+t} \right)^{\frac{\gamma+1}{\gamma}}; \\ (b) \quad \text{for any } p \in (0, \frac{\gamma}{\gamma+1}], \text{ one has } \left| \log \frac{DT^t(x)}{DT^t(y)} \right| \lesssim |T^t(x) - T^t(y)|^p.$$

Proof. (a) Given $i \in \{0, \dots, t-1\}$, let J_{n_i} denote the element of \mathcal{J} containing $T^i(x)$ and $T^i(y)$. Also let $B_{n_i} = 2^\gamma \gamma(\gamma+1) a_{n_i}^{\gamma-1}$ be the maximum value of $|D^2 T|$ and $M_{n_i}^{(j)}$ be the minimum value of $|DT^j|$ on J_{n_i} , respectively. We have

$$(5) \quad \begin{aligned} \left| \log \frac{DT^t(x)}{DT^t(y)} \right| &\leq \sum_{i=0}^{t-1} |\log DT \circ T^i(x) - \log DT \circ T^i(y)| \\ &\leq \sum_{i=0}^{t-1} \frac{B_{n_i}}{M_{n_i}^{(1)}} |T^i(x) - T^i(y)| \leq \sum_{i=0}^{t-1} B_{n_i} |T^i(x) - T^i(y)|. \end{aligned}$$

Following [39], we write $|T^t(x) - T^t(y)| = |DT^{t-i}(z)| |T^i(x) - T^i(y)|$ for some $z \in J_{n_i}$ and use that the expansion $DT^{t-i}(z)$ decreases as n_i increases to conclude that the last sum here is maximized for $n_i = n + t - i$, i.e. $T^i(x), T^i(y) \in J_{n+t-i}$. Using equation (4) we obtain

$$\left| \log \frac{DT^t(x)}{DT^t(y)} \right| \lesssim \sum_{i=0}^{t-1} (n+t-i)^{\frac{1-\gamma}{\gamma}} (n+t-i)^{-\frac{\gamma+1}{\gamma}} = \sum_{i=0}^{t-1} (n+t-i)^{-2} \leq \min\{2, n^{-1}\},$$

where the last inequality follows from $\sum_{i=n+1}^{\infty} i^{-2} \leq \int_n^{+\infty} x^{-2} dx$.

Now, $T^t(x)$ has no preimage in $\bigcup_{i>n+t} J_i$, so the weakest expansion occurs when $x \in J_{n+t}$. The previous distortion estimate implies

$$DT^t(x) \gtrsim \frac{|J_n|}{|J_{n+t}|}.$$

Using equation (4) again, statement (a) easily follows.

(b) Adopting the same notation as in (a) and starting from (5), we fix $p \in (0, \frac{\gamma}{\gamma+1}]$ and write

$$\begin{aligned} \left| \log \frac{DT^t(x)}{DT^t(y)} \right| &\leq \sum_{i=0}^{t-1} B_{n_i} |T^i(x) - T^i(y)|^{1-p} \frac{|T^i(x) - T^i(y)|^p}{|T^t(x) - T^t(y)|^p} |T^t(x) - T^t(y)|^p \\ &\leq \sum_{i=0}^{t-1} \frac{B_{n_i} |J_{n_i}|^{1-p}}{(M_{n_i}^{(t-i)})^p} |T^t(x) - T^t(y)|^p. \end{aligned}$$

Using statement (a) and, as in the previous proof, that the worst case scenario in the upper bounds of equation (4) occurs for $n_i = n + t - i$, we obtain

$$\left| \log \frac{DT^t(x)}{DT^t(y)} \right| \lesssim n^{p\frac{\gamma+1}{\gamma}} \sum_{i=0}^{t-1} (n+t-i)^{-2} |T^t(x) - T^t(y)|^p \leq n^{p\frac{\gamma+1}{\gamma}-1} |T^t(x) - T^t(y)|^p,$$

and statement (b) follows from the assumption $p \leq \frac{\gamma}{\gamma+1}$. \square

Finally, on several occasions in the proofs, we shall require the following estimate.

Lemma 3.2. *For every pair $a, b > 1$, every pair n_0, n_1 such that $n_0 > n_1 + 1$, and for all t such that $n_0 + 1 \leq t + n_1$,*

$$\sum_{n=n_0+1}^{t+n_1} n^{-a} (t-n+n_0)^{-b} \lesssim t^{-\min\{a,b\}}.$$

Proof. According to the inequality $\sum_{n=n_0+1}^{t+n_1} f(n) \leq \int_{n_0}^{t+n_1+1} f(x) dx$ which holds for every $f \geq 0$, we estimate the sum via the following integral:

$$\begin{aligned} \int_{n_0}^{t+n_1+1} x^{-a}(t+n_0-x)^{-b} dx &= \int_{n_0}^{t/2} x^{-a}(t+n_0-x)^{-b} dx \\ &\quad + \int_{t/2}^{t+n_1+1} x^{-a}(t+n_0-x)^{-b} dx. \end{aligned}$$

In the first integral, the second factor in the integrand is at most $(n_0 + \frac{t}{2})^{-b}$, while the first factor integrates to something less than $\frac{n_0^{-(a-1)}}{a-1}$. In the second integral, the first factor is at most $(\frac{t}{2})^{-a}$, while the second factor integrates to something less than $\frac{(n_0-n_1-1)^{-(b-1)}}{b-1}$. The desired estimate immediately follows. \square

3.2. Estimating escape rates - proof of Theorem 2.1. Recall that H is a cylinder in $\mathcal{J}^{(\ell_H)}$ and $H \subseteq J_h$ for some $h \geq 0$. The main estimate of this section is the following lemma.

Lemma 3.3. $m(\dot{I}^t) \lesssim t^{-\frac{1}{\gamma}}$.

The proof of this lemma is based on the fact that an induced map related to T has exponential escape rate. To formulate this property, choose $n_S > h$, let $I_S = [a_{n_S}, 1) \supset H$ and consider the induced map $S = T^R : I_S \rightarrow I_S$, where R is the first return time to I_S .

Let $\dot{I}_S^t = \bigcap_{i=0}^t S^{-i}(I_S \setminus H)$ denote the set of points in I_S which do not enter H before time t under the action of S . The induced open system $S|_{\dot{I}_S^t}$ is uniformly expanding with countably many branches and admits a countable Markov partition which is formed by joining $\mathcal{J}^{(\ell_H)}$ with the partition into sets on which R is constant. The action of S on this partition satisfies the large images condition [16]; hence the following property holds.¹

Lemma 3.4 ([16]). *There exists $\sigma < 1$ such that $m(\dot{I}_S^t) \lesssim \sigma^t$.*

Proof of Lemma 3.3. We first assume there exists $d > 0$ such that $[\frac{1}{2}, \frac{1}{2}+d) \cap H = \emptyset$. The complementary case is much simpler and will be addressed at the end of the proof.

Without loss of generality, we can choose the index n_S that defines I_S sufficiently large so that $\frac{a_{n_S}}{2} < d$; viz. the open system makes full returns to the interval $J_S = [\frac{1}{2}, \frac{1}{2} + \frac{a_{n_S}}{2})$ before entering $I \setminus I_S = [0, a_{n_S})$. In order to obtain the estimate on $m(\dot{I}^t)$, we consider separately the sets $\dot{I}^t \cap I_S$ and $\dot{I}^t \cap (I \setminus I_S)$.

Case I (Estimate for points in $\dot{I}^t \cap I_S$). Consider the decomposition of $\dot{I}^t \cap I_S$ into subsets E_k^t of points having made k passes through $I \setminus I_S$ before time t . After each pass through $I \setminus I_S$, an orbit must spend at least $n_S + 1$ iterates within I_S before making its next pass. It results that the index k here is at most $\lfloor \frac{t-2}{n_S+2} \rfloor$.

¹The full results of [16] also require a “smallness” condition on the size of the hole. This condition is not needed here since we are not invoking any results regarding a spectral gap for the transfer operator associated with S , but just an exponential rate of escape, which does not require the hole to be small.

In order to estimate the measure of the sets E_k^t , we consider separately the cases $k > b \log t$ and $k \leq b \log t$, where $b = \frac{1}{(n_S+1)\gamma \log \sigma^{-1}}$ and σ is from Lemma 3.4.

For $k > b \log t$, we observe that every point in E_k^t must spend at least $(k-1)(n_S+1)$ iterates in I_S before hitting the hole. Hence we have $E_k^t \subset \overset{\circ}{I}_S^{(k-1)(n_S+1)}$, and using Lemma 3.4 and the definition of b , we get

$$\sum_{k > b \log t} m(E_k^t) \lesssim \sum_{k > b \log t} \sigma^{(k-1)(n_S+1)} \lesssim t^{-\frac{1}{\gamma}},$$

as desired.

For $k \leq b \log t$, we first note that the case $k = 0$ is easily estimated using $E_0^t \subset \overset{\circ}{I}_S^t$ and Lemma 3.4. From now on, we assume $k \in \{1, \dots, \lfloor b \log t \rfloor\}$ and observe as before that the subset $E_k^{t,+} \subset E_k^t$ of points whose orbit spends at least $b(n_S+1) \log t$ iterates in I_S up to time t is included in $\overset{\circ}{I}_S^{b(n_S+1) \log t}$. This inclusion implies $m\left(\bigcup_{k=1}^{\lfloor b \log t \rfloor} E_k^{t,+}\right) \lesssim t^{-\frac{1}{\gamma}}$.

It remains to consider the complementary subset $E_k^{t,-} = E_k^t \setminus E_k^{t,+}$ of points whose orbit up to t spends more than $t - b(n_S+1) \log t$ iterates in $I \setminus I_S$. Given $x \in E_k^{t,-}$ and $i \in \{1, \dots, k\}$, let $n_i > n_S$ be such that J_{n_i} is the element of \mathcal{J} where $T^j(x)$ begins its i th pass through $I \setminus I_S$. We must have

$$n_i - n_S > \frac{t - b(n_S+1) \log t}{k}$$

for at least one $i \in \{1, \dots, k\}$; otherwise we would have

$$\sum_{i=1}^k n_i - n_S \leq t - b(n_S+1) \log t,$$

which contradicts the definition of $E_k^{t,-}$. Accordingly, we have

$$(6) \quad m(E_k^{t,-}) \leq \sum_{i=1}^k m\left(x \in E_k^t : n_i - n_S > \frac{t - b(n_S+1) \log t}{k}\right).$$

The sets in this sum can be decomposed using symbolic dynamics. Given two integers $t_1 \leq t_2$ and a symbolic word $\theta_{t_1}^{t_2} \in \{L, R\}^{t_2-t_1+1}$, let $J_{\theta_{t_1}^{t_2}} = \bigcap_{\ell=t_1}^{t_2} T^{-\ell}(J_{\theta_\ell})$.

We have

$$(7) \quad \{x \in E_k^t : n_i = n\} = \bigcup_{j, \{\theta_\ell\}_{\ell=1}^{j-2}, \{\theta_\ell\}_{\ell=j+1}^t} J_{\theta_0^{j-1}} \cap T^{-j}(J_n) \cap J_{\theta_{j+1}^t},$$

where, by an abuse of notation,²

$$J_{\theta_0^{j-1}} := I_S \cap J_{\theta_1^{j-2}} \cap T^{-(j-1)}(J_S)$$

and where the union on $\{\theta_\ell\}_{\ell=0}^{j-2}$ (resp. $\{\theta_\ell\}_{\ell=j+1}^t$) is taken over all admissible words compatible with $i-1$ (resp. $k-i$) passing through $I \setminus I_S$ and avoiding H until at

²For this expression to be meaningful, we should decompose the sets I_S and $T^{-(j-1)}(J_S)$ into (standard) cylinder sets prior to define the intermediate cylinder $J_{\theta_1^{j-2}}$. This abuse of notation has no impact on the reasoning here. (The same comment applies to the set $T^{-j}(J_n)$ and to the decomposition in equation (7).)

least time t . The sets in (7) are pairwise disjoint; hence it suffices to estimate each quantity $m\left(J_{\theta_0^{j-1}} \cap T^{-j}(J_n) \cap J_{\theta_{j+1}^t}\right)$.

To proceed, notice first that the property $T^n(J_n) = J_0$ implies

$$J_{\theta_0^{j-1}} \cap T^{-j}(J_n) = J_{\theta_0^{j-1}} \cap T^{-j}(I \setminus I_S) \cap T^{-(j+n)}(J_0).$$

Moreover, the map T^{j+n} is one-to-one on each element of $T^{-(j+n)}(J_0)$. Assuming $j+n+1 \leq t$ and applying the bounded distortion estimate of the proof of Lemma 3.1, we obtain

$$\begin{aligned} & \frac{m\left(J_{\theta_0^{j-1}} \cap T^{-j}(J_n) \cap J_{\theta_{j+1}^t}\right)}{m\left(J_{\theta_0^{j-1}} \cap T^{-j}(I \setminus I_S) \cap T^{-(j+n)}(J_0)\right)} \\ & \approx \frac{m\left(T^{j+n}\left(J_{\theta_0^{j-1}} \cap T^{-j}(J_n) \cap J_{\theta_{j+1}^t}\right)\right)}{m\left(T^{j+n}\left(J_{\theta_0^{j-1}} \cap T^{-j}(I \setminus I_S) \cap T^{-(j+n)}(J_0)\right)\right)}. \end{aligned}$$

The second ratio here is equal to

$$\frac{m\left(J_0 \cap T^{j+n}(J_{\theta_{j+n+1}^t})\right)}{m(J_0)},$$

from where our first estimate follows:

$$m\left(J_{\theta_0^{j-1}} \cap T^{-j}(J_n) \cap J_{\theta_{j+1}^t}\right) \approx m\left(J_0 \cap T^{j+n}(J_{\theta_{j+n+1}^t})\right) m\left(J_{\theta_0^{j-1}} \cap T^{-j}(J_n)\right).$$

Proceeding similarly for the second factor above and using $T(J_S) = I \setminus I_S$, we get

$$\frac{m\left(J_{\theta_0^{j-1}} \cap T^{-j}(J_n)\right)}{m\left(J_{\theta_0^{j-1}}\right)} \approx \frac{m\left(T^j\left(J_{\theta_0^{j-1}} \cap T^{-j}(J_n)\right)\right)}{m\left(T^j\left(J_{\theta_0^{j-1}}\right)\right)} = \frac{m(J_n)}{a_{n_S}},$$

from which equation (4) implies

$$(8) \quad m\left(J_{\theta_0^{j-1}} \cap T^{-j}(J_n) \cap J_{\theta_{j+1}^t}\right) \approx n^{-\frac{\gamma+1}{\gamma}} m\left(J_0 \cap T^{j+n}(J_{\theta_{j+n+1}^t})\right) m\left(J_{\theta_0^{j-1}}\right).$$

In the case where $j+n+1 > t$ (which happens only when $i = k$), we use the inclusion

$$J_{\theta_0^{j-1}} \cap T^{-j}(J_n) \cap \bigcup_{\{\theta_\ell\}_{\ell=j+1}^t} J_{\theta_{j+1}^t} \subset J_{\theta_0^{j-1}} \cap T^{-j}(J_n)$$

to obtain, using the relation before equation (8),

$$m\left(J_{\theta_0^{j-1}} \cap T^{-j}(J_n) \cap J_{\theta_{j+1}^t}\right) \lesssim n^{-\frac{\gamma+1}{\gamma}} m\left(J_{\theta_0^{j-1}}\right).$$

Now use that imposing $i-1$ passes through $I \setminus I_S$ before time j implies at least $(i-1)(n_S+1)$ iterates in I_S before j to obtain the following relation:

$$m\left(\bigcup_{j, \{\theta_\ell\}_{\ell=1}^{j-2}} J_{\theta_0^{j-1}}\right) \subset m\left(I_S^{(i-1)(n_S+1)}\right) \lesssim \sigma^{(i-1)(n_S+1)}.$$

Similarly, for $j+n+1 \leq t$ (otherwise the consideration here is not needed), let q be the number of iterates that the orbits of points in $J_0 \cap T^{j+n}(J_{\theta_{j+n+1}^t})$ spend in I_S . Each pass in $I \setminus I_S$ from $i+1$ through $k-1$ must be followed by at least n_S+1 iterates in I_S ; hence $q \geq (k-i-1)(n_S+1)$ (also $q \leq t - (j+n) + 1 - (k-i-1)$

where the maximum is obtained when each pass in $I \setminus I_S$ consists of a single iterate) and then by Lemma 3.4,

$$m \left(\bigcup_{j, \{\theta_\ell\}_{\ell=j+1}^t} J_0 \cap T^{j+n}(J_{\theta_{j+n+1}^t}) \right) \leq \sum_{q \geq (k-i-1)(n_S+1)} m(I_S^q) \lesssim \sigma^{(k-i)(n_S+1)}.$$

(Notice that this estimate holds even in the case $i = k$.) Putting these estimates together with (7) and (8), we have obtained

$$m(E_k^t : n_i = n) \lesssim \sigma^{(k-1)(n_S+1)} n^{-\frac{\gamma+1}{\gamma}}.$$

Using the inequality

$$\sum_{n > n_S + \frac{t - b(n_S+1) \log t}{k}} n^{-\frac{\gamma+1}{\gamma}} \leq \left(n_S + \frac{t - b(n_S+1) \log t}{k} \right)^{-\frac{1}{\gamma}}$$

it follows from (6) that

$$m(E_k^{t,-}) \lesssim \sigma^{(k-1)(n_S+1)} k^{\frac{\gamma+1}{\gamma}} (kn_S + t - b(n_S+1) \log t)^{-\frac{1}{\gamma}}.$$

It remains to sum over k . We finally have

$$m \left(\bigcup_{k=1}^{\lfloor b \log t \rfloor} E_k^{t,-} \right) \lesssim (n_S + t - b(n_S+1) \log t)^{-\frac{1}{\gamma}} \sum_{k=1}^{\lfloor b \log t \rfloor} \sigma^{(k-1)(n_S+1)} k^{\frac{\gamma+1}{\gamma}} \lesssim t^{-\frac{1}{\gamma}},$$

as desired, where we used

$$\sum_{k=1}^{b \log t} \sigma^{(k-1)(n_S+1)} k^{\frac{\gamma+1}{\gamma}} < +\infty \quad \text{and} \quad (n_S + t - b(n_S+1) \log t)^{-1} \lesssim t^{-1}.$$

Case II (Estimate for points in $\hat{I}^t \cap (0, a_{n_S})$). Recall that $(0, a_{n_S}) = \bigcup_{n > n_S} J_n$ and by definition of the J_n , we have $\hat{I}^t \supset \bigcup_{n > t+h} J_n$ so that using equation (4) yields

$$m \left(\hat{I}^t \cap \bigcup_{n > t+h} J_n \right) \lesssim (t+h)^{-\frac{1}{\gamma}} \leq t^{-\frac{1}{\gamma}}.$$

It remains to estimate $m \left(\hat{I}^t \cap \bigcup_{n=n_S+1}^{t+h} J_n \right)$. For every $n > n_S$, we have $T^{n-n_S}(\hat{I}^t \cap J_n) = \hat{I}^{t-n+n_S} \cap J_{n_S}$. Using bounded distortion again, we get

$$\frac{m(\hat{I}^t \cap J_n)}{m(J_n)} \approx \frac{m(\hat{I}^{t-n+n_S} \cap J_{n_S})}{m(J_{n_S})},$$

which, together with the inclusion $J_{n_S} \subset I_S$ and the conclusion in Case I, implies

$$(9) \quad m \left(\hat{I}^t \cap \bigcup_{n=n_S+1}^{t+h} J_n \right) \lesssim \sum_{n=n_S+1}^{t+h} n^{-\frac{\gamma+1}{\gamma}} (t-n+n_S)^{-\frac{1}{\gamma}}.$$

Lemma 3.2 implies that the sum up to $t+h-1$ is bounded above by $\lesssim t^{-\frac{1}{\gamma}}$. For the last term, we have $(t+h)^{-\frac{\gamma+1}{\gamma}} (n_S-h)^{-\frac{1}{\gamma}} \lesssim t^{-\frac{1}{\gamma}}$.

It results that $m\left(\dot{I}^t \cap \bigcup_{n=n_S+1}^{t+h} J_n\right) \lesssim t^{-\frac{1}{\gamma}}$, and this concludes the proof of Lemma 3.3 in the case where H is disjoint from $[\frac{1}{2}, \frac{1}{2} + d)$ for some $d > 0$.

The case in which H contains $[\frac{1}{2}, \frac{1}{2} + d)$ for some $d > 0$ is much simpler since points starting in $[2d, 1)$ never enter $[0, 2d)$ before escaping. Thus the estimates of Case I with $k = 0$ together with Case II imply that the upper bound on $m(\dot{I}^t)$ in this case is the same. \square

With Lemma 3.3 established, we are ready to prove Theorem 2.1.

Proof of Theorem 2.1. Given a density f as in the theorem, let $C_0 = \inf_{x \in (0, x_0)} x^\alpha f(x) > 0$ and $n_0 = \min\{n : a_n \leq x_0\}$. We have $\dot{I}^t \supset \bigcup_{n>t+h} J_n$ for all $t \geq 0$; hence for t sufficiently large so that $t + h \geq n_0$, equation (4) implies

$$(10) \quad \mu_f(\dot{I}^t) = \int_{\dot{I}^t} f \, dm \geq C_0 \int_0^{a_{t+h}} x^{-\alpha} \, dx \gtrsim (h+t)^{-\frac{1-\alpha}{\gamma}},$$

from where the lower bound immediately follows.

For the upper bound, we split $[0, 1)$ into 3 intervals: $[0, a_{h+t}) \cup [a_{h+t}, a_{n_S}) \cup I_S$ and estimate the intersection of \dot{I}^t with each of these separately.³

On $[0, a_{h+t})$, we estimate

$$\int_{\dot{I}^t \cap [0, a_{h+t})} f \, dm \lesssim \int_0^{a_{h+t}} x^{-\alpha} \, dx \lesssim (h+t)^{-\frac{1-\alpha}{\gamma}},$$

while on I_S , we have, using that f is bounded on this set and Lemma 3.3,

$$\int_{\dot{I}^t \cap I_S} f \, dm \lesssim m(\dot{I}^t) \lesssim t^{-\frac{1}{\gamma}}.$$

On $[a_{h+t}, a_{n_S}) = \bigcup_{n=n_S+1}^{t+h} J_n$, we proceed as in Case II of the previous proof:

$$\begin{aligned} \int_{\dot{I}^t \cap [a_{h+t}, a_{n_S})} f \, dm &= \sum_{n=n_S+1}^{t+h} \int_{\dot{I}^t \cap J_n} f \, dm \lesssim \sum_{n=n_S+1}^{t+h} a_n^{-\alpha} m(\dot{I}^t \cap J_n) \\ &\lesssim \sum_{n=n_S+1}^{t+h} n^{-\frac{\gamma+1-\alpha}{\gamma}} (t-n+n_S)^{-\frac{1}{\gamma}}, \end{aligned}$$

where we have used (9). As before, the last sum (except its last term) is estimated using Lemma 3.2 to give

$$\sum_{n=n_S+1}^{t+h} n^{-\frac{\gamma+1-\alpha}{\gamma}} (t-n+n_S)^{-\frac{1}{\gamma}} \lesssim t^{-\frac{1-\alpha}{\gamma}}.$$

\square

³For $\alpha = 0$, the upper bound also directly follows from the fact that f is uniformly bounded on I together with Lemma 3.3, namely

$$\mu_f(\dot{I}^t) \lesssim m(\dot{I}^t) \lesssim t^{-\frac{1}{\gamma}}.$$

3.3. Properties of the function spaces \mathcal{F}_p . The definition of the quantity H_J^p before Theorem 2.2 implies the following simple facts about the set \mathcal{F}_p ($p \in \mathbb{R}^+$), whose proof we leave to the reader.

Lemma 3.5. (1) $\|\cdot\|_p$ is scale invariant, i.e. $\|Cf\|_p = \|f\|_p$ for any $C > 0$.
 (2) For any $J \in \mathcal{J}^{(\ell_H)}$, E a subinterval of J and $f \in \mathcal{F}_p$, we have

$$\sup_{x \in E} f(x) \leq e^{H_J^p(f)|E|^p} \inf_{x \in E} f(x) \leq e^{H_J^p(f)|E|^p} |E|^{-1} \int_E f \, dm.$$

(3) If $q > p$, then $\mathcal{F}_q \subset \mathcal{F}_p$ (and $\mathcal{F}_q^0 \subset \mathcal{F}_p^0$).

Now, we equip the set of measures with the topology of weak convergence, consider the ball $\mathcal{B}_p = \{\mu_f : f \in \mathcal{F}_p, \|f\|_p \leq 1\}$ and notice that this ball is not closed. Indeed, given $\ell \in \mathbb{N}$, let the density f_ℓ be defined by

$$f_\ell(x) = \begin{cases} a_\ell^{-1} & \text{if } x \in \bigcup_{n \geq \ell+1} J_n, \\ 0 & \text{elsewhere.} \end{cases}$$

Then we have $f_\ell \geq 0$, $\int f_\ell \, dm = 1$ and $\|f_\ell\|_p = 0$ so that $\mu_{f_\ell} \in \mathcal{B}_p$. However, we clearly have

$$\lim_{\ell \rightarrow \infty} \mu_{f_\ell} = \delta_0 \notin \mathcal{B}_p.$$

The Dirac measure at 0 turns out to be the only possible singular component to where sequences in \mathcal{B}_p can accumulate.

Lemma 3.6. The set $\{(1-s)\mu_f + s\delta_0 : s \in [0, 1], f \in \mathcal{B}_p\}$ is compact.

Proof. Let $\{\mu_{f_\ell}\}_{\ell \in \mathbb{N}} \subset \mathcal{B}_p$ be an arbitrary sequence. Since $\int f_\ell \, dm = 1$, there exists a subsequence $\{\mu_{f_{\ell_k}}\}$ which converges weakly to a probability measure μ_∞ on I . Now fix $J \in \mathcal{J}^{(\ell_H)}$. By Lemma 3.5, the sequence of densities $\{f_{\ell_k}\}$ is a bounded, equicontinuous family on J . By the Arzelà-Ascoli theorem, there exists a subsequence that converges uniformly to a function $f_J^{(\infty)}$ on J .⁴ Note that $H_J^p(f_J^{(\infty)}) \leq 1$.

Diagonalizing, we obtain a subsequence $\{f_{\ell_{k_j}}\}$ converging to $f_J^{(\infty)}$ on each $J \in \mathcal{J}^{(\ell_H)}$. Letting $f^{(\infty)} = \sum_J f_J^{(\infty)}$, we have $f^{(\infty)} \geq 0$, $\|f^{(\infty)}\|_p \leq 1$, and by Fatou's lemma, $\int f^{(\infty)} \, dm \leq 1$.

Let $s = 1 - \int f^{(\infty)} \, dm$. If $s < 1$, let $f_\infty = (1-s)^{-1}f^{(\infty)}$. By the above observations, we have $\mu_{f_\infty} \in \mathcal{B}_p$. Since $\{0\}$ is the only accumulation point of the sequence of sets $\{J\}_{J \in \mathcal{J}^{(\ell_H)}}$, we must have $\mu_\infty = (1-s)\mu_{f_\infty} + s\delta_0$, as required. \square

For the next statement, we need to introduce the (nonlinear) normalized transfer operator and its iterates:

$$(11) \quad \mathring{\mathcal{L}}_1^t f := \frac{\mathring{\mathcal{L}}^t f}{|\mathring{\mathcal{L}}^t f|_1}, \quad \forall t \geq 1.$$

Recall that $\mathcal{F}_p^0 = \{f \in \mathcal{F}_p : \exists x_0 \in (0, 1) \text{ and } \alpha \in [0, 1) \text{ such that (1) holds}\}$.

⁴To be precise, note that $\|f_\ell\|_p \leq 1$ implies that f_ℓ is uniformly continuous on J so that f can be extended to a continuous function \tilde{f}_ℓ on the closure \bar{J} and $\|\tilde{f}_\ell\|_p \leq 1$ still holds.

Proposition 3.7. *Let $p \in (0, \frac{\gamma}{\gamma+1}]$. We have $\mathring{\mathcal{L}}_1(\mathcal{F}_p^0) \subset \mathcal{F}_p^0$. In addition, there exist two constants $C_1, C_2 \geq 0$ such that for every $f \in \mathcal{F}_p^0$,*

$$\|\mathring{\mathcal{L}}_1^t f\|_p \leq C_1 \|f\|_p + C_2 \text{ for all } t \geq 1.$$

Proof. Every $J \subset \mathcal{J}^{(\ell_H)}$ has at most two pre-images under \mathring{T} and each pre-image is included in some element of $\mathcal{J}^{(\ell_H)}$. This implies that $\mathring{\mathcal{L}}(C^0(\mathcal{J}^{(\ell_H)})) \subset C^0(\mathcal{J}^{(\ell_H)})$. Also we obviously have

$$\mathring{\mathcal{L}}f \geq 0 \quad \text{and} \quad |\mathring{\mathcal{L}}f|_1 \leq |f|_1,$$

for every $f \geq 0$.

Now, fix $f \in \mathcal{F}_p^0$ and assume $C_0 \geq 1$ and $\alpha \in [0, 1)$ are such that $C_0^{-1} \leq x^\alpha f(x) \leq C_0$ for $x \in (0, a_{n_0})$, and $f \leq C_0$ on $I \setminus [0, a_{n_0})$. Without loss of generality, we may assume $n_0 > h + 1$. Now letting \mathring{T}_L and \mathring{T}_R denote the left and right branches of \mathring{T} respectively, we have

$$(12) \quad \mathring{\mathcal{L}}f(x) = \frac{f(\mathring{T}_L^{-1}(x))}{DT(\mathring{T}_L^{-1}(x))} + \frac{f(\mathring{T}_R^{-1}(x))}{DT(\mathring{T}_R^{-1}(x))}.$$

If $y = \mathring{T}_L^{-1}(x) \in (0, \frac{1}{2})$, then it follows from the definition of T that $\frac{x}{2} \leq y \leq x$. Thus if $x \in (0, a_{n_0-1})$, then by assumption on f ,

$$\begin{aligned} C_0^{-1}(\mathring{T}_L^{-1}(x))^{-\alpha} &\leq f(\mathring{T}_L^{-1}(x)) \leq C_0(\mathring{T}_L^{-1}(x))^{-\alpha} \\ \implies C_0^{-1}x^{-\alpha} &\leq f(\mathring{T}_L^{-1}(x)) \leq 2C_0x^{-\alpha}. \end{aligned}$$

Combining this estimate with (12) and using the fact that $1 \leq DT_L \leq 3$ and $DT_R = 2$, we have

$$\frac{1}{3}C_0^{-1}x^{-\alpha} \leq \mathring{\mathcal{L}}f(x) \leq 2C_0x^{-\alpha} + \frac{C_0}{2} \leq 3C_0x^{-\alpha},$$

for all $x \in (0, a_{n_0-1})$, which is the required polynomial bound on the behavior of $\mathring{\mathcal{L}}f$ near 0. For $x \in [a_{n_0-1}, 1)$, we use the fact that f is bounded by C_0 at both pre-images of x so that $\mathring{\mathcal{L}}f(x) \leq 2C_0$.

Moreover, the lower bound on $\mathring{\mathcal{L}}f$ given above implies that $|\mathring{\mathcal{L}}f|_1 > 0$ so that $\mathring{\mathcal{L}}_1 f$ is well-defined. Anticipating the proof below that $\|\mathring{\mathcal{L}}f\|_p < +\infty$ for every f with $\|f\|_p < +\infty$, we obtain $\mathring{\mathcal{L}}_1(\mathcal{F}_p^0) \subset \mathcal{F}_p^0$.

In order to check the estimate on $\|\mathring{\mathcal{L}}_1^t f\|_p$, it suffices to prove the inequality for $\|\mathring{\mathcal{L}}^t f\|_p$ due to the scale invariance property from Lemma 3.5(1).

Let $f \in \mathcal{F}_p^0$. Fix $n \geq 0$, $J \in \mathcal{J}^{(\ell_H)}$, $J \subset J_n$ and $x, y \in J$. Let also $t \geq 1$ and $\{x_i\}$ (resp. $\{y_i\}$) be an enumeration of the pre-images $\mathring{T}^{-t}(x)$ (resp. $\mathring{T}^{-t}(y)$) such that each pair x_i, y_i lies in the same branch of \mathring{T}^{-t} . Then

$$(13) \quad \log \mathring{\mathcal{L}}^t f(x) - \log \mathring{\mathcal{L}}^t f(y) = \log \frac{\sum_i f(x_i)/DT^t(x_i)}{\sum_i f(y_i)/DT^t(y_i)} \leq \max_i \log \frac{f(x_i)}{f(y_i)} + \log \frac{DT^t(y_i)}{DT^t(x_i)},$$

where we have used the fact that $\frac{\sum_i b_i}{\sum_i c_i} \leq \max_i \frac{b_i}{c_i}$ for two series of positive terms. The first term on the right hand side of (13) is estimated by

$$\log \frac{f(x_i)}{f(y_i)} \leq \|f\|_p |x_i - y_i|^p,$$

while the second term on the right side of (13) is estimated by

$$\log \frac{DT^t(x_i)}{DT^t(y_i)} \lesssim |T^t(x_i) - T^t(y_i)|^p = |x - y|^p,$$

according to Lemma 3.1(b). Putting these two estimates together and using Lemma 3.1(a), we obtain that there exist $C_1, C_2 > 0$ such that

$$\begin{aligned} \frac{|\log \mathring{\mathcal{L}}^t f(x) - \log \mathring{\mathcal{L}}^t f(y)|}{|x - y|^p} &\leq \|f\|_p \max_i \frac{|x_i - y_i|^p}{|x - y|^p} + C_2 \\ &\leq \|f\|_p C_1 \left(\frac{n+1}{n+t} \right)^{p \frac{\gamma+1}{\gamma}} + C_2 \\ &\leq C_1 \|f\|_p + C_2, \end{aligned}$$

as desired. \square

The last statement of this section provides examples of unbounded densities that belong to the sets \mathcal{F}_p^0 .

Lemma 3.8. *Let $p \in (0, \frac{\gamma}{\gamma+1}]$. Every function $\frac{f}{|f|_1}$ where $f(x) = x^{-\alpha}$ for all $x \in (0, 1)$, $\alpha \in [0, 1)$ belongs to \mathcal{F}_p^0 . The same is true for the density f_{SRB} associated with the SRB measure.*

Proof. Letting $f(x) = x^{-\alpha}$ for all $x \in (0, 1)$, we clearly have $f \in C^0(\mathcal{J}^{(\ell_H)})$, $f \geq 0$ and $|f|_1 < +\infty$ (and (1) holds). Therefore, we only need to check that $\|f\|_p < +\infty$.

Fix $J \in \mathcal{J}^{(\ell_H)}$, $J \subset J_n$ for some $n \geq 0$ and let $x < y \in J$. Then

$$\frac{\log f(x) - \log f(y)}{|x - y|^p} = \alpha \frac{\log(y/x)}{|x - y|^p} \leq \frac{\alpha}{x} |x - y|^{1-p}.$$

Since $x \in J_n$, we have $\frac{1}{x} \lesssim n^{-\frac{1}{\gamma}}$ and $|x - y| \lesssim n^{-\frac{\gamma+1}{\gamma}}$ by equation (4). Thus

$$H_J^p(f) \lesssim n^{\frac{1-(\gamma+1)(1-p)}{\gamma}},$$

and the exponent of n is non-positive when $p \leq \gamma/(\gamma+1)$. Taking the supremum over n , we have $\|f\|_p < \infty$ as required.

As for f_{SRB} , observe that the sequence $\{\mathcal{L}^t 1\}_{t \in \mathbb{N}}$ (transfer operator for the system without the hole) converges to the density f_{SRB} . On the other hand, the proof of Proposition 3.7 can be repeated *mutatis mutandis* to conclude $\mathcal{L}(\mathcal{F}_p^0) \subset \mathcal{F}_p^0$ and $\sup_{t \in \mathbb{N}} \|\mathcal{L}^t f\|_p < +\infty$ for all $f \in \mathcal{F}_p^0$. Since $1 \in \mathcal{F}_p^0$, it follows from Lemma 3.6 that $f_{\text{SRB}} \in \mathcal{F}_p$. However, $f_{\text{SRB}}(x) \approx x^{-\gamma}$; hence we must have $f_{\text{SRB}} \in \mathcal{F}_p^0$ as desired. \square

3.4. Proof of Theorem 2.2. The proof relies on the following strengthening of the volume estimate in Lemma 3.3 on the set of points that enters H precisely at time t .

Lemma 3.9. $m(\mathring{I}^{t-1} \setminus \mathring{I}^t) \lesssim t^{-\frac{\gamma+1}{\gamma}} \log t, \forall t \geq 2$.

The proof is given in Section 3.5 below.

Remark 3.10. We believe one should be able to eliminate the factor $\log t$ and thus obtain Lemma 3.3 via the identity $m(\mathring{I}^t) = \sum_{i=t}^{\infty} m(\mathring{I}^i \setminus \mathring{I}^{i+1})$. Although we are able to prove this upper bound in a special case (see Lemma 3.11), our current techniques do not provide this estimate in the general case, so we will use the weaker version stated above.

Let $f \in \mathcal{F}_p^0$ for some $p > 0$. Using Lemma 3.5(3), we may assume without loss of generality that $p \in (0, \frac{\gamma}{\gamma+1}]$. Assume for now that the exponent α of f from (1) is positive. We are going to derive a bound analogous to Lemma 3.9 for μ_f .

Let n_0 be the smallest n such that $f(x) \approx x^{-\alpha}$ on $(0, a_n)$. As in the proof of Theorem 2.1, we split $[0, 1)$ into 3 intervals, namely $[0, a_t)$, $[a_t, a_{n_0})$ and $[a_{n_0}, 1)$.

We have $(\dot{I}^{t-1} \setminus \dot{I}^t) \cap [0, a_t) \subset J_{t+h}$. Thus

$$\int_{(\dot{I}^{t-1} \setminus \dot{I}^t) \cap [0, a_t)} f \, dm \lesssim \int_{J_{t+h}} x^{-\alpha} \, dx \lesssim (h+t)^{-\frac{\gamma+1}{\gamma} + \frac{\alpha}{\gamma}}.$$

Moreover, since f is bounded on $[a_{n_0}, 1)$, we also have

$$\int_{(\dot{I}^{t-1} \setminus \dot{I}^t) \cap [a_{n_0}, 1)} f \, dm \lesssim m(\dot{I}^{t-1} \setminus \dot{I}^t) \lesssim t^{-\frac{\gamma+1}{\gamma}} \log t.$$

It remains to estimate the μ_f measure of $\bigcup_{n=n_0+1}^t J_n \cap (\dot{I}^{t-1} \setminus \dot{I}^t)$. Since without loss of generality, we may take $n_0 > h$, we have for each $n > n_0$, $T^{n-n_0}(J_n \cap (\dot{I}^{t-1} \setminus \dot{I}^t)) = J_{n_0} \cap (\dot{I}^{t-1-n+n_0} \setminus \dot{I}^{t-n+n_0})$. Using bounded distortion, we obtain

$$\frac{m(J_n \cap (\dot{I}^{t-1} \setminus \dot{I}^t))}{m(J_n)} \approx \frac{m(J_{n_0} \cap (\dot{I}^{t-1-n+n_0} \setminus \dot{I}^{t-n+n_0}))}{m(J_{n_0})},$$

which implies $m(J_n \cap (\dot{I}^{t-1} \setminus \dot{I}^t)) \lesssim n^{-\frac{\gamma+1}{\gamma}} (t-n+n_0)^{-\frac{\gamma+1}{\gamma}} \log(t-n+n_0)$. Now

$$\begin{aligned} \int_{(\dot{I}^{t-1} \setminus \dot{I}^t) \cap [a_t, a_{n_0})} f \, dm &= \sum_{n=n_0+1}^t \int_{(\dot{I}^{t-1} \setminus \dot{I}^t) \cap J_n} f \, dm \lesssim \sum_{n=n_0+1}^t a_n^{-\alpha} m(J_n \cap (\dot{I}^{t-1} \setminus \dot{I}^t)) \\ &\lesssim \sum_{n=n_0+1}^t n^{-\frac{\gamma+1}{\gamma} + \frac{\alpha}{\gamma}} (t-n+n_0)^{-\frac{\gamma+1}{\gamma}} \log(t-n+n_0). \end{aligned}$$

Since $\alpha > 0$, we may dominate $\log(t-n+n_0)$ by $C(t-n+n_0)^{\alpha/\gamma}$ for some $C > 0$. Now using Lemma 3.2, we finally conclude the existence of a constant \bar{C} such that

$$(14) \quad \mu_f(\dot{I}^{t-1} \setminus \dot{I}^t) \leq \bar{C} t^{-\frac{\gamma+1}{\gamma} + \frac{\alpha}{\gamma}}.$$

Notice that using the relation $\mu_f(\dot{I}^t) = \sum_{i=t}^{\infty} \mu_f(\dot{I}^i \setminus \dot{I}^{i+1})$, this inequality implies the estimate $\mu_f(\dot{I}^t) \lesssim t^{-\frac{1-\alpha}{\gamma}}$ for $\alpha > 0$ without using Lemma 3.3. However, for $\alpha = 0$, a similar reasoning yields $\mu_f(\dot{I}^{t-1} \setminus \dot{I}^t) \lesssim t^{-\frac{\gamma+1}{\gamma}} \log t$, from which the conclusion of Theorem 2.1 cannot be deduced, hence the necessity of Lemma 3.3.

In any case, together with the estimate $\mu_f(\dot{I}^t) \geq C_f t^{-\frac{1-\alpha}{\gamma}}$ from Theorem 2.1, (14) yields

$$\begin{aligned} 1 &\geq \frac{\mu_f(\dot{I}^{t+1})}{\mu_f(\dot{I}^t)} = \frac{\mu_f(\dot{I}^t) - \mu_f(\dot{I}^t \setminus \dot{I}^{t+1})}{\mu_f(\dot{I}^t)} \\ &\geq 1 - \frac{\bar{C} t^{-\frac{\gamma+1}{\gamma} + \frac{\alpha}{\gamma}}}{C_f t^{-\frac{1-\alpha}{\gamma}}} = 1 - \frac{\bar{C}}{C_f} t^{-1} \rightarrow 1 \text{ as } t \rightarrow +\infty, \end{aligned}$$

and a similar conclusion holds for $\alpha = 0$ by Lemma 3.9. Consequently, we have proved the following limit for every $k \geq 1$ (and $\alpha \in [0, 1)$):

$$(15) \quad \lim_{t \rightarrow +\infty} \frac{\mu_f(\hat{I}^{t+k})}{\mu_f(\hat{I}^t)} = \prod_{i=0}^{k-1} \left(\lim_{t \rightarrow +\infty} \frac{\mu_f(\hat{I}^{t+i+1})}{\mu_f(\hat{I}^{t+i})} \right) = 1.$$

Now, since $p \in (0, \frac{\gamma}{\gamma+1}]$, we can apply Proposition 3.7 to conclude that the sequence $\left\{ \frac{\hat{T}_*^t \mu_f}{\mu_f(\hat{I}^t)} \right\}_{t \in \mathbb{N}}$ is composed of absolutely continuous probability measures with densities in \mathcal{F}_p^0 . By Lemma 3.6, any of its limit points must have the form $\mu_\infty = (1 - s_\infty)\mu_{f_\infty} + s_\infty\delta_0$ for some $f_\infty \in \mathcal{F}_p$ and $s_\infty \in [0, 1]$. We want to prove that $s_\infty = 1$ for any limit point.

Let $J \in \mathcal{J}^{(\ell_H)}$, let $g_t := \hat{\mathcal{L}}_1^t f$ and consider a converging subsequence $\{g_{t_j}\}_{j \in \mathbb{N}}$ with limit point $(1 - s_\infty)f_\infty$. (Recall that $\hat{\mathcal{L}}_1^t$ is the normalized transfer operator; see equation (11) above.) Since $f_\infty \in \mathcal{F}_p$, the convergence $g_{t_j}|_J \rightarrow (1 - s_\infty)f_\infty|_J$ holds in the uniform topology of functions defined on this interval. In particular, its integrals against any bounded measurable function converge as well on each $J \in \mathcal{J}^{(\ell_H)}$.

Fixing $k \geq 1$, note that the set $\bigcup_{i=0}^k T^{-i}(H)$ is bounded away from 0 and thus intersects only finitely many elements of $\mathcal{J}^{(\ell_H)}$. Thus the sequence $\{g_{t_j}\}_{j \in \mathbb{N}}$ converges uniformly on this set as well. Now, we have

$$\frac{\mu_f(\hat{I}^{k+t_j})}{\mu_f(\hat{I}^{t_j})} = \frac{|\hat{\mathcal{L}}_1^{k+t_j} f|_1}{|\hat{\mathcal{L}}_1^{t_j} f|_1} = \int_I \hat{\mathcal{L}}_1^k g_{t_j} dm = \int_{\hat{I}^k} g_{t_j} dm = 1 - \int_{\bigcup_{i=0}^k T^{-i}(H)} g_{t_j} dm,$$

using the fact that $\int_I g_{t_j} dm = 1$.

Since the limit of the above expression is 1 by (15) and the convergence of g_{t_j} to $(1 - s_\infty)f_\infty$ is uniform on each J , we must have $(1 - s_\infty)f_\infty \equiv 0$ on $\bigcup_{i=0}^k T^{-i}(H)$. Since $f_\infty \in \mathcal{F}_p$ is log-Hölder continuous on each $J \in \mathcal{J}^{(\ell_H)}$, we conclude that $(1 - s_\infty)f_\infty \equiv 0$ on any J such that $J \cap \left(\bigcup_{i=0}^k T^{-i}(H) \right) \neq \emptyset$. Since this holds for all k , the transitivity of T implies that we must have $(1 - s_\infty)f_\infty \equiv 0$ on all $J \in \mathcal{J}^{(\ell_H)}$, viz. $s_\infty = 1$. Since the subsequence is arbitrary, it follows that $s_\infty = 1$ for any limit point as desired. Theorem 2.2 is proved.

3.5. Proof of Lemma 3.9. We first prove the following auxiliary result. Given $t \geq 1$ and $0 \leq s < t$, let

$$E^t = \{x \in I : \min\{k \geq 0 : T^k(x) \in J_h\} = t\}$$

and

$$E_s^{s+t} = \{x \in T^{-s}(J_h) : \min\{k > 0 : T^{s+k}(x) \in J_h\} = t\}.$$

Lemma 3.11. $m(E^t) \lesssim t^{-\frac{\gamma+1}{\gamma}}$ and $m(E_s^{s+t}) \lesssim t^{-\frac{\gamma+1}{\gamma}}$.

Proof. The set E^t consists of a collection of intervals in $T^{-t}(I)$. Using symbolic dynamics as in equation (7) above, we label these intervals according to their itinerary

up to the first time k ($0 \leq k \leq t$) when they enter $[0, a_{h-1})$, viz.

$$(16) \quad E^t = \bigcup_{k=0}^t \bigcup_{\{\theta_\ell\}_{\ell=0}^{k-1}} J_{\theta_0^{k-1}} \cap T^{-k}(J_{h+t-k}),$$

where the words $\{\theta_\ell\}_{\ell=0}^{k-1}$ are such that $J_{\theta_\ell^{k-1}} \subset [a_{h-1}, 1)$ for all $0 \leq \ell \leq k-1$. Notice that the term $k=0$ actually reduces to J_{h+t} .

Using bounded distortion for the map T^k , we get

$$\frac{m\left(J_{\theta_0^{k-1}} \cap T^{-k}(J_{h+t-k})\right)}{m\left(J_{\theta_0^{k-1}} \cap T^{-k}([0, a_{h-1}))\right)} \approx \frac{m(J_{h+t-k})}{m([0, a_{h-1}))} \approx (h+t-k)^{-\frac{\gamma+1}{\gamma}}.$$

Moreover, recall the induced map S and the associated sets \mathring{I}_S^t defined at the beginning of Section 3.2, which we now consider for $n_S = h-1$. For $k \geq 1$, we actually have

$$\bigcup_{\{\theta_\ell\}_{\ell=0}^{k-1}} J_{\theta_0^{k-1}} \cap T^{-k}([0, a_{h-1})) \subset \mathring{I}_S^{k-1};$$

hence Lemma 3.4 implies the existence of $\sigma_h \in (0, 1)$ such that

$$m\left(\bigcup_{\{\theta_\ell\}_{\ell=0}^{k-1}} J_{\theta_0^{k-1}} \cap T^{-k}([0, a_{h-1}))\right) \lesssim \sigma_h^k.$$

For $k=0$, we obviously have $m([0, a_{h-1})) \leq 1$. Grouping all terms together, we finally get

$$m(E^t) \lesssim \sum_{k=0}^t (h+t-k)^{-\frac{\gamma+1}{\gamma}} \sigma_h^k.$$

We now show that this last sum is $\lesssim t^{-\beta}$ where $\beta = \frac{\gamma+1}{\gamma}$. The terms $k=0$, $k=1$ and $k=t$ satisfy this estimate. Moreover, the remaining sum from $k=2$ to $t-1$ is not larger than the following integral, which we compute by integration by parts:

$$\begin{aligned} \int_1^t (h+t-x)^{-\beta} \sigma_h^x dx &= \frac{\sigma_h (h+t-1)^{-\beta}}{\log \sigma_h^{-1}} - \frac{h^{-\beta} \sigma_h^t}{\log \sigma_h^{-1}} \\ &+ \int_1^{\frac{t}{2}} \beta \frac{(h+t-x)^{-\beta-1} \sigma_h^x}{\log \sigma_h^{-1}} dx + \int_{\frac{t}{2}}^t \beta \frac{(h+t-x)^{-\beta-1} \sigma_h^x}{\log \sigma_h^{-1}} dx. \end{aligned}$$

The first two terms in the right hand side here are $\lesssim t^{-\beta}$. Moreover, on the first integral, the integrand is at most $\frac{\sigma_h}{\log \sigma_h^{-1}} (h + \frac{t}{2})^{-\beta-1}$; hence the desired estimate follows. On the second integral, the integrand is at most $\frac{h^{-\beta-1}}{\log \sigma_h^{-1}} \sigma_h^t$, and then the integral is also dominated as $\lesssim t^{-\beta}$. The proof of the estimate on $m(E^t)$ is complete.

For $m(E_s^{s+t})$, the argument is similar. We decompose the set as follows:

$$E_s^{s+t} = \bigcup_{k=h+1}^t \bigcup_{\{\theta_\ell\}_{\ell=0}^{s+k-1}} J_{\theta_0^{s+k-1}} \cap T^{-(k+s)}(J_{h+t-k}),$$

where now the words $\{\theta_\ell\}_{\ell=0}^{s+k-1}$ are such that $J_{\theta_\ell^{s+k-1}} \subset J_h$ and $J_{\theta_\ell^{s+k-1}} \subset [a_{h-1}, 1)$ for all $s < \ell \leq s+k-1$. Using the same bounded distortion estimates as above, we get

$$m(E_s^{s+t}) \lesssim \sum_{k=h+1}^t (h+t-k)^{-\frac{\gamma+1}{\gamma}} \sigma_h^k,$$

from which the conclusion immediately follows. \square

We will prove Lemma 3.9 by keeping track of passes through J_h and concatenating estimates of the form in Lemma 3.11. Unfortunately, in estimating the contribution of points that make k passes through J_h before entering H , we obtain a factor C_0^k , which is potentially disastrous since k can increase with t . In order to overcome the effect of this constant, we slightly weaken the rate of decay as in the statement of the lemma.

Proof of Lemma 3.9. Let $C_0 > 0$ be the constant in the estimates of Lemma 3.11 and choose n_0 large enough that

$$(17) \quad \rho := C_0 2^{\frac{1}{\gamma}+2} \gamma (n_0 - 1)^{-\frac{1}{\gamma}} < 1.$$

We call a return to J_h *long* if there have been at least n_0 iterates since the last entry to J_h (or for the first entry, if it occurs after time $n_0 - 1$). A return that is not long is called *short*.

Given $x \in \dot{I}^{t-1} \setminus \dot{I}^t$, consider the decomposition of $[0, t]$ into segments, labelled alternately $s_1, \ell_1, s_2, \ell_2, \dots, s_k, \ell_k, s_{k+1}$, where the segments ℓ_i are long returns to J_h and the segments s_i are comprised of one or more short returns to J_h (but the total length s_i of the segment may be greater than n_0). Note we must have $\ell_i \geq n_0$ for each i but some of the s_i may be 0.

Let $(s_1, \dots, s_{k+1}; \ell_1, \dots, \ell_k; t) \subset \dot{I}^{t-1} \setminus \dot{I}^t$ denote the set of points with the specified trajectory which fall into the hole at time t with k long returns to J_h . Let also $\ell = \sum_{i=1}^k \ell_i$.

Recall once again the induced map S and the associated sets \dot{I}_S^t , which we now consider for $n_S = n_0 + h$. Lemma 3.4 implies the existence of $C_3 > 0$ and $\sigma_0 < 1$ such that $m(\dot{I}_S^t) \leq C_3 \sigma_0^t$ for all t . Now choose $C_4 > 0$ sufficiently large such that

$$C_4 \log \sigma_0^{-1} \geq \frac{\gamma+1}{\gamma}.$$

We divide our estimate into two cases.

Case 1 ($t - \ell > C_4 \log t$). During short returns, iterates remain in the interval $[a_{n_0+h}, 1)$. Therefore, the orbits of every point in $(s_1, \dots, s_{k+1}; \ell_1, \dots, \ell_k; t)$ spend at least $t - \ell$ iterates inside $[a_{n_0+h}, 1)$, i.e. we have $(s_1, \dots, s_{k+1}; \ell_1, \dots, \ell_k; t) \subset \dot{I}_S^{t-\ell}$, which implies

$$m(x \in \dot{I}^{t-1} \setminus \dot{I}^t : t - \ell > C_4 \log t) \leq C_3 \sigma_0^{C_4 \log t} \leq C_3 t^{-\frac{\gamma+1}{\gamma}}.$$

Case 2 ($t - \ell \leq C_4 \log t$). We fix $k \geq 1$ and estimate the contribution to $\hat{I}^{t-1} \setminus \hat{I}^t$ from points making k long returns to J_h . Now

$$m(s_1, \dots, s_{k+1}; \ell_1, \dots, \ell_k; t) = m(s_{k+1} \mid (s_1, \dots, s_k; \ell_1, \dots, \ell_k)) \\ \cdot \prod_{i=1}^k m(\ell_i \mid (s_1, \dots, s_i; \ell_1, \dots, \ell_{i-1})) m(s_i \mid (s_1, \dots, s_{i-1}; \ell_1, \dots, \ell_{i-1})),$$

where the truncated lists $(s_1, \dots, s_i; \ell_1, \dots, \ell_{i-1})$ denote the set of points whose entries to J_h follow the specified itinerary. Note that by Lemma 3.11 we have

$$m(\ell_i \mid (s_1, \dots, s_i; \ell_1, \dots, \ell_{i-1})) \leq C_0 \ell_i^{-\frac{\gamma+1}{\gamma}}.$$

Also, we have

$$\sum_{\text{relevant } s_i} m(s_i \mid (s_1, \dots, s_{i-1}; \ell_1, \dots, \ell_{i-1})) \leq 1$$

for each i . Since each sum over s_i holds for s_1, \dots, s_{i-1} fixed, we apply the bound above recursively to obtain

$$\sum_{s_1, \dots, s_{k+1}} \prod_{i=1}^{k+1} m(s_i \mid (s_1, \dots, s_{i-1}; \ell_1, \dots, \ell_{i-1})) \leq 1,$$

where the sum runs over relevant s_1, \dots, s_{k+1} . Finally, summing the products over the relevant lengths ℓ_i , we must estimate the following expression, where as before $\beta = \frac{\gamma+1}{\gamma}$:

$$(18) \quad \sum_{\ell=\lceil t-C_4 \log t \rceil}^t \sum_{\ell_1=n_0}^{\ell-(k-1)n_0} \sum_{\ell_2=n_0}^{\ell-\ell_1-(k-2)n_0} \dots \\ \sum_{\ell_{k-1}=n_0}^{\ell-\ell_1-\dots-\ell_{k-2}-n_0} C_0^k (\ell_1 \dots \ell_{k-1})^{-\beta} (\ell - \ell_1 - \dots - \ell_{k-1})^{-\beta},$$

where we have used the fact that $\ell_k = \ell - \ell_1 - \dots - \ell_{k-1}$. We will estimate the iterated sums one at a time and show the calculation in detail in order to verify that we can control the effect of the constant C_0^k .

Let $s = \ell - \ell_1 - \dots - \ell_{k-2}$. Then the first sum we must estimate is

$$(19) \quad \sum_{\ell_{k-1}=n_0}^{s-n_0} (s - \ell_{k-1})^{-\beta} \ell_{k-1}^{-\beta} \leq 2 \sum_{\ell_{k-1}=n_0}^{\lfloor s/2 \rfloor} (s - \ell_{k-1})^{-\beta} \ell_{k-1}^{-\beta} \leq 2^{1+\beta} s^{-\beta} \sum_{\ell_{k-1}=n_0}^{\lfloor s/2 \rfloor} \ell_{k-1}^{-\beta} \\ \leq 2^{1+\beta} s^{-\beta} \int_{n_0-1}^{\infty} x^{-\beta} dx \leq 2^{1+\beta} s^{-\beta} \frac{(n_0-1)^{1-\beta}}{\beta-1} \\ = 2^{\frac{1}{\gamma}+2} s^{-\beta} \gamma (n_0-1)^{-\frac{1}{\gamma}} \leq C_0^{-1} \rho s^{-\beta},$$

where the last inequality follows from equation (17). Turning now to the second sum in (18), we must estimate

$$\sum_{\ell_{k-2}=n_0}^{\ell-\ell_1-\dots-\ell_{k-3}-2n_0} (\ell - \ell_1 - \dots - \ell_{k-2})^{-\beta} \ell_{k-2}^{-\beta},$$

and setting $s = \ell - \ell_1 - \dots - \ell_{k-3}$, we see that this sum has the same form as that in (19) and thus satisfies the same bound (note, we are not hurt by the fact that we are subtracting $2n_0$ rather than n_0 in the upper limit of this sum since subtracting $2n_0$ just makes the sum smaller). Proceeding this way $k - 3$ more times, we arrive at the expression

$$\begin{aligned} m(x \in \overset{\circ}{I}^{t-1} \setminus \overset{\circ}{I}^t : k \text{ long passes with } t - \ell \leq C_4 \log t) &\leq \sum_{\ell=\lceil t-C_4 \log t \rceil}^t C_0 \rho^{k-1} \ell^{-\beta} \\ &\leq C_0 \rho^{k-1} \lfloor C_4 \log t + 1 \rfloor \lceil t - C_4 \log t \rceil^{-\frac{\gamma+1}{\gamma}}. \end{aligned}$$

Finally, summing over k yields the existence of $C > 0$ such that

$$m(\overset{\circ}{I}^{t-1} \setminus \overset{\circ}{I}^t : t - \ell \leq C_4 \log t) \leq C t^{-\frac{\gamma+1}{\gamma}} \log t.$$

Putting together our estimates from Cases 1 and 2 completes the proof of the lemma. \square

3.6. General open systems - proof of Theorem 2.3. The proof requires a preliminary statement. We say a set $A \subset \mathbb{N}$ has density $\rho \in [0, 1]$ if $\limsup_{n \rightarrow +\infty} \frac{\#(A \cap [1, n])}{n} = \rho$, where $\#$ denotes the cardinality of a set.

Define $\beta_t = \frac{\mu(\overset{\circ}{X}^{t+1})}{\mu(\overset{\circ}{X}^t)}$ for $t \geq 0$ and $\beta_{-1} = \mu(\overset{\circ}{X}^0)$. We obviously have $\beta_t \leq 1$ for all $t \geq -1$, and the desired preliminary statement specifies the density of any set of indexes t in which β_t remains bounded away from 1.

Lemma 3.12. *For $\lambda \in (0, 1)$, define $A_\lambda = \{k \in \mathbb{N} : \beta_k \leq \lambda\}$. Then A_λ must have zero density.*

Proof. Fix $\lambda \in (0, 1)$ and note that if A_λ is finite, it has zero density. So assume A_λ is infinite. Then $\limsup_{\substack{k \rightarrow +\infty \\ k \in A_\lambda}} \beta_k \leq \lambda$. Equation (2) implies

$$(20) \quad \lim_{t \rightarrow +\infty} \mu(\overset{\circ}{X}^t)^{\frac{1}{t}} = \lim_{t \rightarrow +\infty} \exp\left(\frac{1}{t} \log \mu(\overset{\circ}{X}^t)\right) = 1.$$

To derive a contradiction, assume A_λ has density $\rho > 0$. The relation $\mu(\overset{\circ}{X}^t) = \prod_{k=-1}^{t-1} \beta_k$ then implies

$$\mu(\overset{\circ}{X}^t)^{\frac{1}{t}} \leq \lambda^{\frac{\#(A_\lambda \cap [0, t-1])}{t}} \text{ for all } t > 0,$$

from which it follows that $\liminf_{t \rightarrow \infty} \mu(\overset{\circ}{X}^t)^{\frac{1}{t}} \leq \lambda^\rho < 1$, contradicting equation (20). \square

Passing now to the proof of the theorem, we consider the sequence of probability measures $\{\mu_t\}_{t \in \mathbb{N}}$ where μ_t is defined by equation (3). By assumption on X , this sequence is compact and any limit point μ_∞ is necessarily a (regular Borel) probability measure.

Our proof has two steps: (A) μ_∞ gives zero weight to $\text{int}\left(\bigcup_{j=0}^i T^{-j}(H)\right)$ for every $i \geq 0$; (B) if μ_∞ gives zero weight to the discontinuity set of $\overset{\circ}{T}$, then μ_∞ is an invariant measure for T .

Since T is assumed to be non-singular with respect to μ , the associated transfer operator \mathcal{L} acting on $L^1(\mu)$ is well-defined. As before, we also consider the transfer operator $\mathring{\mathcal{L}}$ for the system with hole defined by $\mathring{\mathcal{L}}^t f = \mathcal{L}^t(f1_{\dot{X}^t})$ for all $t \geq 1$.

(A) First note that $\mathring{T}_*^t \mu$ is supported on $X \setminus H$ for each $t \geq 0$ so that $\mu_\infty(H) = 0$. Now define $g_k = \frac{\mathring{\mathcal{L}}^k 1}{\mu(\dot{X}^k)}$ for $k \geq 1$. For each $i \geq 1$, $k \geq 1$, we have (using the composition property pointed out at the end of Section 1.1)

$$\beta_{k+i-1} \cdots \beta_k = \frac{\int_X \mathring{\mathcal{L}}^i(\mathring{\mathcal{L}}^k 1) d\mu}{\mu(\dot{X}^k)} = \int_{\dot{X}^i} g_k d\mu = 1 - \int_{\bigcup_{j=0}^i T^{-j}(H)} g_k d\mu,$$

since $\int_{X \setminus H} g_k = 1$. Choose $\lambda \in (0, 1)$ and let A_λ be as defined in Lemma 3.12. Given $i \geq 1$, let

$$(21) \quad A_\lambda^i = A_\lambda - [0, i-1]$$

denote the translates of elements of A_λ by some integer at most $i-1$. Note that A_λ^i still has frequency zero. Then since $\liminf_{k \rightarrow +\infty} \beta_{k+i-1} \cdots \beta_k \geq \lambda^i$, as long as we avoid the exceptional set A_λ^i , we have

$$\limsup_{\substack{k \rightarrow +\infty \\ k \notin A_\lambda^i}} \int_{\bigcup_{j=0}^i T^{-j}(H)} g_k d\mu \leq 1 - \lambda^i.$$

Since $d\mu_{t_j} = \frac{1}{t_j} \sum_{k=0}^{t_j-1} g_k d\mu$ and A_λ^i has zero density, we conclude $\mu_\infty\left(\text{int}\left(\bigcup_{j=0}^i T^{-j}(H)\right)\right) \leq 1 - \lambda^i$. Since λ is arbitrary, it must be that $\mu_\infty\left(\text{int}\left(\bigcup_{j=0}^i T^{-j}(H)\right)\right) = 0$. Since i is arbitrary, μ_∞ must be supported on $X \setminus \left(\bigcup_{i=0}^\infty \text{int}\left(\bigcup_{j=0}^i T^{-j}(H)\right)\right)$ and therefore is singular with respect to μ by assumption (2).

(B) For any test function $\varphi \in C^0(X)$, we have

$$\begin{aligned} \mathring{T}_* \mu_{t_j}(\varphi) &= \frac{1}{t_j} \sum_{k=0}^{t_j-1} \frac{\int \mathring{\mathcal{L}}^{k+1} 1 \varphi d\mu}{\mu(\dot{X}^k)} = \frac{1}{t_j} \sum_{k=0}^{t_j-1} \frac{\int \mathring{\mathcal{L}}^{k+1} 1 \varphi d\mu}{\mu(\dot{X}^{k+1})} \beta_k \\ &= \frac{1}{t_j} \sum_{k=0}^{t_j-1} \frac{\int \mathring{\mathcal{L}}^k 1 \varphi d\mu}{\mu(\dot{X}^k)} \beta_{k-1} + \frac{1}{t_j} \left(\frac{\int \mathring{\mathcal{L}}^{t_j} 1 \varphi d\mu}{\mu(\dot{X}^{t_j-1})} - \int_{\dot{X}^0} \varphi d\mu \right). \end{aligned}$$

Again choose $\lambda \in (0, 1)$, define A_λ as in Lemma 3.12 and A_λ^1 as in (21). The discontinuity set of $\varphi \circ \mathring{T}$ is contained in the discontinuity set of \mathring{T} , and since we assume that μ_∞ gives zero measure to this set, we may pass to the limit (see

e.g. [2, Theorem 5.2(iii)]) to obtain

$$\begin{aligned} |\hat{T}_* \mu_\infty(\varphi) - \mu_\infty(\varphi)| &\leq \lim_{j \rightarrow +\infty} \frac{1}{t_j} \sum_{\substack{k=0 \\ k \in A_\lambda^1}}^{t_j-1} \frac{|\int \hat{\mathcal{L}}^k 1 \varphi d\mu|}{|\hat{\mathcal{L}}^k 1|_1} |\beta_{k-1} - 1| \\ &\quad + \frac{1}{t_j} \left(\frac{|\varphi|_\infty |\hat{\mathcal{L}}^{t_j} 1|_1}{|\hat{\mathcal{L}}^{t_j-1} 1|_1} + |\varphi|_\infty \right) \\ &\quad + \lim_{j \rightarrow +\infty} \frac{1}{t_j} \sum_{\substack{k=0 \\ k \notin A_\lambda^1}}^{t_j-1} \frac{|\int \hat{\mathcal{L}}^k 1 \varphi d\mu|}{|\hat{\mathcal{L}}^k 1|_1} |\beta_{k-1} - 1| \\ &\leq |\varphi|_\infty (1 - \lambda), \end{aligned}$$

where we have used the fact that the set of k for which $\beta_{k-1} \leq \lambda$ has density 0, $\beta_k \leq 1$, and $\frac{|\int \hat{\mathcal{L}}^k 1 \varphi d\mu|}{|\hat{\mathcal{L}}^k 1|_1} \leq |\varphi|_\infty$ for each k . Since λ is arbitrary, we must have $\hat{T}_* \mu_\infty(\varphi) = \mu_\infty(\varphi)$.

Furthermore, since $\partial H \cup \hat{T}^{-1}(\partial H)$ is contained in the discontinuity set for \hat{T} , by assumption we have $\mu_\infty(\partial H \cup \hat{T}^{-1}(\partial H)) = 0$, and this together with part (A) implies $\mu_\infty(H \cup T^{-1}(H)) = 0$, i.e. $\mu_\infty(\hat{X}^1) = 1$. Thus for $\varphi \in C^0(X)$,

$$\mu_\infty(\varphi) = \hat{T}_* \mu_\infty(\varphi) = \mu_\infty(\varphi \circ T \cdot 1_{\hat{X}^1}) = \mu_\infty(\varphi \circ T) = T_* \mu_\infty(\varphi).$$

Since this identity holds for all $\varphi \in C^0(X)$ and μ_∞ is a regular Borel measure, it follows from the uniqueness statement of the Riesz-Markov Theorem [35, Theorem 13.23] that $T_* \mu_\infty = \mu_\infty$ so that μ_∞ is an invariant measure for T . Now the invariance of μ_∞ implies $\mu_\infty(T^{-j}(H)) = 0$ for each $j \in \mathbb{N}$ so that $\mu_\infty(\hat{X}^\infty) = 1$.

ACKNOWLEDGEMENTS

The authors are grateful to L.-S. Young for stimulating discussions. The second author thanks the Courant Institute, where part of this work was done, for its hospitality.

REFERENCES

- [1] V. S. Afraimovich and L. A. Bunimovich, *Which hole is leaking the most: a topological approach to study open systems*, Nonlinearity **23** (2010), no. 3, 643–656, DOI 10.1088/0951-7715/23/3/012. MR2593912 (2011e:37080)
- [2] Patrick Billingsley, *Convergence of probability measures*, John Wiley & Sons, Inc., New York-London-Sydney, 1968. MR0233396 (38 #1718)
- [3] Henk Bruin, Mark Demers, and Ian Melbourne, *Existence and convergence properties of physical measures for certain dynamical systems with holes*, Ergodic Theory Dynam. Systems **30** (2010), no. 3, 687–728, DOI 10.1017/S0143385709000200. MR2643708 (2011g:37107)
- [4] L.A. Bunimovich and C.P. Dettmann, *Open circular billiards and the Riemann hypothesis*, Phys. Rev. Lett. **94** (2005), 100201.
- [5] L. A. Bunimovich and C. P. Dettmann, *Peeping at chaos: nondestructive monitoring of chaotic systems by measuring long-time escape rates*, Europhys. Lett. EPL **80** (2007), no. 4, Art. 40001, 6, DOI 10.1209/0295-5075/80/40001. MR2354135 (2008g:82066)
- [6] Leonid A. Bunimovich and Alex Yurchenko, *Where to place a hole to achieve a maximal escape rate*, Israel J. Math. **182** (2011), 229–252, DOI 10.1007/s11856-011-0030-8. MR2783972 (2012e:37012)
- [7] N. N. Čencova, *Natural invariant measure for the Smale horseshoe* (Russian), Dokl. Akad. Nauk SSSR **256** (1981), no. 2, 294–298. MR600950 (82d:58033)

- [8] N. Chernov and R. Markarian, *Ergodic properties of Anosov maps with rectangular holes*, Bol. Soc. Brasil. Mat. (N.S.) **28** (1997), no. 2, 271–314, DOI 10.1007/BF01233395. MR1479505 (98k:58140)
- [9] N. Chernov and R. Markarian, *Anosov maps with rectangular holes. Nonergodic cases*, Bol. Soc. Brasil. Mat. (N.S.) **28** (1997), no. 2, 315–342, DOI 10.1007/BF01233396. MR1479506 (99b:58142)
- [10] N. Chernov, R. Markarian, and S. Troubetzkoy, *Conditionally invariant measures for Anosov maps with small holes*, Ergodic Theory Dynam. Systems **18** (1998), no. 5, 1049–1073, DOI 10.1017/S0143385798117492. MR1653291 (99k:58111)
- [11] N. Chernov, R. Markarian, and S. Troubetzkoy, *Invariant measures for Anosov maps with small holes*, Ergodic Theory Dynam. Systems **20** (2000), no. 4, 1007–1044, DOI 10.1017/S0143385700000560. MR1779391 (2001f:37042)
- [12] H. van den Bedem and N. Chernov, *Expanding maps of an interval with holes*, Ergodic Theory Dynam. Systems **22** (2002), no. 3, 637–654, DOI 10.1017/S0143385702000329. MR1908547 (2003c:37046)
- [13] Pierre Collet, Servet Martínez, and Véronique Maume-Deschamps, *On the existence of conditionally invariant probability measures in dynamical systems*, Nonlinearity **13** (2000), no. 4, 1263–1274, DOI 10.1088/0951-7715/13/4/315. MR1767958 (2001e:37007)
- [14] P. Collet, S. Martínez, and V. Maume-Deschamps, *Corrigendum: “On the existence of conditionally invariant probability measures in dynamical systems”* [Nonlinearity **13** (2000), no. 4, 1263–1274; MR1767958], Nonlinearity **17** (2004), no. 5, 1985–1987, DOI 10.1088/0951-7715/17/5/C01. MR2086159 (2005e:37002)
- [15] Pierre Collet, Servet Martínez, and Bernard Schmitt, *The Yorke-Pianigiani measure and the asymptotic law on the limit Cantor set of expanding systems*, Nonlinearity **7** (1994), no. 5, 1437–1443. MR1294552 (95i:28011)
- [16] Mark F. Demers, *Markov extensions for dynamical systems with holes: an application to expanding maps of the interval*, Israel J. Math. **146** (2005), 189–221, DOI 10.1007/BF02773533. MR2151600 (2006d:37062)
- [17] Mark F. Demers, *Markov extensions and conditionally invariant measures for certain logistic maps with small holes*, Ergodic Theory Dynam. Systems **25** (2005), no. 4, 1139–1171, DOI 10.1017/S0143385704000963. MR2158400 (2007d:37057)
- [18] Mark F. Demers and Lai-Sang Young, *Escape rates and conditionally invariant measures*, Nonlinearity **19** (2006), no. 2, 377–397, DOI 10.1088/0951-7715/19/2/008. MR2199394 (2006i:37051)
- [19] Mark F. Demers and Carlangelo Liverani, *Stability of statistical properties in two-dimensional piecewise hyperbolic maps*, Trans. Amer. Math. Soc. **360** (2008), no. 9, 4777–4814, DOI 10.1090/S0002-9947-08-04464-4. MR2403704 (2009f:37021)
- [20] Mark Demers, Paul Wright, and Lai-Sang Young, *Escape rates and physically relevant measures for billiards with small holes*, Comm. Math. Phys. **294** (2010), no. 2, 353–388, DOI 10.1007/s00220-009-0941-y. MR2579459 (2011a:37077)
- [21] Mark F. Demers and Paul Wright, *Behaviour of the escape rate function in hyperbolic dynamical systems*, Nonlinearity **25** (2012), no. 7, 2133–2150, DOI 10.1088/0951-7715/25/7/2133. MR2947939
- [22] Mark F. Demers, *Dispersing billiards with small holes*, Ergodic theory, open dynamics, and coherent structures, Springer Proc. Math. Stat., vol. 70, Springer, New York, 2014, pp. 137–170, DOI 10.1007/978-1-4939-0419-8_8. MR3213499
- [23] Mark F. Demers, *Escape rates and physical measures for the infinite horizon Lorentz gas with holes*, Dyn. Syst. **28** (2013), no. 3, 393–422, DOI 10.1080/14689367.2013.814946. MR3170623
- [24] Carl P. Dettmann and Orestis Georgiou, *Survival probability for the stadium billiard*, Phys. D **238** (2009), no. 23–24, 2395–2403, DOI 10.1016/j.physd.2009.09.019. MR2576082 (2011a:37078)
- [25] C. P. Dettmann and E. D. Leonel, *Escape and transport for an open bouncer: Stretched exponential decays*, Physica D **241** (2012), 403–408.
- [26] P. A. Ferrari, H. Kesten, S. Martinez, and P. Picco, *Existence of quasi-stationary distributions. A renewal dynamical approach*, Ann. Probab. **23** (1995), no. 2, 501–521. MR1334159 (96c:60089)

- [27] Gary Froyland, Rua Murray, and Ognjen Stancevic, *Spectral degeneracy and escape dynamics for intermittent maps with a hole*, *Nonlinearity* **24** (2011), no. 9, 2435–2463, DOI 10.1088/0951-7715/24/9/003. MR2819930 (2012e:37042)
- [28] Ale Jan Homburg and Todd Young, *Intermittency in families of unimodal maps*, *Ergodic Theory Dynam. Systems* **22** (2002), no. 1, 203–225, DOI 10.1017/S0143385702000093. MR1889571 (2003h:37050)
- [29] Gerhard Keller and Carlangelo Liverani, *Rare events, escape rates and quasistationarity: some exact formulae*, *J. Stat. Phys.* **135** (2009), no. 3, 519–534, DOI 10.1007/s10955-009-9747-8. MR2535206 (2011a:37012)
- [30] Carlangelo Liverani and Véronique Maume-Deschamps, *Lasota-Yorke maps with holes: conditionally invariant probability measures and invariant probability measures on the survivor set* (English, with English and French summaries), *Ann. Inst. H. Poincaré Probab. Statist.* **39** (2003), no. 3, 385–412, DOI 10.1016/S0246-0203(02)00005-5. MR1978986 (2004c:37009)
- [31] Carlangelo Liverani, Benoît Saussol, and Sandro Vaienti, *A probabilistic approach to intermittency*, *Ergodic Theory Dynam. Systems* **19** (1999), no. 3, 671–685, DOI 10.1017/S0143385799133856. MR1695915 (2000d:37029)
- [32] Artur Lopes and Roberto Markarian, *Open billiards: invariant and conditionally invariant probabilities on Cantor sets*, *SIAM J. Appl. Math.* **56** (1996), no. 2, 651–680, DOI 10.1137/S0036139995279433. MR1381665 (97a:58108)
- [33] Giulio Pianigiani and James A. Yorke, *Expanding maps on sets which are almost invariant. Decay and chaos*, *Trans. Amer. Math. Soc.* **252** (1979), 351–366, DOI 10.2307/1998093. MR534126 (80i:58030)
- [34] P. A. Richardson, Jr., *Natural measures on the unstable and invariant manifolds of open billiard dynamical systems*, *Doctoral Dissertation*, Department of Mathematics, University of North Texas (1999).
- [35] H. L. Royden, *Real analysis*, 3rd ed., Macmillan Publishing Company, New York, 1988. MR1013117 (90g:00004)
- [36] E. Seneta and D. Vere-Jones, *On quasi-stationary distributions in discrete-time Markov chains with a denumerable infinity of states*, *J. Appl. Probability* **3** (1966), 403–434. MR0207047 (34 #6863)
- [37] D. Vere-Jones, *Geometric ergodicity in denumerable Markov chains*, *Quart. J. Math. Oxford Ser. (2)* **13** (1962), 7–28. MR0141160 (25 #4571)
- [38] A. M. Yaglom, *Certain limit theorems of the theory of branching random processes* (Russian), *Doklady Akad. Nauk SSSR (N.S.)* **56** (1947), 795–798. MR0022045 (9,149e)
- [39] Lai-Sang Young, *Recurrence times and rates of mixing*, *Israel J. Math.* **110** (1999), 153–188, DOI 10.1007/BF02808180. MR1750438 (2001j:37062)

DEPARTMENT OF MATHEMATICS, FAIRFIELD UNIVERSITY, FAIRFIELD, CONNECTICUT 06824

CENTRE DE PHYSIQUE THÉORIQUE, CNRS - AIX-MARSEILLE UNIVERSITÉ - UNIVERSITÉ DE TOULON, CAMPUS DE LUMINY, 13288 MARSEILLE CEDEX 9, FRANCE